



Research article

Analytical and numerical negative boundedness of fractional differences with Mittag–Leffler kernel

Pshtiwan Othman Mohammed^{1*}, Rajendra Dahal², Christopher S. Goodrich³, Y. S. Hamed⁴ and Dumitru Baleanu^{5,6,7,*}

- ¹ Department of Mathematics, College of Education, University of Sulaimani, Sulaimani 46001, Iraq
² Department of Mathematics and Statistics, Coastal Carolina University, Conway, SC 29526, United States of America
³ School of Mathematics and Statistics, UNSW Sydney, Sydney, NSW 2052, Australia
⁴ Department of Mathematics and Statistics, College of Science, Taif University, P.O. Box 11099, Taif 21944, Saudi Arabia
⁵ Department of Mathematics, Cankaya University, 06530 Balgat, Ankara, Turkey
⁶ Institute of Space Sciences, R76900 Magurele-Bucharest, Romania
⁷ Department of Natural Sciences, School of Arts and Sciences, Lebanese American University, Beirut 11022801, Lebanon

* **Correspondence:** Email: pshtiwansangawi@gmail.com; dumitru@cankaya.edu.tr.

Abstract: We show that a class of fractional differences with Mittag–Leffler kernel can be negative and yet monotonicity information can still be deduced. Our results are complemented by numerical approximations. This adds to a growing body of literature illustrating that the sign of a fractional difference has a very complicated and subtle relationship to the underlying behavior of the function on which the fractional difference acts, regardless of the particular kernel used.

Keywords: discrete fractional calculus; Mittag–Leffler type kernel; analytical and numerical monotonicity results

Mathematics Subject Classification: 26A33, 26A51, 33B10, 39A12, 39B62, 65D15, 65Q20

1. Introduction

Define the sets \mathbb{N}_a and \mathbb{N}_a^b , for any $a, b \in \mathbb{R}$ with $b - a$ a nonnegative integer, to be $\mathbb{N}_a := \{a, a + 1, a + 2, \dots\}$ and $\mathbb{N}_a^b := \{a, a + 1, a + 2, \dots, b\}$, respectively. It worth recalling that the nabla (or

backward) difference of a function $g : \mathbb{N}_a \rightarrow \mathbb{R}$ can be expressed as follows

$$(\nabla g)(t) := g(t) - g(t-1), \quad \text{for } t \in \mathbb{N}_{a+1}.$$

As is known, there is a strong correlation between the sign of the nabla operator and whether g is either monotone increasing or monotone decreasing. For example, if $(\nabla g)(t) \geq 0$, then g is increasing function on \mathbb{N}_a .

In recent years a nonlocal version of the discrete calculus has been proposed. This nonlocal version is known as the “discrete fractional calculus”, a research area popularized by the seminal papers of Atici and Eloe [9–11] in the late 2000s and then further extended by the subsequent work of Lizama [25]. One of the reasons for the interest in the discrete fractional calculus is its emerging applications in biological mathematics – see, for example, the recent work of Atici, et al. [7, 8], in which the authors apply discrete fractional calculus to the modeling of tumors.

To understand the nonlocal nature of the fractional calculus, consider a commonly utilized version of the discrete fractional difference – namely, for $\nu > 0$ the ν -th order Riemann-Liouville difference, which for a function $g : \mathbb{N}_a \rightarrow \mathbb{R}$ is denoted by ${}^{\text{RL}}\Delta^\nu g$ and defined pointwise by

$$({}^{\text{RL}}\Delta^\nu g)(t) := \sum_{s=a}^{t+\nu} \frac{\Gamma(t-s)}{\Gamma(-\nu)\Gamma(t-s+\nu+1)} g(s), \quad t \in \mathbb{N}_{a+N-\nu}, \quad (1.1)$$

where a positive integer N satisfying $N-1 < \nu \leq N$. The key fact about (1.1) is that it is nonlocal, very much unlike the integer-order difference mentioned in the first paragraph of this section. Consequently, the relationships between the sign of $({}^{\text{RL}}\Delta^\nu g)(t)$ and the monotone behavior of g are quite muddled and complex. This question was initially investigated by Dahal and Goodrich [13] in 2014, and then subsequently investigated by many authors including Abdeljawad and Baleanu [1], Bravo, Lizama, and Rueda [12], Goodrich and Jonnalagadda [16], Goodrich and Lizama [18, 19], Goodrich and Muellner [22], and Jia, Erbe, and Peterson [15, 24].

Very recently Goodrich, Lyons, and Velcsov [20] together with Jonnalagadda [17] and Scapellato [21] demonstrated that a function can increase (under certain conditions) even if its fractional difference is negative. This is something that plainly cannot happen in the integer-order case. And this phenomenon provides further evidence of the highly complicating nature of the nonlocal structure of fractional-order difference operators. It also has serious implications for the use of fractional calculus in modeling since one of the most important uses of calculus in modeling is to identify where functions are increasing or decreasing.

So, all in all, there is a large body of evidence that nonlocal discrete operators behave in ways that are very complicated, particularly as concerns their ability to detect the qualitative behaviors of the functions on which they operate. At the same time, there are a variety of definitions for discrete fractional differences and sums. Therefore, it is relevant to determine whether these aberrant behaviors are exhibited by all such nonlocal difference operators – or only some of them.

Consequently, in this brief note, we propose to continue investigating these questions in the specific context of the fractional difference with exponential-type kernels (see Definitions 2.1 and 2.2). In particular, we demonstrate that, as with other definitions of the fractional difference, the type of difference studied here can be negative even though the function on which it acts is increasing, and this observation confirms that fractional differences with Mittag-Leffler kernels exhibit the same

sort of aberrant behavior as other nonlocal difference operators. This complements not only the already mentioned reference [17] but also builds upon earlier work by Abdeljawad, Al-Mdallal, and Hajji [6]. And it continues to demonstrate some of the surprising properties of this class of discrete nonlocal operators. To see the development of fractional calculus with Mittag-Leffler kernels we advise the readers to see the recently published articles [28, 29].

2. Essential tools and main results

In this section, we begin with recalling the necessary fundamental discrete operators for our main results briefly. The interested reader may visit the monograph [23] by Goodrich and Peterson and recently published articles [1–3, 26] for additional information and mathematical background regarding the discrete fractional calculus.

The first and well-known definition in this article is the discrete Mittag-Leffler functions. We then provide the definition of the discrete fractional difference defined using the Mittag-Leffler kernel on the set \mathbb{N}_a . The discrete Mittag-Leffler function of 2-parameters is given by (see [27]):

$$E_{\nu, \beta}(\lambda, t) := \sum_{k=0}^{\infty} \lambda^k \frac{t^{\overline{k\nu + \beta - 1}}}{\Gamma(k\nu + \beta)},$$

for $\lambda \in \mathbb{R}$ such that $1 > |\lambda|$, and $\nu, \beta, t \in \mathbb{C}$ such that $\operatorname{Re}(\nu) > 0$. It is essential to see that $t^{\overline{\nu}}$ is the rising function and given as follows

$$t^{\overline{\nu}} := \frac{\Gamma(t + \nu)}{\Gamma(t)},$$

for $\nu \in \mathbb{R}$ and $t \in \mathbb{R}$ apart from the elements $\{\dots, -2, -1, 0\}$. As a special case of the above definition, the discrete Mittag-Leffler function of 1-parameter is given as follows

$$E_{\nu}(\lambda, t) := \sum_{k=0}^{\infty} \lambda^k \frac{t^{\overline{k\nu}}}{\Gamma(k\nu + 1)} \quad (\text{for } |\lambda| < 1).$$

Remark 2.1. Considering Remark 1 in [4], we can obtain the following for $\lambda_1 = -\frac{\nu-1}{2-\nu}$ and $1 < \nu < \frac{3}{2}$:

- $E_{\nu-1}(\lambda_1, 0) = 1$,
- $E_{\nu-1}(\lambda_1, 1) = 2 - \nu$,
- $E_{\nu-1}(\lambda_1, 2) = \nu(2 - \nu)^2$,
- $E_{\nu-1}(\lambda_1, 3) = \frac{2-\nu}{2} \left[(\nu-1)^3(2\nu-3) - 3(\nu-1)^2 + 2 \right]$,
- $0 < E_{\nu-1}(\lambda_1, t) < 1$ for each $1 < \nu < \frac{3}{2}$ and $t = 1, 2, 3, \dots$. At the same time, we have that $E_{\nu-1}(\lambda_1, t)$ is monotonically decreasing for each $1 < \nu < \frac{3}{2}$ and $t = 0, 1, 2, \dots$.

Definition 2.1. (see [1, Definition 2.24]) Let $0 < \nu < \frac{1}{2}$ and $\lambda_0 = -\frac{\nu}{1-\nu}$. Then, the discrete fractional difference operators with Mittag-Leffler kernels of order ν denoted by ${}^{\text{ABC}}_a \nabla^{\nu} g$ and ${}^{\text{ABR}}_a \nabla^{\nu} g$, respectively,

defined by

$$\left({}^{\text{ABC}}_a \nabla^\nu g\right)(t) := \frac{H(\nu)}{1-\nu} \nabla_t \sum_{r=a+1}^t g(r) E_{\bar{\nu}}(\lambda_0, t-r+1),$$

and

$$\left({}^{\text{ABR}}_a \nabla^\nu g\right)(t) := \frac{H(\nu)}{1-\nu} \sum_{r=a+1}^t (\nabla_r g)(r) E_{\bar{\nu}}(\lambda_0, t-r+1),$$

for each $t \in \mathbb{N}_{a+1}$. Here the function $\nu \mapsto H(\nu)$ is a normalization constant satisfying $0 < H(\nu)$.

Definition 2.2. (see [5]) For $g : \mathbb{N}_{a-\aleph} \rightarrow \mathbb{R}$ with $\aleph < \nu \leq \aleph + \frac{1}{2}$ and $\aleph \in \mathbb{N}_0$, Then, the discrete fractional difference operators with Mittag-Leffler kernels of order ν , respectively, are defined by

$$\left({}^{\text{ABC}}_a \nabla^\nu g\right)(t) = \left({}^{\text{ABC}}_a \nabla^{\nu-\aleph} \nabla^\aleph g\right)(t) := \frac{H(\nu-\aleph)}{\aleph+1-\nu} \nabla_t \sum_{r=a+1}^t \left(\nabla_r^\aleph g\right)(r) E_{\bar{\nu-\aleph}}(\lambda_\aleph, t-r+1),$$

and

$$\left({}^{\text{ABR}}_a \nabla^\nu g\right)(t) = \left({}^{\text{ABR}}_a \nabla^{\nu-\aleph} \nabla^\aleph g\right)(t) := \frac{H(\nu-\aleph)}{\aleph+1-\nu} \sum_{r=a+1}^t \left(\nabla_r^{\aleph+1} g\right)(r) E_{\bar{\nu-\aleph}}(\lambda_\aleph, t-r+1),$$

for $t \in \mathbb{N}_{a+1}$. Here $\lambda_\aleph = -\frac{\nu-\aleph}{\aleph+1-\nu}$.

The following is the essential lemma which brings us to the main results.

Lemma 2.1. Let the function g be defined on \mathbb{N}_a and $1 < \nu < \frac{3}{2}$. Then we have that

$$\begin{aligned} \left({}^{\text{ABR}}_a \nabla^\nu g\right)(t) &= H(\nu-1) \left\{ (\nabla g)(t) + \frac{1}{2-\nu} \left[E_{\bar{\nu-1}}(\lambda_1, t-a) - E_{\bar{\nu-1}}(\lambda_1, t-a-1) \right] (\nabla g)(a+1) \right. \\ &\quad \left. + \frac{1}{2-\nu} \sum_{r=a+2}^{t-1} \left[E_{\bar{\nu-1}}(\lambda_1, t-r+1) - E_{\bar{\nu-1}}(\lambda_1, t-r) \right] (\nabla_r g)(r) \right\}, \end{aligned}$$

for each $t \in \mathbb{N}_{a+3}$.

Proof. From Definitions 2.1 and 2.2, the following can be deduced for $1 < \nu < \frac{3}{2}$:

$$\begin{aligned} \left({}^{\text{ABR}}_a \nabla^\nu g\right)(t) &= \frac{H(\nu-1)}{2-\nu} \left\{ \sum_{r=a+1}^t E_{\bar{\nu-1}}(\lambda_1, t-r+1) (\nabla_r g)(r) - \sum_{r=a+1}^{t-1} E_{\bar{\nu-1}}(\lambda_1, t-r) (\nabla_r g)(r) \right\} \\ &= \frac{H(\nu-1)}{2-\nu} \left\{ (2-\nu) (\nabla g)(t) + \sum_{r=a+1}^{t-1} \left[E_{\bar{\nu-1}}(\lambda_1, t-r+1) - E_{\bar{\nu-1}}(\lambda_1, t-r) \right] (\nabla_r g)(r) \right\} \\ &= H(\nu-1) \left\{ (\nabla g)(t) + \frac{1}{2-\nu} \left[E_{\bar{\nu-1}}(\lambda_1, t-a) - E_{\bar{\nu-1}}(\lambda_1, t-a-1) \right] (\nabla g)(a+1) \right. \\ &\quad \left. + \frac{1}{2-\nu} \sum_{r=a+2}^{t-1} \left[E_{\bar{\nu-1}}(\lambda_1, t-r+1) - E_{\bar{\nu-1}}(\lambda_1, t-r) \right] (\nabla_r g)(r) \right\}, \end{aligned}$$

for each $t \in \mathbb{N}_{a+3}$, and hence the proof is complete. \square

The first main result we present, Theorem 2.1, demonstrates that $({}^{\text{ABR}}_a \nabla^\nu g)(a+3)$ can be negative even though $(\nabla g)(a+3) > 0$ – cf., [14, Theorem 3.1].

Theorem 2.1. *Let the function g be defined on \mathbb{N}_a , and let $1 < \nu < 1.5$ and $\varepsilon > 0$. Assume that*

$$({}^{\text{ABR}}_a \nabla^\nu g)(a+3) > -\varepsilon (\nabla g)(a+1) H(\nu-1). \quad (2.1)$$

If $(\nabla g)(a+1) \geq 0$, $(\nabla g)(a+2) \geq 0$, and $\frac{1}{2}(\nu-1)^2(2\nu^2-5\nu+2) < -\varepsilon$, then $(\nabla g)(a+3) \geq 0$.

Proof. Due to Lemma 2.1 and the condition (2.1) we get

$$\begin{aligned} (\nabla g)(t) \geq & -(\nabla g)(a+1) \left\{ \frac{1}{2-\nu} \left[\mathbb{E}_{\nu-1}(\lambda_1, t-a) - \mathbb{E}_{\nu-1}(\lambda_1, t-a-1) \right] + \varepsilon \right\} \\ & - \frac{1}{2-\nu} \sum_{r=a+2}^{t-1} \left[\mathbb{E}_{\nu-1}(\lambda_1, t-r+1) - \mathbb{E}_{\nu-1}(\lambda_1, t-r) \right] (\nabla_r g)(r), \quad (2.2) \end{aligned}$$

for each $t \in \mathbb{N}_{a+3}$. Setting $t = a+3$ in (2.2), yields

$$\begin{aligned} (\nabla g)(a+3) \geq & -(\nabla g)(a+1) \left\{ \frac{1}{2-\nu} \left[\mathbb{E}_{\nu-1}(\lambda_1, 3) - \mathbb{E}_{\nu-1}(\lambda_1, 2) \right] + \varepsilon \right\} \\ & - \frac{1}{2-\nu} \sum_{r=a+2}^{a+2} \left[\mathbb{E}_{\nu-1}(\lambda_1, a+4-r) - \mathbb{E}_{\nu-1}(\lambda_1, a+3-r) \right] (\nabla_r g)(r). \end{aligned}$$

Since $(\nabla g)(a+2) \geq 0$ by assumption, it follows that

$$\begin{aligned} - \frac{1}{2-\nu} \sum_{r=a+2}^{a+2} \left[\mathbb{E}_{\nu-1}(\lambda_1, a+4-r) - \mathbb{E}_{\nu-1}(\lambda_1, a+3-r) \right] (\nabla_r g)(r) \\ = - \frac{1}{2-\nu} \underbrace{\left[-(2-\nu)(\nu-1)^2 \right]}_{>0} \underbrace{(\nabla g)(a+2)}_{\geq 0} \geq 0. \quad (2.3) \end{aligned}$$

Also, we know that $(\nabla g)(a+1) \geq 0$. So, we can use the inequalities (2.2) and (2.3) to deduce that $(\nabla g)(a+3) \geq 0$, using especially that

$$\frac{1}{2-\nu} \left[\mathbb{E}_{\nu-1}(\lambda_1, 3) - \mathbb{E}_{\nu-1}(\lambda_1, 2) \right] + \varepsilon = \frac{1}{2}(\nu-1)^2(2\nu^2-5\nu+2) + \varepsilon < 0$$

by the assumption given in the statement of the theorem, and this ends the proof. \square

Remark 2.2. *Figure 1 shows the graph of $\nu \mapsto -\frac{1}{2}(\nu-1)^2(2\nu^2-5\nu+2)$ for $\nu \in \left(1, \frac{3}{2}\right)$. Observe that in order for Theorem 2.1 to be applied it must hold that $\varepsilon \in \left(0, -\frac{1}{2}(\nu-1)^2(2\nu^2-5\nu+2)\right)$ for a fixed $\nu \in \left(1, \frac{3}{2}\right)$. This admissible region for ε is shown by the light grey region in the figure.*




Figure 1. Graph of $-v \mapsto \frac{1}{2}(v-1)^2(2v^2 - 5v + 2)$ for $v \in \left[1, \frac{3}{2}\right]$. The admissible range of ε for v fixed is shaded in light grey.

Now, define the set $\mathcal{F}_{k,\varepsilon} \subseteq \left(1, \frac{3}{2}\right)$ as follows:

$$\mathcal{F}_{k,\varepsilon} := \left\{ v \in \left(1, \frac{3}{2}\right) : \frac{1}{2-v} \left[\mathbb{E}_{v-1}(\lambda_1, k-a) - \mathbb{E}_{v-1}(\lambda_1, k-a-1) \right] < -\varepsilon \right\} \subseteq (1, 1.5), \quad \forall k \in \mathbb{N}_{a+3}.$$

Lemma 2.2 proves that the nested collection $\{\mathcal{F}_{k,\varepsilon}\}_{k=a+1}^{\infty}$ is decreasing (whenever $\varepsilon > 0$). This is a phenomenon that has been observed in similar contexts (e.g., [14, Lemma 3.3], [20, Lemma 3.2], [21, Lemma 3.2]).

Lemma 2.2. *Let $1 < v < \frac{3}{2}$. Then for each $\varepsilon > 0$ and $k \in \mathbb{N}_{a+3}$ we have that $\mathcal{F}_{k+1,\varepsilon} \subseteq \mathcal{F}_{k,\varepsilon}$.*

Proof. Let $\varepsilon > 0$ and $v \in \mathcal{F}_{k+1,\varepsilon}$ be arbitrary for some fixed $k \in \mathbb{N}_{a+3}$. Then we have

$$\frac{1}{2-v} \left[\mathbb{E}_{v-1}(\lambda_1, k+1-a) - \mathbb{E}_{v-1}(\lambda_1, k-a) \right] = \frac{\lambda_1}{2-v} \mathbb{E}_{v-1,v-1}(\lambda_1, k+1-a) < -\varepsilon.$$

Since $\mathbb{E}_{v-1,v-1}(\lambda_1, k+1-a)$ is decreasing for each $k \in \mathbb{N}_{a+3}$ (see [4]), $1 < v < \frac{3}{2}$, and $\lambda_1 < 0$, we have

$$\frac{\lambda_1}{2-v} \mathbb{E}_{v-1,v-1}(\lambda_1, k-a) < \frac{\lambda_1}{2-v} \mathbb{E}_{v-1,v-1}(\lambda_1, k+1-a) < -\varepsilon.$$

This implies that $v \in \mathcal{F}_{k,\varepsilon}$, and, therefore, $\mathcal{F}_{k+1,\varepsilon} \subseteq \mathcal{F}_{k,\varepsilon}$. Thus, we have accomplished the result. \square

Theorem 2.1 and Lemma 2.2 now lead to the following corollary, which is the principal analytical result of this note – cf., [14, Corollary 3.4]. Corollary 2.1 asserts that the same pathological behavior observed with other discrete fractional differences carries over to the Mittag-Leffler kernel setting.

Corollary 2.1. *Let the assumptions of Theorem 2.1 be fulfilled together with*

$$\left({}^{\text{ABR}}_a \nabla^v g \right) (t) > -\varepsilon \mathbb{H}(v-1) (\nabla g)(a+1), \quad (2.4)$$

for each $\nu \in \left(1, \frac{3}{2}\right)$, $t \in \mathbb{N}_{a+3}^s$ and some fixed $s \in \mathbb{N}_{a+3}$. Now, if we assume that $(\nabla g)(a+1) \geq 0$, $(\nabla g)(a+2) \geq 0$, and $\nu \in \mathcal{F}_{s,\varepsilon}$, then we have $(\nabla g)(t) \geq 0$, for all $t \in \mathbb{N}_{a+1}^s$.

Proof. Due to the assumption that $\nu \in \mathcal{F}_{s,\varepsilon}$ and Lemma 2.2, we have

$$\nu \in \mathcal{F}_{s,\varepsilon} = \mathcal{F}_{s,\varepsilon} \cap \bigcap_{k=a+3}^{s-1} \mathcal{F}_{k,\varepsilon}.$$

This leads to

$$\frac{1}{2-\nu} \left[\mathbb{E}_{\nu-1}(\lambda_1, k+1-a) - \mathbb{E}_{\nu-1}(\lambda_1, k-a) \right] < -\varepsilon, \quad (2.5)$$

for each $k \in \mathbb{N}_{a+3}^s$.

We now can proceed by induction to complete the proof as follows. At first, for $t = a+3$ we can obtain $(\nabla g)(a+3) \geq 0$ immediately as in Theorem 2.1 with the help of inequalities (2.4) and (2.5) just as in the proof of Theorem 2.1. Accordingly, we can continue and inductively iterate inequality (2.2) to get $(\nabla g)(t) \geq 0$, for all $t \in \mathbb{N}_{a+2}^s$ as requested. Note that in this last step we are using the fact that $\mathbb{E}_{\nu-1}(\lambda_1, t-r+1) - \mathbb{E}_{\nu-1}(\lambda_1, t-r) \geq 0$, for each $(r, t) \in \mathbb{N}_{a+2}^{t-1} \times \mathbb{N}_{a+3}^s$, which is true since the partial function $t \mapsto \mathbb{E}_{\nu-1}(\lambda_1, t)$ is decreasing – see Remark 2.1. Thus, we have completed the proof. \square

We next provide an example in order to demonstrate the application of the preceding ideas.

Example 2.1. Considering Lemma 2.1 with $t := a+3$:

$$\begin{aligned} \left({}^{\text{ABR}}_a \nabla^\nu g \right) (a+3) &= H(\nu-1) \left\{ (\nabla g)(a+3) + \frac{1}{2-\nu} \left[\mathbb{E}_{\nu-1}(\lambda_1, 3) - \mathbb{E}_{\nu-1}(\lambda_1, 2) \right] (\nabla g)(a+1) \right. \\ &\quad \left. + \frac{1}{2-\nu} \sum_{r=a+2}^{a+2} \left[\mathbb{E}_{\nu-1}(\lambda_1, a+4-r) - \mathbb{E}_{\nu-1}(\lambda_1, a+3-r) \right] (\nabla_r g)(r) \right\}. \end{aligned}$$

For $a = 0$, it follows that

$$\begin{aligned} &\left({}^{\text{ABR}}_0 \nabla^\nu g \right) (3) \\ &= H(\nu-1) \left\{ (\nabla g)(3) + \frac{1}{2}(\nu-1)^2(2\nu^2 - 5\nu + 2)(\nabla g)(1) \right. \\ &\quad \left. + \frac{1}{2-\nu} \sum_{r=2}^2 \left[\mathbb{E}_{\nu-1}(\lambda_1, 4-r) - \mathbb{E}_{\nu-1}(\lambda_1, 3-r) \right] (\nabla_r g)(r) \right\} \\ &= H(\nu-1) \left\{ (\nabla g)(3) + \frac{1}{2}(\nu-1)^2(2\nu^2 - 5\nu + 2)(\nabla g)(1) - (\nu-1)^2(\nabla g)(2) \right\} \\ &= H(\nu-1) \left\{ g(3) - g(2) + \frac{1}{2}(\nu-1)^2(2\nu^2 - 5\nu + 2)[g(1) - g(0)] - (\nu-1)^2[g(2) - g(1)] \right\}. \end{aligned}$$

If we take $\nu = 1.99$, $g(0) = 0.01$, $g(1) = 1.01$, $g(2) = 1.001$, $g(3) = 1.005$, and $\varepsilon = 0.002$, we have

$$\left({}^{\text{ABR}}_0 \nabla^{1.99} g \right) (3) = -0.0018 H(0.99) > -0.002 H(0.99) = -\varepsilon H(0.99) (\nabla g)(1).$$

Note that $\left({}^{\text{ABR}}_0 \nabla^{1.99} g \right) (3) < 0$. Yet, as Theorem 2.1 correctly predicts, it, nonetheless, holds that $(\nabla g)(1) > 0$. Thus, the collection of functions to which Theorem 2.1 applies is non-void.

We conclude this note by providing a brief numerical analysis of the set $\mathcal{F}_{k,\varepsilon}$, which plays a key role in Corollary 2.1; throughout we take $a = 0$ purely for convenience. Let us first consider Figure 2 above. This is a heat map, which identifies the cardinality of the set $\{k \in \mathbb{N} : \nu \in \mathcal{F}_{k,\varepsilon}\}$. It is worth mentioning that the warmer colors (i.e., oranges and reds) are associated to higher cardinality values as indicated by the vertical sidebar in Figure 2. We see that there is a concentration of larger cardinalities in a roughly triangular region as indicated in the figure. The largest cardinalities seem to be concentrated for $1.07 \lesssim \nu \lesssim 1.10$ and $\varepsilon > 0$ close to zero; this implies that the analytical results presented earlier (i.e., Corollary 2.1) should be valid for the greatest number of time steps t when ν and ε are in this region of the (ν, ε) -parameter space.

On the other hand, Figures 3–6 plot the interval of k values such that $\mathcal{F}_{k,\varepsilon} \neq \emptyset$ for different choices of both ν and ε . Consistent with the heat map in Figure 2, we see that there is a relative maximum when ν is away from the boundary values $\nu = 1$ and $\nu = 1.5$, though the precise value depends on the value of ε . In particular, as $\varepsilon \rightarrow 0^+$, the maximum seems to approach about 1.07, just as indicated by Figure 2. In addition, we see that the length of the intervals drops off sharply both as $\nu \rightarrow 1^+$ and as $\nu \rightarrow 1.5^-$ – again, precisely as depicted in Figure 2.

Finally, the data contained in Figures 2–6 is not entirely dissimilar from the observations in [14, Figures 1–4], which analyzed the Riemann-Liouville fractional nabla difference. In each of the Mittag-Leffler kernel and the Riemann-Liouville settings the ν -values for which the respective monotonicity-type theorems – i.e., Corollary 2.1 and [14, Corollary 3.4] – seem to be most applicable are apparently concentrated for ν close to 1. A possible, albeit non-rigorous, explanation for this common observation is that when $\nu \approx 1$ the fractional difference is “more like” the first-order difference, which is closely connected to monotonicity. But we do not have a precise analytical explanation for this numerical observation, and we hope that this sort of curiosity provides motivation to analyze further these types of fractional difference operators in the future. Nonetheless, the results of this note show that this is a common feature across multiple types of fractional difference operators




Figure 2. Heat map for the cardinality of $\{k : \nu \in \mathcal{F}_{k,\varepsilon}\}$.




Figure 3. $\mathcal{F}_{k,0.01}$.




Figure 4. $\mathcal{F}_{k,0.005}$.




Figure 5. $\mathcal{F}_{k,0.0005}$.




Figure 6. $\mathcal{F}_{k,0.0001}$.

3. Conclusions

In this brief note we have demonstrated that the fractional difference with Mittag-Leffler kernel behaves in an aberrant manner, similar to that of other classes of nonlocal difference operators. In particular, we have shown that such a difference acting on a function can be negative even if the function on which it acts is increasing. This sort of unusual behavior is not possible when considering a local difference operator, but it seems to be an almost defining feature of nonlocal discrete operators as the results of this note demonstrate.

Acknowledgements

This research was supported by Taif University Researchers Supporting Project (No. TURSP-2020/155), Taif University, Taif, Saudi Arabia.

Conflicts of interest

The authors declare that they have no conflicts interests.

References

1. T. Abdeljawad, Different type kernel h -fractional differences and their fractional h -sums, *Chaos Solit. Fract.*, **116** (2018), 146–156. <https://doi.org/10.1016/j.chaos.2018.09.022>
2. T. Abdeljawad, F. Jarad, A. Atangana, P. O. Mohammed, On a new type of fractional difference operators on h -step isolated time scales, *J. Frac. Calc. Nonlinear Sys.*, **1** (2021), 46–74.
3. T. Abdeljawad, On delta and nabla caputo fractional differences and dual identities, *Discrete Dyn. Nat. Soc.*, **2013** (2013), 406910. <https://doi.org/10.1155/2013/406910>
4. T. Abdeljawad, D. Baleanu, Monotonicity analysis of a nabla discrete fractional operator with discrete Mittag-Leffler kernel, *Chaos Solit. Fract.*, **102** (2017), 106–110. <https://doi.org/10.1016/j.chaos.2017.04.006>
5. T. Abdeljawad, F. Madjidi, Lyapunov-type inequalities for fractional difference operators with discrete Mittag-Leffler kernel of order $2 < \alpha < 5/2$, *Eur. Phys. J. Spec. Top.*, **226** (2017), 3355–3368. <https://doi.org/10.1140/epjst/e2018-00004-2>
6. T. Abdeljawad, Q. M. Al-Mdallal, M. A. Hajji, Arbitrary order fractional difference operators with discrete exponential kernels and applications, *Discrete Dyn. Nat. Soc.*, **2017** (2017), 4149320. <https://doi.org/10.1155/2017/4149320>
7. F. M. Atici, M. Atici, M. Belcher, D. Marshall, A new approach for modeling with discrete fractional equations, *Fund. Inform.*, **151** (2017), 313–324. <https://doi.org/10.3233/FI-2017-1494>
8. F. M. Atici, M. Atici, N. Nguyen, T. Zhorojev, G. Koch, A study on discrete and discrete fractional pharmacokinetics-pharmacodynamics models for tumor growth and anti-cancer effects, *Comput. Math. Biophys.*, **7** (2019), 10–24. <https://doi.org/10.1515/cmb-2019-0002>
9. F. M. Atici, P. W. Eloe, A transform method in discrete fractional calculus, *Int. J. Differ. Equ.*, **2** (2007), 165–176.
10. F. M. Atici, P. W. Eloe, Discrete fractional calculus with the nabla operator, *Electron. J. Qual. Theory Differ. Equ.*, 2009.
11. F. M. Atici, P. W. Eloe, Initial value problems in discrete fractional calculus, *Proc. Amer. Math. Soc.*, **137** (2009), 981–989.
12. J. Bravo, C. Lizama, S. Rueda, Second and third order forward difference operator: what is in between? *RACSAM*, **115** (2021), 86. <https://doi.org/10.1007/s13398-021-01015-5>
13. R. Dahal, C. S. Goodrich, A monotonicity result for discrete fractional difference operators, *Arch. Math.*, **102** (2014), 293–299. <https://doi.org/10.1007/s00013-014-0620-x>
14. R. Dahal, C. S. Goodrich, Theoretical and numerical analysis of monotonicity results for fractional difference operators, *Appl. Math. Lett.*, **117** (2021), 107104. <https://doi.org/10.1016/j.aml.2021.107104>
15. F. F. Du, B. Jia, L. Erbe, A. Peterson, Monotonicity and convexity for nabla fractional q -differences, *Dynam. Syst. Appl.*, **22** (2016), 1224–1243. <https://doi.org/10.1080/10236198.2016.1188089>

16. C. S. Goodrich, J. M. Jonnalagadda, An analysis of polynomial sequences and their application to discrete fractional operators, *J. Differ. Equ. Appl.*, **27** (2021), 1081–1102. <https://doi.org/10.1080/10236198.2021.1965132>
17. C. S. Goodrich, J. M. Jonnalagadda, B. Lyons, Convexity, monotonicity and positivity results for sequential fractional nabla difference operators with discrete exponential kernels, *Math. Method. Appl. Sci.*, **44** (2021), 7099–7120. <https://doi.org/10.1002/mma.7247>
18. C. S. Goodrich, C. Lizama, A transference principle for nonlocal operators using a convolutional approach: Fractional monotonicity and convexity, *Israel J. Math.*, **236** (2020), 533–589. <https://doi.org/10.1007/s11856-020-1991-2>
19. C. S. Goodrich, C. Lizama, Positivity, monotonicity, and convexity for convolution operators, *Discrete Contin. Dyn. Syst.*, **40** (2020), 4961–4983. <https://doi.org/10.3934/dcds.2020207>
20. C. S. Goodrich, B. Lyons, M. T. Velcsov, Analytical and numerical monotonicity results for discrete fractional sequential differences with negative lower bound, *Commun. Pur. Appl. Anal.*, **20** (2021), 339–358. <https://doi.org/10.3934/cpaa.2020269>
21. C. S. Goodrich, B. Lyons, A. Scapellato, M. T. Velcsov, Analytical and numerical convexity results for discrete fractional sequential differences with negative lower bound, *J. Differ. Equ. Appl.*, **27** (2021), 317–341. <https://doi.org/10.1080/10236198.2021.1894142>
22. C. S. Goodrich, M. Muellner, An analysis of the sharpness of monotonicity results via homotopy for sequential fractional operators, *Appl. Math. Lett.*, **98** (2019), 446–452. <https://doi.org/10.1016/j.aml.2019.07.003>
23. C. Goodrich, A. C. Peterson, *Discrete fractional calculus*, New York: Springer, 2015.
24. B. Jia, L. Erbe, A. Peterson, Two monotonicity results for nabla and delta fractional differences, *Arch. Math.*, **104** (2015), 589–597.
25. C. Lizama, The Poisson distribution, abstract fractional difference equations, and stability, *Proc. Amer. Math. Soc.*, **145** (2017), 3809–3827.
26. P. O. Mohammed, T. Abdeljawad, F. K. Hamasalh, On Riemann-Liouville and Caputo fractional forward difference monotonicity analysis, *Mathematics*, **9** (2021), 1303. <https://doi.org/10.3390/math9111303>
27. P. O. Mohammed, T. Abdeljawad, Discrete generalized fractional operators defined using h-discrete Mittag-Leffler kernels and applications to AB fractional difference systems, *Math. Meth. Appl. Sci.*, 2020, 1–26. <https://doi.org/10.1002/mma.7083>
28. M. Yavuz, T. A. Sulaiman, A. Yusuf, T. Abdeljawad, The Schrödinger-KdV equation of fractional order with Mittag-Leffler nonsingular kernel, *Alex. Eng. J.*, **60** (2021), 2715–2724. <https://doi.org/10.1016/j.aej.2021.01.009>
29. K. Zhao, Existence, stability and simulation of a class of nonlinear fractional Langevin equations involving nonsingular Mittag-Leffler kernel, *Fractal Fract.*, **6** (2021), 469. <https://doi.org/10.3390/fractalfract6090469>