



Research article

Simulating systems of Itô SDEs with split-step (α, β) -Milstein scheme

Hassan Ranjbar¹, Leila Torkzadeh^{1,*}, Dumitru Baleanu^{2,3,4} and Kazem Nouri¹

¹ Department of Mathematics, Faculty of Mathematics, Statistics and Computer Sciences, Semnan University, P.O. Box, 35195–363, Semnan, Iran

² Department of Mathematics, Faculty of Arts and Sciences, Cankaya University, Ankara, Turkey

³ Institute of Space Sciences, P.O. Box, MG-23, R 76900 Magurele-Bucharest, Romania

⁴ Department of Medical Research, China Medical University Hospital, China Medical University, Taichung, Taiwan

* **Correspondence:** Email: torkzadeh@semnan.ac.ir.

Abstract: In the present study, we provide a new approximation scheme for solving stochastic differential equations based on the explicit Milstein scheme. Under sufficient conditions, we prove that the split-step (α, β) -Milstein scheme strongly convergence to the exact solution with order 1.0 in mean-square sense. The mean-square stability of our scheme for a linear stochastic differential equation with single and multiplicative commutative noise terms is studied. Stability analysis shows that the mean-square stability of our proposed scheme contains the mean-square stability region of the linear scalar test equation for suitable values of parameters α, β . Finally, numerical examples illustrate the effectiveness of the theoretical results.

Keywords: Itô stochastic ordinary differential equations; mean-square convergence; mean-square stability; split-step Milstein scheme

Mathematics Subject Classification: 34F05, 41A25, 60H10, 93E15

1. Introduction

In this paper, we deal with the following Itô stochastic differential equations (SDEs)

$$dW_t = A(W_t, t)dt + \sum_{p=1}^P B_p(W_t, t)d\varpi_t^p, t \geq 0. \tag{1.1}$$

Here $A, B_p : \mathbb{R}^d \times [t_0, T] \rightarrow \mathbb{R}^d$, are drift and diffusion functions, respectively. Also $\varpi_t^p, p = 1, \dots, P$ is an one-dimensional Wiener process. This type of SDE (1.1) are widely applied for describing many

real-life phenomena such as economics, epidemiology, chemistry, meteorology and etc, see [1–8], for example. For many of them, there is no analytical closed-form solution, so numerical techniques and analysis will become important. For SDE (1.1), Milstein [9] proposed an explicit numerical scheme with strong convergence order 1.0, namely

$$\mathcal{W}_{l+1}^{\text{Mil}} = \mathcal{W}_l^{\text{Mil}} + hA(\mathcal{W}_l^{\text{Mil}}, t) + \sum_{p=1}^P B_p(\mathcal{W}_l^{\text{Mil}}, t) \mathcal{I}_{(p)} + \sum_{p_1=1}^P \sum_{p_2=1}^P L^{p_1} B_{p_2}(\mathcal{W}_l^{\text{Mil}}, t) \mathcal{I}_{(p_1, p_2)}, \quad (1.2)$$

with

$$\mathcal{I}_{(p)} = \int_{t_l}^{t_l+h} d\varpi_p(r) \cong \Delta\varpi_{p,l} = \sqrt{h}\xi_p, \quad \mathcal{I}_{(p_1, p_2)} = \int_{t_l}^{t_l+h} \left(\int_{t_l}^{r_1} d\varpi_{p_1}(r_2) \right) d\varpi_{p_2}(r_1),$$

and

$$L^{p_1} = \sum_{i=1}^d B_{i, p_1} \frac{\partial}{\partial \mathcal{W}_{i,l}^{\text{Mil}}},$$

such that

$$\sum_{p_1, p_2=1}^P L^{p_1} B_{p_2}(\mathcal{W}_l^{\text{Mil}}, t_l) = \sum_{p_1=1}^P \sum_{p_2=1}^P \sum_{i=1}^d \frac{\partial}{\partial \mathcal{W}_{i,l}^{\text{Mil}}} \begin{bmatrix} B_{p_2,1} \\ \vdots \\ B_{p_2,d} \end{bmatrix} B_{p_1,i} = \sum_{i=1}^d \begin{bmatrix} \frac{\partial B_{1,1}}{\partial \mathcal{W}_{i,l}^{\text{Mil}}} & \cdots & \frac{\partial B_{p_1,1}}{\partial \mathcal{W}_{i,l}^{\text{Mil}}} \\ \vdots & \ddots & \vdots \\ \frac{\partial B_{1,d}}{\partial \mathcal{W}_{i,l}^{\text{Mil}}} & \cdots & \frac{\partial B_{p_1,d}}{\partial \mathcal{W}_{i,l}^{\text{Mil}}} \end{bmatrix} \begin{bmatrix} B_{1,i} \\ \vdots \\ B_{p_1,i} \end{bmatrix},$$

where $t_l = t_0 + lh$, $l = 0, 1, \dots, N$, a time step $h = (T - t_0)/N$ with a fixed natural number N and $\xi_p \sim \mathcal{N}(0, 1)$. Furthermore, drifting split-step backward Milstein (DSSBM) scheme is given by [10]

$$\begin{aligned} \widetilde{\mathcal{W}}_l &= \mathcal{W}_l + hA(\widetilde{\mathcal{W}}_l, t_l), \\ \mathcal{W}_{l+1} &= \widetilde{\mathcal{W}}_l + \sum_{p=1}^P B_p(\widetilde{\mathcal{W}}_l, t_l) \mathcal{I}_{(p)} + \sum_{p_1=1}^P \sum_{p_2=1}^P L^{p_1} B_{p_2}(\widetilde{\mathcal{W}}_l, t_l) \mathcal{I}_{(p_1, p_2)}. \end{aligned} \quad (1.3)$$

The drifting split-step Adams-Moulton Milstein scheme is another modification of the classical Milstein scheme, which was initialized in [11]. Recently, Jiang et al. [12] propose a new split two-step Milstein scheme as follow

$$\begin{aligned} \widetilde{\mathcal{W}}_l &= \mathcal{W}_l - h\beta_2 A(\widetilde{\mathcal{W}}_{l-1}) + h\beta_0 A(\widetilde{\mathcal{W}}_l) - \frac{\sigma}{2} hL^1 B(\widetilde{\mathcal{W}}_l), \\ \mathcal{W}_{l+1} &= \mathcal{W}_l + B(\widetilde{\mathcal{W}}_l) \mathcal{I}_{(1)} + L^1 B(\widetilde{\mathcal{W}}_l) (|\mathcal{I}_{(1)}|^2 - h), \end{aligned} \quad (1.4)$$

for autonomous stochastic differential systems with one-dimensional Wiener process. Especially, they named method (1.4), Adams-Bashforth Milstein (ABM) scheme when $\beta_2 = 0$, $\beta_0 = -1/2$ and $\sigma = 1$. Furthermore, method (1.4) called Adams-Moulton Milstein (AMM) scheme if $\beta_2 = 5/12$, $\beta_0 = -1/2$ and $\sigma = 1/2$. Also, the family of drift-implicit Milstein schemes has been found in [13, 14], which is adapted for stiff stochastic problems. Other families of numerical schemes were also studied, for instance [15–21]. We just mention some of them here and refer the readers to the references therein, among others.

One way to judge numerical schemes is to use the mean-square (MS-) stability properties. Saito and Mitsui [22] studied MS-stability properties of some numerical schemes for linear test equations,

as well. Also, in [23] the same authors, analyzed the MS-stability of the Euler-Maruyama scheme for linear systems of SDEs. For more study of the MS-stability of various linear systems of SDEs applied to numerical schemes, the reader may refer to [24–27] and the references therein.

We emphasize that this paper is motivated by the Higham et al. [28], where proposed a new Milstein scheme and they investigated its efficiency in many financial models.

For SDE (1.1), the numerical scheme for the split-step (α, β) -Milstein (SSABM) approximation is defined by

$$\overline{\mathcal{W}}_l = \mathcal{W}_l + \alpha h A(\overline{\mathcal{W}}_l, t_l), \quad (1.5a)$$

$$\widehat{\mathcal{W}}_l = \overline{\mathcal{W}}_l - \frac{h}{2} \beta \sum_{p=1}^P L^p B_p(\widehat{\mathcal{W}}_l, t_l), \quad (1.5b)$$

$$\mathcal{W}_{l+1} = \mathcal{W}_l + h A(\widehat{\mathcal{W}}_l, t_l) + \sum_{p=1}^P B_p(\widehat{\mathcal{W}}_l, t_l) \mathcal{I}_{(p)} + \sum_{p_1=1}^P \sum_{p_2=1}^P L^{p_1} B_{p_2}(\widehat{\mathcal{W}}_l, t_l) \mathcal{I}_{(p_1, p_2)}, \quad (1.5c)$$

where $\alpha, \beta \in [0, 1]$. By taking $\alpha = \beta = 0$ in (1.5) Milstein method [9] is obtained. We gives the split-step theta-Milstein (SSTM) method [29], when $\alpha = \theta, \beta = 0$. Furthermore if $\alpha = 1, \beta = 0$, we obtain DSSBM scheme [10]. Obviously, deterministic equations (1.5a) and (1.5b) are implicit in $\overline{\mathcal{W}}_l$ and $\widehat{\mathcal{W}}_l$ when $\alpha, \beta \in (0, 1]$ must be solved to obtain the intermediate approximation $\overline{\mathcal{W}}_l$ and $\widehat{\mathcal{W}}_l$, respectively.

In this work the consider the strong convergence properties of the numerical scheme (1.5) in MS sense, we shall follow [30–32], where assumed that the drift and diffusion coefficients function of SDE (1.1) satisfies the following conditions.

Proposition 1.1. *There exist constants $\ell_1 > 0$ and $\ell_2 > 0$ such that*

-Lipschitz conditions:

$$\begin{aligned} & |A(s_1, t) - A(s_2, t)|^2 \vee \sum_{p=1}^P |B_p(s_1, t) - B_p(s_2, t)|^2 \vee \sum_{p_1=1}^P \sum_{p_2=1}^P |L^{p_1} B_{p_2}(s_1, t) - L^{p_1} B_{p_2}(s_2, t)|^2 \\ & \leq \ell_1 |s_1 - s_2|^2. \end{aligned} \quad (1.6)$$

-Linear growth bounds:

$$|A(s_1, t)|^2 \vee \sum_{p=1}^P |B_p(s_1, t)|^2 \vee \sum_{p_1=1}^P \sum_{p_2=1}^P |L^{p_1} B_{p_2}(s_1, t)|^2 \leq \ell_2 (1 + |s_1|^2), \quad (1.7)$$

for all $s_1, s_2 \in \mathbb{R}^d$.

This paper is constructed as follows. We will devote to our main results about the MS convergence of the numerical scheme (1.5) in Section 2. And then, the MS-stability properties of the SSABM scheme (1.5) are established in Section 3. Section 4 contains examples. The conclusion is stated in Section 5.

2. Mean-square convergence analysis

To recover the strong convergence order 1.0 for the scheme (1.5), we need Proposition 1.1, following Lemma and the local mean error and mean-square error Milstein scheme (1.2):

$$\left| \mathbb{E} \left[(\mathcal{W}_{l+1}^{\text{Mil}} - W(t_{l+1})) \middle| F_l \right] \right| \leq Kh^2 \sqrt{1 + |\mathcal{W}_l|^2}, \quad (2.1a)$$

$$\left| \mathbb{E} \left[|\mathcal{W}_{l+1}^{\text{Mil}} - W(t_{l+1})|^2 \middle| F_l \right] \right|^{\frac{1}{2}} \leq Kh^{3/2} \sqrt{1 + |\mathcal{W}_l|^2}, \quad (2.1b)$$

respectively.

Lemma 2.1. [33] Assume for a one-step discrete time approximation \mathcal{W} that the local mean error and mean-square error for all $N = 1, 2, \dots$, and $l = 0, 1, \dots, N - 1$ satisfy the estimates

$$\left| \mathbb{E} [(\mathcal{W}_{l+1} - W_{t_{l+1}}) | F_l] \right| \leq Kh^{\kappa_1} \sqrt{1 + |\mathcal{W}_l|^2}, \quad (2.2)$$

and

$$\left| \mathbb{E} \left[|\mathcal{W}_{l+1} - W_{t_{l+1}}|^2 | F_l \right] \right|^{1/2} \leq Kh^{\kappa_2} \sqrt{1 + |\mathcal{W}_l|^2}, \quad (2.3)$$

when $\kappa_2 \geq \frac{1}{2}$ and $\kappa_1 \geq \kappa_2 + \frac{1}{2}$. Then

$$\left| \mathbb{E} \left[|\mathcal{W}_r - W_{t_r}|^2 | F_0 \right] \right|^{1/2} \leq Kh^{\kappa_2 - 1/2} \sqrt{1 + |\mathcal{W}_r|^2},$$

holds for each $r = 0, 1, \dots, N$.

Theorem 2.1. Suppose Proposition 1.1 holds. Then the numerical scheme SSABM (1.5) strongly converges to SDE (1.1) in the MS sense with order 1.0.

Proof. First, we compute the local mean error of our scheme in the following

$$\begin{aligned} \delta_1 &= \left| \mathbb{E} \left[(\mathcal{W}_{l+1} - W_{t_{l+1}}) \middle| F_l \right] \right| \\ &\leq \left| \mathbb{E} \left[(\mathcal{W}_{l+1}^{\text{Mil}} - W_{t_{l+1}}) \middle| F_l \right] \right| + \left| \mathbb{E} \left[(\mathcal{W}_{l+1} - \mathcal{W}_{l+1}^{\text{Mil}}) \middle| F_l \right] \right| \leq Kh^2 \sqrt{1 + |\mathcal{W}_l|^2} + \delta_2, \end{aligned} \quad (2.4)$$

where

$$\begin{aligned} \delta_2 &= \left| \mathbb{E} \left[(\mathcal{W}_{l+1} - \mathcal{W}_{l+1}^{\text{Mil}}) \middle| F_l \right] \right| \\ &= \left| \mathbb{E} \left[\mathcal{W}_l - \mathcal{W}_l^{\text{Mil}} + h \left(A(\widehat{\mathcal{W}}_l, t_l) - A(\mathcal{W}_l^{\text{Mil}}, t_l) \right) + \sum_{p=1}^P \left(B_p(\widehat{\mathcal{W}}_l, t_l) - B_p(\mathcal{W}_l^{\text{Mil}}, t_l) \right) \mathcal{I}_{(p)} \right. \right. \\ &\quad \left. \left. + \sum_{p_1=1}^P \sum_{p_2=1}^P \left(L^{p_1} B_{p_2}(\widehat{\mathcal{W}}_l, t_l) - L^{p_1} B_{p_2}(\mathcal{W}_l^{\text{Mil}}, t_l) \right) \mathcal{I}_{(p_1, p_2)} \right] \middle| F_l \right| \\ &\leq h \sqrt{\ell_1} \left| \widehat{\mathcal{W}}_l - \mathcal{W}_l \right|. \end{aligned} \quad (2.5)$$

Notice that for obtain of inequality (2.5), used global Lipschitz condition (1.6), $\mathbb{E}[\mathcal{I}_{(p)}] = \mathbb{E}[\mathcal{I}_{(p_1, p_2)}] = 0$. Also, applying (1.5a), (1.5b), Proposition 1.1 and

$$\sqrt{L_1 + L_2 + \dots + L_p} \leq \sqrt{L_1} + \sqrt{L_2} + \dots + \sqrt{L_p}, \quad \{L_i\}_{i=1}^p \geq 0,$$

yields

$$|\widehat{\mathcal{W}}_l - \mathcal{W}_l| \leq |\widehat{\mathcal{W}}_l - \overline{\mathcal{W}}_l| + |\overline{\mathcal{W}}_l - \mathcal{W}_l|, \quad (2.6)$$

where

$$\begin{aligned} |\widehat{\mathcal{W}}_l - \overline{\mathcal{W}}_l| &\leq \left| -\frac{\beta h}{2} \sum_{p=1}^P L^p B_p(\widehat{\mathcal{W}}_l, t_l) \right| \\ &\leq \frac{\beta h}{2} \sum_{p=1}^P |L^p B_p(\widehat{\mathcal{W}}_l, t_l) - L^p B_p(\overline{\mathcal{W}}_l, t_l)| \\ &\quad + \frac{\beta h}{2} \sum_{p=1}^P |L^p B_p(\overline{\mathcal{W}}_l, t_l) - L^p B_p(\mathcal{W}_l, t_l)| + \frac{\beta h}{2} \sum_{p=1}^P |L^p B_p(\mathcal{W}_l, t_l)| \\ &\leq \frac{\beta h}{2} \sqrt{\ell_1} |\widehat{\mathcal{W}}_l - \overline{\mathcal{W}}_l| + \frac{\beta h}{2} \sqrt{\ell_1} |\overline{\mathcal{W}}_l - \mathcal{W}_l| + \frac{\beta h}{2} \sqrt{\ell_2} \sqrt{1 + |\mathcal{W}_l|^2}, \end{aligned}$$

and

$$\begin{aligned} |\overline{\mathcal{W}}_l - \mathcal{W}_l| &\leq |\alpha h A(\overline{\mathcal{W}}_l, t_l)| \leq \alpha h |A(\overline{\mathcal{W}}_l, t_l) - A(\mathcal{W}_l, t_l)| + \alpha h |A(\mathcal{W}_l, t_l)| \\ &\leq \alpha h \sqrt{\ell_1} |\overline{\mathcal{W}}_l - \mathcal{W}_l| + \alpha h \sqrt{\ell_2} \sqrt{1 + |\mathcal{W}_l|^2}. \end{aligned}$$

From the above inequalities, we conclude that

$$|\overline{\mathcal{W}}_l - \mathcal{W}_l| \leq h \frac{\alpha \sqrt{\ell_2}}{1 - h\alpha \sqrt{\ell_1}} \sqrt{1 + |\mathcal{W}_l|^2}, \quad (2.7)$$

and

$$|\widehat{\mathcal{W}}_l - \overline{\mathcal{W}}_l| \leq h \frac{\beta \sqrt{\ell_2}}{2(1 - h\frac{\beta}{2} \sqrt{\ell_1})(1 - h\alpha \sqrt{\ell_1})} \sqrt{1 + |\mathcal{W}_l|^2}. \quad (2.8)$$

Now for $(1 - h\frac{\beta}{2} \sqrt{\ell_1})(1 - h\alpha \sqrt{\ell_1}) > 0$, we have from (2.4)–(2.8), $\varkappa_1 = h^2$. Similarly by standard arguments, we can prove

$$\begin{aligned} \delta_3 &= \left| \mathbb{E} \left[\left| \mathcal{W}_{l+1} - W_{l+1} \right|^2 \middle| F_t \right] \right|^{\frac{1}{2}} \\ &\leq \left| \mathbb{E} \left[\left| W_{l+1}^{\text{Mil}} - W_{l+1} \right|^2 \middle| F_t \right] \right|^{\frac{1}{2}} + \left| \mathbb{E} \left[\left| \mathcal{W}_{l+1} - W_{l+1}^{\text{Mil}} \right|^2 \middle| F_t \right] \right|^{\frac{1}{2}} \leq Kh^{3/2} \sqrt{1 + |\mathcal{W}_l|^2} + \sqrt{|\delta_4|}. \end{aligned} \quad (2.9)$$

By $\mathbb{E}[\mathcal{I}_{(p)}^2 | F_t] \leq O(h)$, $\mathbb{E}[\mathcal{I}_{(p_1, p_2)}^2 | F_t] \leq O(h^2)$ [6, Lemma 5.7.2] and inequality

$$(L_1 + L_2 + \dots + L_p)^2 \leq P(L_1^2 + L_2^2 + \dots + L_p^2), \quad (2.10)$$

we can obtain

$$\begin{aligned}
\delta_4 &= \mathbb{E} \left[\left| \mathcal{W}_{l+1} - W_{l+1}^{\text{Mil}} \right|^2 \middle| F_t \right] \\
&= \mathbb{E} \left[\left| \mathcal{W}_l - W_l^{\text{Mil}} + h \left(A(\widehat{\mathcal{W}}_l, t_l) - A(W_l^{\text{Mil}}, t_l) \right) + \sum_{p=1}^P \left(B_p(\widehat{\mathcal{W}}_l, t_l) - B_p(W_l^{\text{Mil}}, t_l) \right) \mathcal{I}_{(p)} \right. \right. \\
&\quad \left. \left. + \sum_{p_1=1}^P \sum_{p_2=1}^P \left(L^{p_1} B_{p_2}(\widehat{\mathcal{W}}_l, t_l) - L^{p_1} B_{p_2}(W_l^{\text{Mil}}, t_l) \right) \mathcal{I}_{(p_1, p_2)} \right|^2 \middle| F_t \right] \\
&\leq h \left(1 + P + P^2 \right) \left(h \left| A(\widehat{\mathcal{W}}_l, t_l) - A(W_l, t_l) \right|^2 + \sum_{p=1}^P \left| B_p(\widehat{\mathcal{W}}_l, t_l) - B_p(W_l, t_l) \right|^2 \right. \\
&\quad \left. + h \sum_{p=1}^P \left| L^{p_1} B_{p_2}(\widehat{\mathcal{W}}_l, t_l) - L^{p_1} B_{p_2}(W_l, t_l) \right|^2 \right) \\
&\leq h(1+2h)(1+P+P^2)\ell_1 \left| \widehat{\mathcal{W}}_l - \mathcal{W}_l \right|^2.
\end{aligned} \tag{2.11}$$

Using the (1.5a), (1.5b), Proposition 1.1 and inequality (2.10), we can get:

$$\left| \widehat{\mathcal{W}}_l - \mathcal{W}_l \right|^2 \leq 2 \left| \widehat{\mathcal{W}}_l - \overline{\mathcal{W}}_l \right|^2 + 2 \left| \overline{\mathcal{W}}_l - \mathcal{W}_l \right|^2, \tag{2.12}$$

where

$$\begin{aligned}
\left| \widehat{\mathcal{W}}_l - \overline{\mathcal{W}}_l \right|^2 &\leq \left| -\frac{\beta h}{2} \sum_{p=1}^P L^p B_p(\widehat{\mathcal{W}}_l, t_l) \right|^2 \\
&\leq 3P \frac{(\beta h)^2}{4} \sum_{p=1}^P \left| L^p B_p(\widehat{\mathcal{W}}_l, t_l) - L^p B_p(\overline{\mathcal{W}}_l, t_l) \right|^2 \\
&\quad + 3P \frac{(\beta h)^2}{4} \sum_{p=1}^P \left| L^p B_p(\overline{\mathcal{W}}_l, t_l) - L^{q_1} B_{p_2}(\mathcal{W}_l, t_l) \right|^2 + 3P \frac{(\beta h)^2}{4} \sum_{p=1}^P \left| L^p B_p(\mathcal{W}_l, t_l) \right|^2 \\
&\leq 3P \frac{(\beta h)^2}{4} \ell_1 \left| \widehat{\mathcal{W}}_l - \overline{\mathcal{W}}_l \right|^2 + 3P \frac{(\beta h)^2}{4} \ell_1 \left| \overline{\mathcal{W}}_l - \mathcal{W}_l \right|^2 + 3P \frac{(\beta h)^2}{4} \ell_2 (1 + |\mathcal{W}_l|^2),
\end{aligned}$$

and

$$\begin{aligned}
\left| \overline{\mathcal{W}}_l - \mathcal{W}_l \right|^2 &\leq \left| \alpha h A(\overline{\mathcal{W}}_l, t_l) \right|^2 \leq 2(\alpha h)^2 \left| A(\overline{\mathcal{W}}_l, t_l) - A(\mathcal{W}_l, t_l) \right|^2 + 2(\alpha h)^2 |A(\mathcal{W}_l, t_l)|^2 \\
&\leq 2(\alpha h)^2 \ell_1 \left| \overline{\mathcal{W}}_l - \mathcal{W}_l \right|^2 + 2(\alpha h)^2 \ell_2 (1 + |\mathcal{W}_l|^2).
\end{aligned}$$

From the above inequalities, we obtain

$$\left| \overline{\mathcal{W}}_l - \mathcal{W}_l \right|^2 \leq h^2 \frac{\alpha^2 \ell_2}{1 - (\alpha h)^2 \ell_1} (1 + |\mathcal{W}_l|^2), \tag{2.13}$$

and

$$\left| \widehat{\mathcal{W}}_l - \overline{\mathcal{W}}_l \right| \leq h^2 \frac{3P \ell_2 (\beta)^2}{4 \left(1 - 3P \frac{(\beta h)^2}{4} \ell_1 \right) (1 - (\alpha h)^2 \ell_1)} (1 + |\mathcal{W}_l|^2). \tag{2.14}$$

Combining (2.9)–(2.14), implies that for $(1 - 3P\frac{(\beta h)^2}{4}\ell_1)(1 - (\alpha h)^2\ell_1) > 0$, $\kappa_2 = \frac{3}{2}$. Thus, we can choose in Lemma 2.1 $\kappa_1 = 2$, $\kappa_2 = \frac{3}{2}$ and can prove that the strong order of SSABM scheme is 1.0. \square

3. Linear mean-square stability

In this part of the paper, we consider the scalar linear SDE with a multi-dimensional Wiener process of form

$$dW(t) = \nu W(t)dt + \sum_{p=1}^P \nu_p W(t)d\varpi_p(t), \quad (3.1)$$

where $\nu, \nu_p \in \mathbb{R}$, $W(0) \neq 0 \in \mathbb{R}$. We know that if the coefficient of test Eq (3.1) is satisfied

$$2\nu + \sum_{p=1}^P \nu_p^2 < 0, \quad (3.2)$$

then the trivial solution is asymptotically MS-stable [7, 22], i.e. $\lim_{t \rightarrow \infty} \mathbb{E}[|W(t)|^2] = 0$. If applied a SSABM scheme (1.5) to test Eq (3.1), obtained the difference equation

$$\mathcal{W}_{l+1} = \mathcal{D}(\nu, \nu, h) \mathcal{W}_l, \quad (3.3)$$

with MS-stability function

$$\mathcal{D}(\nu, \nu, h) = 1 + \frac{\nu h + \sqrt{h} \sum_{p=1}^P \nu_p \xi_{p,l} + \sum_{p_1=1}^P \sum_{p_2=1}^P \nu_{p_1} \nu_{p_2} \mathcal{I}_{(p_1, p_2)}}{(1 - \alpha \nu h) \left(1 + \frac{1}{2} h \beta \sum_{p=1}^P \nu_p^2 \right)}. \quad (3.4)$$

Theorem 3.1. *For the test Eq (3.1) with a one-dimensional Wiener process ($P = 1$), the SSABM scheme (1.5) is MS-stable, if and only if $3/2 \leq \alpha + \beta \leq 2$.*

Proof. The MS-stability function of SSABM scheme applied to the test Eq (3.1) with $P = 1$ reads

$$\mathcal{D}(\nu, \nu, h) = 1 + \frac{\nu h + \sqrt{h\nu} \xi_l + h\nu^2(\xi_l^2 - 1)}{(1 - \alpha \nu h) \left(1 + \frac{1}{2} h \beta \nu^2 \right)}. \quad (3.5)$$

The stochastic difference Eq (3.3) with (3.5) is MS-stable if and only if $\mathbb{E}[|\mathcal{D}(\nu, \nu, h)|^2] < 1$. So, we can write

$$a_1 + a_2 + a_3 + a_4 < 1, \quad (3.6)$$

where

$$a_1 = \frac{(\nu h + (1 - \alpha \nu h) \left(1 + \frac{1}{2} h \beta \nu^2 \right) - \frac{1}{2} h \nu^2)^2}{(1 - \alpha \nu h)^2 \left(1 + \frac{1}{2} h \beta \nu^2 \right)^2},$$

$$a_2 = \frac{h \nu^2}{(1 - \alpha \nu h)^2 \left(1 + \frac{1}{2} h \beta \nu^2 \right)^2},$$

$$a_3 = \frac{\frac{3}{4}h^2v^4}{(1 - \alpha vh)^2 \left(1 + \frac{1}{2}h\beta v^2\right)^2},$$

$$a_4 = \frac{h \left(vh + (1 - \alpha vh) \left(1 + \frac{1}{2}h\beta v^2\right) - \frac{1}{2}hv^2 \right) v^2}{(1 - \alpha vh)^2 \left(1 + \frac{1}{2}h\beta v^2\right)^2}.$$

After a little algebra, the condition (3.6) becomes

$$vh((3 - 2\alpha)v + \beta v^2) + 2v + v^2 - \alpha\beta(vh)^2v^2 + \frac{1}{2}h(v^2 - 2v)(v^2 + 2v) < 0. \quad (3.7)$$

It is easy to deduce from (3.7) that $3 - 2\alpha \geq 2\beta$. Also, we know $\alpha + \beta \leq 2$. Thus we complete the proof. \square

Theorem 3.2. For the test Eq (3.1) with commutative noises, SSABM scheme (1.5) is MS-stable, if and only if $3/2 \leq \alpha + \beta \leq 2$.

Proof. The commutativity condition on the diffusion coefficient of test Eq (3.1) reads as $v_{p_1}v_{p_2} = v_{p_2}v_{p_1}$ for all $p_1, p_2 = 1, 2, \dots, P$ [6,25]. Together with the identity $\mathcal{I}_{(p_1, p_2)} + \mathcal{I}_{(p_2, p_1)} = \mathcal{I}_{p_1}\mathcal{I}_{p_2}$ the MS-stability function of SSABM scheme in (3.4) converts to

$$\mathcal{D}(v, v, h) = 1 + \frac{vh + \sqrt{h} \sum_{p=1}^P v_p \xi_{p,l} + \frac{1}{2}h \sum_{p_1=1}^P \sum_{p_2=1}^P v_{p_1} v_{p_2} \xi_{p_1,l} \xi_{p_2,l}}{(1 - \alpha vh) \left(1 + \frac{1}{2}h\beta \sum_{p=1}^P v_p^2\right)}. \quad (3.8)$$

According Theorem 3.1, our scheme is MS-stable for test Eq (3.1) with with commutative noises if and only if

$$v^2h + 2v(1 - \alpha vh) \left(1 + \frac{1}{2}h\beta \sum_{p=1}^P v_p^2\right) + \sum_{p=1}^P v_p^2 + \frac{1}{2} \sum_{p=1}^P v_p^4 + \frac{1}{4}h \left(\sum_{\substack{p_1=1 \\ p_1 \neq p_2}}^P \sum_{p_2=1}^P v_{p_1} v_{p_2} \right)^2 < 0,$$

which, this is equivalent to

$$vh \left((3 - 2\alpha)v + \beta \sum_{p=1}^P v_p^2 \right) - \alpha\beta(vh)^2 \sum_{p=1}^P v_p^2 + 2v + \sum_{p=1}^P v_p^2 + \frac{1}{2}h \left(\sum_{p=1}^P v_p^2 - 2v \right) \left(\sum_{p=1}^P v_p^2 + 2v \right) < 0.$$

It can be easily seen that the above inequality holds if $3 - 2\alpha \geq 2\beta$, which implies the desired assertions. \square

In Figure 1, the behavior of the MS-stability functions of our scheme (3.7) and test Eq (3.2) are compared when $P = 1$. Results of this figure show that the scheme is A-stable if $3/2 \leq \alpha + \beta \leq 2$. Similarly, such results can be obtained for the state of commutative noises in Figure 2.




Figure 1. MS-stability regions for the SSABM scheme (1.5) applied to SDE (3.1) with $P = 1$.




Figure 2. MS-stability regions for the SSABM scheme (1.5) applied to SDE (3.1) with commutative noises.

4. Numerical results

In this section, we compare the convergence rate and the computational performance of our new scheme (1.5) with various α, β , the Milstein and DSSBM (1.3) schemes. In this work, all the computations are performed by using a MATLAB platform.

Example 4.1. Consider the one-dimensional nonlinear SDE,

$$dW(t) = -(a + b^2 W(t))(1 - W^2(t))dt + b(1 - W^2(t))d\varpi(t), \quad W_0 = \frac{1}{2}. \quad (4.1)$$

The exact solution is given by [6]

$$W(t) = \frac{(1 + W_0) \exp(-2at + 2b\varpi(t)) + W_0 - 1}{(1 + W_0) \exp(-2at + 2b\varpi(t)) - W_0 + 1}.$$

The means square errors (MSE) of the SSABM (1.5), Milstein, DSSBM (1.3) (SSTM with $\theta = 1$ [29]), ABM (1.4) and MMA (1.4) schemes can then be obtained in Figure 3. As shown in Figure 3, the SSABM scheme has better than other schemes for $a = b = 0.5$ and if it is $a = 1.0$ and $b = 0.25$ in (4.1), ABM (1.4) has better than other schemes.




Figure 3. MSE estimations of the numerical schemes for nonlinear SDE (4.1).

Example 4.2. Consider the following stiff stochastic system [34]

$$\begin{aligned} dW_1(t) &= 10W_2(t)dt + 0.2(W_1(t) + W_2(t))d\varpi(t), \\ dW_2(t) &= -10W_1(t)dt + 0.2(W_1(t) + W_2(t))d\varpi(t), \\ W_1(t) &= 1, \quad W_2(t) = 0, \quad t \in [0, 20]. \end{aligned} \quad (4.2)$$

By choosing $h = 0.01$, Figure 4 depicts the numerical simulations of the SSABM and Milstein schemes. These figures confirmed that the stability properties of the SSABM scheme are better than the Milstein scheme.




Figure 4. Numerical simulation of the Milstein and SSABM (1.5) schemes for SDE system (4.2).

5. Conclusions

This work has been devoted to the numerical solution to stochastic differential systems (1.1) by the new implicit Milstein scheme. Under given conditions, the strong convergence of the approach has been theoretically investigated and proved that the split-step (α, β) -Milstein scheme has a convergence order of 1.0 in MS sense. Furthermore, the MS-stability of the SSABM scheme has been discussed in this paper. For SDE (3.1) with a single noise term, we show that our scheme is mean-square A-stability for any value $3/2 \leq \alpha + \beta \leq 2$. Also, this result satisfies for SDE (3.1) with multiplicative commutative noise terms. In the last part of this article, the presented scheme has superior efficiency and accuracy to the Milstein and DSSBM [10] schemes.

Conflict of interest

The authors declare no conflicts of interest.

References

1. Z. Korpınar, M. Inc, A. S. Alshomrani, D. Baleanu, The deterministic and stochastic solutions of the Schrödinger equation with time conformable derivative in birefringent fibers, *AIMS Mathematics*, **5** (2020), 2326–2345. <https://doi.org/10.3934/math.2020154>
2. K. Nouri, H. Ranjbar, L. Torkzadeh, The explicit approximation approach to solve stiff chemical langevin equations, *Eur. Phys. J. Plus*, **135** (2020), 758. <https://doi.org/10.1140/epjp/s13360-020-00765-2>
3. K. Nouri, H. Ranjbar, D. Baleanu, L. Torkzadeh, Investigation on ginzburg-landau equation via a tested approach to benchmark stochastic davis-skodje system, *Alexandria Eng. J.*, **60** (2021), 5521–5526. <https://doi.org/10.1016/j.aej.2021.04.040>
4. D. J. Higham, P. E. Kloeden, *An introduction to the numerical simulation of stochastic differential equations*, Society for Industrial and Applied Mathematics, 2021.
5. K. Nouri, F. Fahimi, L. Torkzadeh, D. Baleanu, Stochastic epidemic model of Covid-19 via the reservoir-people transmission network, *Comput. Mater. Contin.*, **72** (2022), 1495–1514. <https://doi.org/10.32604/cmc.2022.024406>
6. P. E. Kloeden, E. Platen, Numerical solution of stochastic differential equations, In: *Applications of mathematics*, Berlin: Springer-Verlag, **23** (1992).
7. X. Mao, *Stochastic differential equations and applications*, Chichester: Horwood Publishing Limited, 2008.
8. K. Nouri, F. Fahimi, L. Torkzadeh, D. Baleanu, Numerical method for pricing discretely monitored double barrier option by orthogonal projection method, *AIMS Mathematics*, **6** (2021), 5750–5761. <https://doi.org/10.3934/math.2021339>
9. G. N. Milstein, Approximate integration of stochastic differential equations, *Theory Prob. Appl.*, **19** (1975), 557–562. <https://doi.org/10.1137/1119062>
10. P. Wang, Z. Liu, Split-step backward balanced Milstein methods for stiff stochastic systems, *Appl. Numer. Math.*, **59** (2009), 1198–1213. <https://doi.org/10.1016/j.apnum.2008.06.001>
11. D. A. Voss, A. Q. M. Khaliq, Split-step Adams-Moulton Milstein methods for systems of stiff stochastic differential equations, *Int. J. Comput. Math.*, **92** (2015), 995–1011. <https://doi.org/10.1080/00207160.2014.915963>
12. F. Jiang, X. Zong, C. Yue, C. Huang, Double-implicit and split two-step Milstein schemes for stochastic differential equations, *Int. J. Comput. Math.*, **93** (2016), 1987–2011. <https://doi.org/10.1080/00207160.2015.1081182>
13. S. S. Ahmad, N. Chandra Parida, S. Raha, The fully implicit stochastic- α method for stiff stochastic differential equations, *J. Comput. Phys.*, **228** (2009), 8263–8282. <https://doi.org/10.1016/j.jcp.2009.08.002>
14. V. Reshniak, A. Q. M. Khaliq, D. A. Voss, G. Zhang, Split-step Milstein methods for multi-channel stiff stochastic differential systems, *Appl. Numer. Math.*, **89** (2015), 1–23. <https://doi.org/10.1016/j.apnum.2014.10.005>

15. T. Tripura, M. Imran, B. Hazra, S. Chakraborty, Change of measure enhanced nearexact euler-maruyama scheme for the solution to nonlinear stochastic dynamical systems, *J. Eng. Mech.*, **148** (2022). [http://doi.org/10.1061/\(ASCE\)EM.1943-7889.0002107](http://doi.org/10.1061/(ASCE)EM.1943-7889.0002107)
16. X. Wang, S. Gan, D. Wang, A family of fully implicit Milstein methods for stiff stochastic differential equations with multiplicative noise, *BIT*, **52** (2012), 741–772. <https://doi.org/10.1007/s10543-012-0370-8>
17. M. S. Semary, M. T. M. Elbarawy, A. F. Fareed, Discrete Temimi-Ansari method for solving a class of stochastic nonlinear differential equations, *AIMS Mathematics*, **7** (2022), 5093–5105. <https://doi.org/10.3934/math.2022283>
18. J. Yao, S. Gan, Stability of the drift-implicit and double-implicit Milstein schemes for nonlinear SDEs, *Appl. Math. Comput.*, **339** (2018), 294–301. <https://doi.org/10.1016/j.amc.2018.07.026>
19. Z. Yin, S. Gan, An improved Milstein method for stiff stochastic differential equations, *Adv. Differ. Equ.*, **369** (2015). <http://doi.org/10.1186/s13662-015-0699-9>
20. R. Kasinathan, R. Kasinathan, D. Baleanu, A. Annamalai, Well posedness of second-order impulsive fractional neutral stochastic differential equations, *AIMS Mathematics*, **6** (2021), 9222–9235. <https://doi.org/10.3934/math.2021536>
21. X. Zong, F. Wu, C. Huang, Convergence and stability of the semi-tamed Euler scheme for stochastic differential equations with non-Lipschitz continuous coefficients, *Appl. Math. Comput.* **228** (2014), 240–250. <https://doi.org/10.1016/j.amc.2013.11.100>
22. Y. Saito, T. Mitsui, Stability analysis of numerical schemes for stochastic differential equations, *SIAM J. Numer. Anal.*, **33** (1996), 2254–2267. <https://doi.org/10.1137/S0036142992228409>
23. Y. Saito, T. Mitsui, Mean-square stability of numerical schemes for stochastic differential systems, *Vietnam J. Math.*, **30** (2002), 551–560.
24. E. Buckwar, C. Kelly, Towards a systematic linear stability analysis of numerical methods for systems of stochastic differential equations, *SIAM J. Numer. Anal.*, **48** (2010), 298–321. <https://doi.org/10.1137/090771843>
25. E. Buckwar, T. Sickenberger, A structural analysis of asymptotic mean-square stability for multi-dimensional linear stochastic differential systems, *Appl. Numer. Math.*, **62** (2012), 842–859. <https://doi.org/10.1016/j.apnum.2012.03.002>
26. D. J. Higham, A-stability and stochastic mean-square stability, *BIT*, **40** (2000), 404–409. <http://doi.org/10.1023/A:1022355410570>
27. A. Tocino, M. J. Senosiain, MS-stability of nonnormal stochastic differential systems, *J. Comput. Appl. Math.*, **379** (2020), 112950. <https://doi.org/10.1016/j.cam.2020.112950>
28. D. J. Higham, X. Mao, L. Szpruch, Convergence, non-negativity and stability of a new Milstein scheme with applications to finance, *Discrete Contin. Dyn. B*, **18** (2013), 2083–2100. <https://doi.org/10.3934/dcdsb.2013.18.2083>
29. X. Zong, F. Wu, G. Xu, Convergence and stability of two classes of theta-Milstein schemes for stochastic differential equations, *J. Comput. Appl. Math.*, **336** (2018), 8–29. <https://doi.org/10.1016/j.cam.2017.12.025>

30. K. Nouri, H. Ranjbar, L. Torkzadeh, Improved Euler-Maruyama method for numerical solution of the Itô stochastic differential systems by composite previous-current-step idea, *Mediterr. J. Math.*, **15** (2018), 140.
31. K. Nouri, H. Ranjbar, L. Torkzadeh, Modified stochastic theta methods by ODEs solvers for stochastic differential equations, *Commun. Nonlinear Sci. Numer. Simul.*, **68** (2019), 336–346. <https://doi.org/10.1016/j.cnsns.2018.08.013>
32. K. Nouri, H. Ranjbar, L. Torkzadeh, Study on split-step Rosenbrock type method for stiff stochastic differential systems, *Int. J. Comput. Math.*, **97** (2020), 816–836. <https://doi.org/10.1080/00207160.2019.1589459>
33. G. N. Milstein, M. V. Tretyakov, *Stochastic numerics for mathematical physics*, Berlin: Springer-Verlag, 2004.
34. S. Singh, S. Raha, Five-stage milstein methods for SDEs, *Int. J. Comput. Math.*, **89** (2012), 760–779. <http://doi.org/10.1080/00207160.2012.657629>



AIMS Press

©2023 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)