



Research article

Strong interaction of Jafari decomposition method with nonlinear fractional-order partial differential equations arising in plasma via the singular and nonsingular kernels

Saima Rashid^{1,*}, Rehana Ashraf² and Fahd Jarad^{3,4,5,*}

¹ Department of Mathematics, Government College University, Faisalabad, Pakistan

² Department of Mathematics, Lahore College Women University, 54000, Lahore, Pakistan

³ Department of Mathematics, Çankaya University, Ankara, Turkey

⁴ Department of Medical Research, China Medical University Hospital, China Medical University, Taichung, Taiwan

⁵ Department of Mathematics, King Abdulaziz University, Jeddah, Saudi Arabia

* **Correspondence:** Email: saimarashid@gcuf.edu.pk, fahd@cankaya.edu.tr.

Abstract: This research utilizes the Jafari transform and the Adomian decomposition method to derive a fascinating explicit pattern for the outcomes of the KdV, mKdV, K(2,2) and K(3,3) models that involve the Caputo fractional derivative operator and the Atangana-Baleanu fractional derivative operator in the Caputo sense. The novel exact-approximate solutions are derived from the formulation of trigonometric, hyperbolic, and exponential function forms. Laser and plasma sciences may benefit from these solutions. It is demonstrated that this approach produces a simple and effective mathematical framework for tackling nonlinear problems. To provide additional context for these ideas, simulations are performed, employing a computationally packaged program to assist in comprehending the implications of solutions.

Keywords: Jafari transform; Adomian decomposition method; KdV and mKdV model; K(2,2) and K(3,3) equations

Mathematics Subject Classification: 46S40, 47H10, 54H25

1. Introduction

Due to its broad relevance and propensity to incorporate many repercussions of actual concerns, the idea of fractional calculus (FC) has garnered considerable prominence in previous decades. Classical calculus has remained a small segment of FC, despite the fact that it can demonstrate

numerous critical challenges and assist us in forecasting the behaviour of intricate occurrences in impulsive integro-differential equations [1], neural networking [2], thermal energy [3], non-Newtonian fluids [4] and heat flux [5]. Despite the fact that innovators offer numerous novel concepts, several aspects should always be deduced in order to guarantee all categories of phenomenon, that will be accomplished by conquering the restrictions posed by mathematicians and scientists. This is highly pertinent when investigating of MHD electro-osmotically flow [6], epidemics [7–9], stability and instability of special functions [10–13], inequalities [14–16] as well as other disciplines. When it tends to arrive to the exploration of repercussions that assist in resolving major difficulties (such as the current global challenges), there is always room for improvements, innovation, creativeness, and extensions in analysis, and so many investigators have inferred provoking outcomes with the assistance of FC, and by incorporating efficacious methodologies with the assistance of underlying FC findings [17–19].

Numerical models investigation and analysis of corresponding features are often a high priority in mathematical modeling when the relevant techniques are implemented. This is certainly pertinent in epidemic research, bifurcation, thermodynamics, electrostatistics modeling, fluid flow, plasma physics, and other fields. Several approaches, including N-solitons [20–23], solitary waves [24–27], Tan-Cot function method [28], Adomian decomposition method [29], homotopy perturbation method [30], q-homotopy analysis method [31], variation iteration method [32], collocation-shooting method [33], G/G' expansion method [34], improved $\tan(\phi(\tau)/2)$ -expansion method [35], Lie symmetry analysis method [36], wavelet method [37] have been employed and refined by researchers to achieve the analytic, semi-analytic, and numerical solution of nonlinear PDEs. The Adomian decomposition method [29, 38] is one of them, and it offers an efficient approach for exact-analytical solutions across a broad and comprehensive domain of specific aspects that simulate real-world issues. This strategy transforms a basic, incredibly straightforward problem into the complicated problem under investigation, and when combined with Adomian components, it offers a tremendous mathematical instrument. Numerous aspects of the Adomian decomposition method have also received considerable focus recently.

In this analysis, we examine a nonlinear framework that explains powerful interactions between interior disturbances in the water. The Korteweg de-Vries (KdV) equations are frequently utilized to illustrate acoustic wave behaviour and its physical relevance. Due to the immense amplitude of lengthy longitudinal waves and prolonged rotating impacts, we employ a modified KdV equation. We now evaluate the following models using the isopycnic surface $\mathcal{W}(x, t)$, which depicts the KdV and modified KdV equations [39], respectively:

$$\frac{\partial \mathcal{W}(x, t)}{\partial t} + a_1 \mathcal{W}(x, t) \frac{\partial \mathcal{W}(x, t)}{\partial x} + a_2 \mathcal{W}^2(x, t) \frac{\partial \mathcal{W}(x, t)}{\partial x} + b_1 \frac{\partial^3 \mathcal{W}(x, t)}{\partial x^3} = 0, \quad (1.1)$$

where a_1 and a_2 signifies the quadratic and cubic non-linear coefficients, respectively. Also, the coefficient of small-scale dispersion is denoted by b_1 . Here, a_1 and a_1 presents the proportional factors associating in the aforesaid equation, and is due to the nonlinear hydrodynamic system, and it appears classically, see [40, 41].

Furthermore, we compute the exact-analytical solution of nonlinear dispersive equations $K(n, n)$:

$$\frac{\partial \mathcal{W}(x, t)}{\partial t} + \frac{\partial}{\partial x}(\mathcal{W}^n) + \frac{\partial^3}{\partial x^3}(\mathcal{W}^n) = 0, \quad n > 1. \quad (1.2)$$

Equation (1.2) is the evolutionary model for compactons. Compactons are characterized as solitons with bounded wave lengths or solitons without exponential tails in solitary wave theory (Rosenau and Hyma, 1993). Compactons are formed by the intricate coupling of nonlinear convection $\frac{\partial}{\partial x}(\mathcal{W}^n)$ and nonlinear dispersion $\frac{\partial^3}{\partial x^3}(\mathcal{W}^n)$ in (1.2).

Amidst Gorge Adomian's massive boost in 1980, the Adomian decomposition method introduced a well-noted terminology. It has been intensively implemented for a diverse set of nonlinear PDEs, for instance, the Korteweg-De Vries model [42], Fisher's model [43], Zakharov-Kuznetsov equation [44] and so on. The ADM was determined to be significantly related to a variety of integral transforms, including Laplace, Swai, Mohand, Aboodh, Elzaki, and others. Humanity is continuously striving to improve performance and minimizing the method's intricacy through invention, modernity, and experimentation. In connection with this, Jafari [45] propounded a well-known integral transform which is known to generalized integral transform. The dominant feature of this transformation is that it has the ability to recapture several existing transformations, see Remark 1.

Motivated by the above propensity, we aim to establish a semi-analytical approach by mingling the Jafari transform with the Adomian decomposition method, namely the Jafari decomposition method (JDM). With the assistance of fractional derivative operators, we constructed the approximate-analytical solutions for KdV, MKdV, K(2,2) and K(3,3). The suggested methodology helps us increase flexibility in determining the initial conditions, and its novelty is that it has a straightforward solution technique. This approach is straightforward and encompasses all of JDM accomplishments, as well as encouraging several scholars to investigate a broad spectrum of applications and processes. The tool to overcome computational complexity without any constraints, perturbations, or transformations from nonlinear to linear, or partial to ordinary differential equations, is the distinctive characteristic of the proposed approach. Furthermore, it is connected to factors that are extremely useful in bringing the findings to a favourable conclusion. It also is coupled to well-posed transformation, which tries to diminish the technique's intricacy while increasing its application and dependability.

2. Preliminaries

In this section, we evoke some essential concepts, notions, and definitions concerning fractional derivative operators depending on power and Mittag-Leffer as a kernel, along with the detailed consequences of the Jafari transform.

Definition 2.1. ([17]) The Caputo fractional derivative (CFD) is described as follows:

$${}^c_0\mathbf{D}_t^\lambda = \begin{cases} \frac{1}{\Gamma(r-\lambda)} \int_0^t \frac{\mathcal{W}^{(r)}(x)}{(t-x)^{\lambda+1-r}} dx, & r-1 < \lambda < r, \\ \frac{d^r}{dt^r} \mathcal{W}(t), & \lambda = r. \end{cases} \quad (2.1)$$

Definition 2.2. ([18]) The Atangana-Baleanu fractional derivative operator in the Caputo form (ABC) is stated as follows:

$${}^{ABC}_{\eta_1} \mathbf{D}_t^\lambda(\mathcal{W}(t)) = \frac{\mathbb{A}(\lambda)}{1-\lambda} \int_{\eta_1}^t \mathcal{W}'(t) E_\lambda \left[-\frac{\lambda(t-x)^\lambda}{1-\lambda} \right] dx, \quad (2.2)$$

where $\mathcal{W} \in \mathcal{H}^1(a_1, a_2)$ (Sobolev space), $a_1 < a_2$, $\lambda \in [0, 1]$ and $\mathbb{A}(\lambda)$ signifies a normalization function as $\mathbb{A}(\lambda) = \mathbb{A}(0) = \mathbb{A}(1) = 1$.

Definition 2.3. ([18]) The fractional integral of the ABC-operator is described as follows:

$${}_{\eta_1}^{ABC} I_t^\lambda(\mathcal{W}(t)) = \frac{1-\lambda}{\mathbb{A}(\lambda)} \mathcal{W}(t) + \frac{\lambda}{\Gamma(\lambda)\mathbb{A}(\lambda)} \int_{\eta_1}^t \mathcal{W}(x)(t-x)^{\lambda-1} dx. \quad (2.3)$$

Definition 2.4. ([45]) Consider an integrable mapping $\mathcal{W}(t)$ defined on a set \mathcal{P} , then

$$\mathcal{P} = \left\{ \mathcal{W}(t) : \exists M > 0, \kappa > 0, |\mathcal{W}(t)| < M \exp(\kappa t), \text{ if } t \geq 0 \right\}. \quad (2.4)$$

Definition 2.5. ([45]) Suppose the mappings $\phi(s), \psi(s) : \mathbb{R}^+ \mapsto \mathbb{R}^+$ such that $\phi(s) \neq 0 \forall s \in \mathbb{R}^+$. The Jafari transform of the mapping $\mathcal{W}(t)$ presented by $\mathbf{Q}(s)$ is described as

$$\mathbb{J}\{\mathcal{W}(t), s\} = \mathbf{Q}(s) = \phi(s) \int_0^\infty \mathcal{W}(t) \exp(-\psi(s)t) dt. \quad (2.5)$$

Theorem 2.6. ([45]) (Convolution property). For Jafari transform, the subsequent holds true:

$$\mathbb{J}\{\mathcal{W}_1 * \mathcal{W}_2\} = \frac{1}{\phi(s)} \mathbf{Q}_1(s) * \mathbf{Q}_2(s). \quad (2.6)$$

Definition 2.7. The Jafari transform of the CFD operator is stated as follows:

$$\mathbb{J}\left\{{}_0^c \mathbf{D}_t^\lambda(\mathcal{W}(t)), s\right\} = \psi^\lambda(s) \mathbf{Q}(s) - \phi(s) \sum_{\kappa=0}^{\lambda-1} \psi^{\lambda-\kappa-1}(s) \mathcal{W}^{(\kappa)}(0), \quad r-1 < \lambda < r, \phi, \psi > 0. \quad (2.7)$$

Remark 1. Definition 2.7 leads to the following conclusions:

- 1) Taking $\phi(s) = 1$ and $\psi(s) = s$, then we acquire the Laplace transform [46].
- 2) Taking $\phi(s) = \frac{1}{s}$ and $\psi(s) = \frac{1}{s}$, then we acquire the α -Laplace transform [47].
- 3) Taking $\phi(s) = \frac{1}{s}$ and $\psi(s) = \frac{1}{s}$, then we acquire the Sumudu transform [48].
- 4) Taking $\phi(s) = \frac{1}{s}$ and $\psi(s) = 1$, then we acquire the Aboodh transform [49].
- 5) Taking $\phi(s) = s$ and $\psi(s) = s^2$, then we acquire the Pourreza transform [50, 51].
- 6) Taking $\phi(s) = s$ and $\psi(s) = \frac{1}{s}$, then we acquire the Elzaki transform [52].
- 7) Taking $\phi(s) = \mathbf{u}_2$ and $\psi(s) = \frac{s}{\mathbf{u}_2}$, then we acquire the Natural transform [53].
- 8) Taking $\phi(s) = s^2$ and $\psi(s) = s$, then we acquire the Mohand transform [54].
- 9) Taking $\phi(s) = \frac{1}{s^2}$ and $\psi(s) = \frac{1}{s}$, then we acquire the Swai transform [55].
- 10) Taking $\phi(s) = 1$ and $\psi(s) = \frac{1}{s}$, then we get the Kamal transform [56].
- 11) Taking $\phi(s) = s^\alpha$ and $\psi(s) = \frac{1}{s}$, then we acquire the G_- transform [57, 58].

Definition 2.8. ([59]) The Jafari transform of the ABC fractional derivative operator is described as:

$$\mathbb{J}\left\{{}_0^{ABC} \mathbf{D}_t^\lambda(\mathcal{W}(t)), s\right\}(\lambda) = \frac{\mathbb{A}(\lambda)\psi^\lambda(s)}{\lambda + (1-\lambda)\psi^\lambda(s)} \left(\mathbf{Q}(s) - \frac{\phi(s)}{\psi(s)} \mathcal{W}(0) \right). \quad (2.8)$$

Remark 2. Definition 2.8 leads to the following conclusions:

- 1) Taking $\phi(s) = 1$ and $\psi(s) = s$, then we acquire the Laplace transform of ABC fractional derivative operator [60, 61].
- 2) Taking $\phi(s) = s$ and $\psi(s) = \frac{1}{s}$, then we acquire the Elzaki transform of ABC fractional derivative operator [62].
- 3) Taking $\phi(s) = \psi(s) = \frac{1}{s}$, then we get the Sumudu transform of ABC fractional derivative operator [63].
- 4) Taking $\phi(s) = 1$ and $\psi(s) = s/\mathbf{u}_2$, then we get the Shehu transform of ABC fractional derivative operator [63].

Definition 2.9. ([64]) The Mittag-Leffler function for single parameter is described as

$$E_\lambda(z) = \sum_{\kappa=0}^{\infty} \frac{z_1^\kappa}{\Gamma(\kappa\lambda + 1)}, \quad \lambda, z_1 \in \mathbb{C}, \quad \Re(\lambda) \geq 0. \quad (2.9)$$

3. Description of the Jafari decomposition method involving singular and nonsingular kernels

Consider the generic fractional form of PDE:

$$\mathbf{D}_t^\lambda \mathcal{W}(x, t) + \mathfrak{L}\mathcal{W}(x, t) + \mathfrak{N}\mathcal{W}(x, t) = \mathcal{F}(x, t), \quad t > 0, \quad 0 < \lambda \leq 1 \quad (3.1)$$

with ICs

$$\mathcal{W}(x, 0) = \mathcal{G}(x), \quad (3.2)$$

where $\mathbf{D}_t^\lambda = \frac{\partial^\lambda \mathcal{W}(x, t)}{\partial t^\lambda}$ symbolizes the Caputo and ABC fractional derivative of order $\lambda \in (0, 1]$ while \mathfrak{L} and \mathfrak{N} denotes the linear and nonlinear factors, respectively. Also, $\mathcal{F}(x, t)$ represents the source term. Taking into account the Jafari transform to (3.1), and we acquire

$$\mathbb{J}[\mathbf{D}_t^\lambda \mathcal{W}(x, t) + \mathfrak{L}\mathcal{W}(x, t) + \mathfrak{N}\mathcal{W}(x, t)] = \mathbb{J}[\mathcal{F}(x, t)].$$

Firstly, applying the differentiation rule of Jafari transform with respect to CFD, then we apply the ABC fractional derivative operator as follows:

$$\psi^\lambda(s)\mathcal{U}(x, s) = \phi(s) \sum_{\kappa=0}^{\ell-1} \psi^{\lambda-1-\kappa}(s)\mathcal{W}^{(\kappa)}(0) + \mathbb{J}[\mathfrak{L}\mathcal{W}(x, t) + \mathfrak{N}\mathcal{W}(x, t)] + \mathbb{J}[\mathcal{F}(x, t)], \quad (3.3)$$

and

$$\frac{\psi^\lambda(s)\mathbb{A}(\lambda)}{\lambda + (1-\lambda)\psi^\lambda(s)}\mathcal{U}(x, s) = \frac{\phi(s)}{\psi(s)} \frac{\psi^\lambda(s)\mathbb{A}(\lambda)}{\lambda + (1-\lambda)\psi^\lambda(s)}\mathcal{W}(0) + \mathbb{J}[\mathfrak{L}\mathcal{W}(x, t) + \mathfrak{N}\mathcal{W}(x, t)] + \mathbb{J}[\mathcal{F}(x, t)]. \quad (3.4)$$

The inverse Jafari transform of (3.3) and (3.4) yields

$$\mathcal{W}(x, t) = \mathbb{J}^{-1} \left[\phi(s) \sum_{\kappa=0}^{\ell-1} \psi(s)^{\lambda-\kappa-1} \mathcal{W}^{(\kappa)}(0) + \frac{1}{\psi^\lambda(s)} \mathbb{J}[\mathcal{F}(x, t)] \right] - \mathbb{J}^{-1} \left[\frac{1}{\psi^\lambda(s)} \mathbb{J}[\mathfrak{L}\mathcal{W}(x, t) + \mathfrak{N}\mathcal{W}(x, t)] \right]. \quad (3.5)$$

and

$$\mathcal{W}(x, t) = \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(0) + \frac{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})}{\psi^\lambda(\mathfrak{s})\mathbb{A}(\lambda)} \mathbb{J}[\mathcal{F}(x, t)] \right] - \mathbb{J}^{-1} \left[\frac{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})}{\psi^\lambda(\mathfrak{s})\mathbb{A}(\lambda)} \mathbb{J}[\mathfrak{L}\mathcal{W}(x, t) + \mathfrak{N}\mathcal{W}(x, t)] \right]. \quad (3.6)$$

The generalized decomposition method solution $\mathcal{W}(x, t)$ is represented by the following infinite series

$$\mathcal{W}(x, t) = \sum_{\ell=0}^{\infty} \mathcal{W}_\ell(x, t). \quad (3.7)$$

Thus, the nonlinear term $\mathfrak{N}(x, t)$ can be evaluated by the Adomian decomposition method prescribed as

$$\mathfrak{N}\mathcal{W}(x, t) = \sum_{\ell=0}^{\infty} \tilde{A}_\ell(\mathcal{W}_0, \mathcal{W}_1, \dots), \quad \ell = 0, 1, \dots, \quad (3.8)$$

where

$$\tilde{A}_\ell(\mathcal{W}_0, \mathcal{W}_1, \dots) = \frac{1}{\ell!} \left[\frac{d^\ell}{d\mathfrak{s}^\ell} \mathfrak{N} \left(\sum_{j=0}^{\infty} \mathfrak{s}^j \mathcal{W}_j \right) \right]_{\mathfrak{s}=0}, \quad \ell > 0.$$

Inserting (3.7) and (3.8) into (3.5) and (3.6), respectively, we have

$$\sum_{\ell=0}^{\infty} \mathcal{W}_\ell(x, t) = \mathcal{G}(x) + \tilde{\mathcal{G}}(x) - \mathbb{J}^{-1} \left[\frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J}[\mathfrak{L}\mathcal{W}(x, t) + \sum_{\ell=0}^{\infty} \tilde{A}_\ell] \right] \quad (3.9)$$

and

$$\sum_{\ell=0}^{\infty} \mathcal{W}_\ell(x, t) = \mathcal{G}(x) + \tilde{\mathcal{G}}(x) - \mathbb{J}^{-1} \left[\frac{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})}{\mathbb{A}(\lambda)\psi^\lambda(\mathfrak{s})} \mathbb{J}[\mathfrak{L}\mathcal{W}(x, t) + \sum_{\ell=0}^{\infty} \tilde{A}_\ell] \right]. \quad (3.10)$$

Consequently, the recursive technique for (3.9) and (3.10) are established as:

$$\begin{aligned} \mathcal{W}_0(x, t) &= \mathcal{G}(x) + \tilde{\mathcal{G}}(x), \quad \ell = 0, \\ \mathcal{W}_{\ell+1}(x, t) &= -\mathbb{J}^{-1} \left[\frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J}[\mathfrak{L}(\mathcal{W}_\ell(x, t)) + \sum_{\ell=0}^{\infty} \tilde{A}_\ell] \right], \quad \ell \geq 1, \\ \mathcal{W}_{\ell+1}(x, t) &= -\mathbb{J}^{-1} \left[\frac{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})}{\mathbb{A}(\lambda)\psi^\lambda(\mathfrak{s})} \mathbb{J}[\mathfrak{L}(\mathcal{W}_\ell(x, t)) + \sum_{\ell=0}^{\infty} \tilde{A}_\ell] \right], \quad \ell \geq 1. \end{aligned} \quad (3.11)$$

4. Physical interpretation of Jafari decomposition method

In what follows, we present the various kinds of partial differential equations with the CFD and AB-fractional derivative operators, respectively.

4.1. Fractional KdV equation

Example 4.1. Assume that the time-fractional KdV equation

$$\mathbf{D}_t^\lambda \mathcal{W}(x, t) - 6\mathcal{W}\mathcal{W}_x(x, t) + \mathcal{W}_{xxx}(x, t) = 0, \quad (4.1)$$

with IC:

$$\mathcal{W}_0(x, 0) = -2 \frac{\sigma^2 \exp(\sigma x)}{(1 + \exp(\sigma x))^2}. \quad (4.2)$$

Proof. Foremost, we provide the solution of (4.1) in two general cases.

Case I. Firstly, we apply the Caputo fractional derivative operator coupled with the Jafari transform and Adomian decomposition method. Applying the Jafari transform to (4.1).

$$\psi^\lambda(\mathfrak{s})\mathcal{U}(x, \mathfrak{s}) - \phi(\mathfrak{s}) \sum_{\kappa=0}^{m-1} \psi^{\lambda-\kappa-1}(\mathfrak{s})\mathcal{W}^{(\kappa)}(0) = \mathbb{J} \left[6\mathcal{W}\mathcal{W}_x(x, t) - \mathcal{W}_{xxx}(x, t) \right]. \quad (4.3)$$

Taking into consideration the IC given in (4.2), we have

$$\mathcal{U}(x, \mathfrak{s}) = \frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})}\mathcal{W}(x, 0) + \frac{1}{\psi^\lambda(\mathfrak{s})}\mathbb{J} \left[6\mathcal{W}\mathcal{W}_x(x, t) - \mathcal{W}_{xxx}(x, t) \right].$$

Employing the inverse Jafari transform, we obtain

$$\mathcal{W}(x, t) = \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})}\mathcal{W}(x, 0) + \frac{1}{\psi^\lambda(\mathfrak{s})}\mathbb{J} \left[6\mathcal{W}\mathcal{W}_x(x, t) - \mathcal{W}_{xxx}(x, t) \right] \right]. \quad (4.4)$$

Thanks to the JDM, we find

$$\begin{aligned} \mathcal{W}_0(x, t) &= \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})}\mathcal{W}(x, 0) \right] = -2\mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \frac{\sigma^2 \exp(\sigma x)}{(1 + \exp(\sigma x))^2} \right] \\ &= -2 \frac{\sigma^2 \exp(\sigma x)}{(1 + \exp(\sigma x))^2}. \end{aligned}$$

Here, we surmise that the unknown function $\mathcal{W}(x, t)$ can be written by an infinite series of the form

$$\mathcal{W}(x, t) = \sum_{\ell=0}^{\infty} \mathcal{W}_\ell(x, t).$$

Also, the non-linearity $\mathcal{F}(\mathcal{W})$ can be decomposed by an infinite series of polynomials represented by

$$\mathcal{F}(\mathcal{W}) = \mathcal{W}\mathcal{W}_x = \sum_{\ell=0}^{\infty} \mathcal{A}_\ell,$$

where $\mathcal{W}_\ell(x, t)$ will be evaluated recurrently, and \mathcal{A}_ℓ is the so-called polynomial of $\mathcal{W}_0, \mathcal{W}_1, \dots, \mathcal{W}_\ell$ established by [65].

$$\sum_{\ell=0}^{\infty} \mathcal{W}_{\ell+1}(x, t) = \mathbb{J}^{-1} \left[\frac{1}{\psi^\lambda(\mathfrak{s})}\mathbb{J} \left[6 \sum_{\ell=0}^{\infty} (\mathcal{A})_\ell + \sum_{\ell=0}^{\infty} (\mathcal{W}_{xxx})_\ell \right] \right], \quad \ell = 0, 1, 2, \dots$$

The first few Adomian polynomials are presented as follows:

$$\mathcal{A}_\ell(\mathcal{W}\mathcal{W}_x) = \begin{cases} \mathcal{W}_0\mathcal{W}_{0x}, & \ell = 0, \\ \mathcal{W}_{0x}\mathcal{W}_1 + \mathcal{W}_{1x}\mathcal{W}_0, & \ell = 1, \\ \mathcal{W}_2\mathcal{W}_{0x} + \mathcal{W}_1\mathcal{W}_{1x} + \mathcal{W}_0\mathcal{W}_{2x}, & \ell = 2, \end{cases} \quad (4.5)$$

For $\ell = 0, 1, 2, 3, \dots$

$$\begin{aligned} \mathcal{W}_1(x, t) &= \mathbb{J}^{-1} \left[\frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[6\mathcal{A}_0 + \mathcal{W}_{0xxx} \right] \right] \\ &= -2 \frac{\sigma^5 \exp(\sigma x) (\exp(\sigma x) - 1)}{(1 + \exp(\sigma x_1))^3} \frac{t^\lambda}{\Gamma(\lambda + 1)}, \\ \mathcal{W}_2(x, t) &= \mathbb{J}^{-1} \left[\frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[6\mathcal{A}_1 + \mathcal{W}_{1xxx} \right] \right] \\ &= -2 \frac{\sigma^8 \exp(\sigma x) (\exp(2\sigma x) - 4 \exp(\sigma x) + 1)}{(1 + \exp(\sigma x_1))^4} \frac{t^{2\lambda}}{\Gamma(2\lambda + 1)}, \\ &\vdots \end{aligned}$$

The approximate solution for Example 4.1 is expressed as:

$$\begin{aligned} \mathcal{W}(x, t) &= \mathcal{W}_0(x, t) + \mathcal{W}_1(x, t) + \mathcal{W}_2(x, t) + \mathcal{W}_3(x, t) + \dots, \\ &= -2 \frac{\sigma^2 \exp(\sigma x)}{(1 + \exp(\sigma x))^2} - 2 \frac{\sigma^5 \exp(\sigma x) (\exp(\sigma x) - 1)}{(1 + \exp(\sigma x_1))^3} \frac{t^\lambda}{\Gamma(\lambda + 1)} \\ &\quad - 2 \frac{\sigma^8 \exp(\sigma x) (\exp(2\sigma x) - 4 \exp(\sigma x) + 1)}{(1 + \exp(\sigma x_1))^4} \frac{t^{2\lambda}}{\Gamma(2\lambda + 1)} + \dots \end{aligned} \quad (4.6)$$

Case II. Here, we surmise ABC fractional derivative operator coupled with the Jafari transform and Adomian decomposition method. Applying the Jafari transform for Example 4.1.

$$\frac{\psi^\lambda(\mathfrak{s})\mathbb{A}(\lambda)}{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})} \mathcal{U}(x, \mathfrak{s}) - \phi(\mathfrak{s}) \sum_{\kappa=0}^{m-1} \psi^{\lambda-\kappa-1}(\mathfrak{s}) \mathcal{W}^{(\kappa)}(0) = \mathbb{J} \left[6\mathcal{W}\mathcal{W}_x(x, t) - \mathcal{W}_{xxx}(x, t) \right]. \quad (4.7)$$

Taking into consideration the IC given in (4.2), we have

$$\mathcal{U}(x, \mathfrak{s}) = \frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) + \frac{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})}{\psi^\lambda(\mathfrak{s})\mathbb{A}(\lambda)} \mathbb{J} \left[6\mathcal{W}\mathcal{W}_x(x, t) - \mathcal{W}_{xxx}(x, t) \right].$$

Employing the inverse Jafari transform, we obtain

$$\mathcal{W}(x, t) = \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) + \frac{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})}{\psi^\lambda(\mathfrak{s})\mathbb{A}(\lambda)} \mathbb{J} \left[6\mathcal{W}\mathcal{W}_x(x, t) - \mathcal{W}_{xxx}(x, t) \right] \right]. \quad (4.8)$$

Thanks to the JDM, we find

$$\mathcal{W}_0(x, t) = \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) \right] = -2 \mathbb{J}^{-1} \left[\frac{\psi(\mathfrak{s}_1)}{\phi(\mathfrak{s})} \frac{\sigma^2 \exp(\sigma x)}{(1 + \exp(\sigma x))^2} \right]$$

$$= -2 \frac{\sigma^2 \exp(\sigma x)}{(1 + \exp(\sigma x))^2}.$$

Here, we surmise that the unknown function $\mathcal{W}(x, t)$ can be written by an infinite series of the form

$$\mathcal{W}(x, t) = \sum_{\ell=0}^{\infty} \mathcal{W}_{\ell}(x, t).$$

Also, the non-linearity $\mathcal{F}_1(\mathcal{W})$ can be decomposed by an infinite series of polynomials represented by

$$\mathcal{F}_1(\mathcal{W}) = \mathcal{W}\mathcal{W}_x = \sum_{\ell=0}^{\infty} \mathcal{A}_{\ell},$$

where $\mathcal{W}_{\ell}(x, t)$ will be evaluated recurrently, and \mathcal{A}_{ℓ} is the so-called polynomial of $\mathcal{W}_0, \mathcal{W}_1, \dots, \mathcal{W}_{\ell}$ defined in (4.5). Then, we have

For $\ell = 0, 1, 2, 3, \dots$

$$\begin{aligned} \mathcal{W}_1(x, t) &= \mathbb{J}^{-1} \left[\frac{\lambda + (1 - \lambda)\psi^{\lambda}(\mathfrak{s})}{\psi^{\lambda}(\mathfrak{s})\mathbb{A}(\lambda)} \mathbb{J} \left[6\mathcal{A}_0 + \mathcal{W}_{0xxx} \right] \right] \\ &= -\frac{2}{\mathbb{A}(\lambda)} \frac{\sigma^5 \exp(\sigma x)(\exp(\sigma x) - 1)}{(1 + \exp(\sigma x_1))^3} \left[\frac{\lambda t^{\lambda}}{\Gamma(\lambda + 1)} + (1 - \lambda) \right], \\ \mathcal{W}_2(x, t) &= \mathbb{J}^{-1} \left[\frac{\lambda + (1 - \lambda)\psi^{\lambda}(\mathfrak{s})}{\psi^{\lambda}(\mathfrak{s})\mathbb{A}(\lambda)} \mathbb{J} \left[6\mathcal{A}_1 + \mathcal{W}_{1xxx} \right] \right] \\ &= -\frac{2}{\mathbb{A}^2(\lambda)} \frac{\sigma^8 \exp(\sigma x)(\exp(2\sigma x) - 4 \exp(\sigma x) + 1)}{(1 + \exp(\sigma x_1))^4} \left[\frac{\lambda^2 t^{2\lambda}}{\Gamma(2\lambda + 1)} + 2\lambda(1 - \lambda) \frac{t^{\lambda}}{\Gamma(\lambda + 1)} + (1 - \lambda)^2 \right], \\ &\vdots \end{aligned}$$

The approximate solution for Example 4.1 is expressed as:

$$\begin{aligned} \mathcal{W}(x, t) &= \mathcal{W}_0(x, t) + \mathcal{W}_1(x, t) + \mathcal{W}_2(x, t) + \mathcal{W}_3(x, t) + \dots, \\ &= -2 \frac{\sigma^2 \exp(\sigma x)}{(1 + \exp(\sigma x))^2} - 2 \frac{\sigma^5 \exp(\sigma x)(\exp(\sigma x) - 1)}{\mathbb{A}(\lambda)(1 + \exp(\sigma x_1))^3} \left[\frac{\lambda t^{\lambda}}{\Gamma(\lambda + 1)} + (1 - \lambda) \right] \\ &\quad - 2 \frac{\sigma^8 \exp(\sigma x)(\exp(2\sigma x) - 4 \exp(\sigma x) + 1)}{\mathbb{A}^2(\lambda)(1 + \exp(\sigma x_1))^4} \left[\frac{\lambda^2 t^{2\lambda}}{\Gamma(2\lambda + 1)} + 2\lambda(1 - \lambda) \frac{t^{\lambda}}{\Gamma(\lambda + 1)} + (1 - \lambda)^2 \right] \quad (4.9) \end{aligned}$$

For $\lambda = 1$, we obtained the exact solution of Example 4.1 as

$$\mathcal{W}(x, t) = -\frac{\sigma^2}{2} \sec h^2 \frac{\sigma}{2} (x - \sigma^2 t).$$

Figure 1 shows the evolutionary outcomes for the explicit and approximate solutions of Example 4.1 for the particular instance $\lambda = 1$. The result generated by the proposed technique is remarkably similar to the exact solution, as shown in Figure 1.

Then, by considering only the first few elements of the linear equations features are integrated, we can deduce that we have accomplished a reasonable estimation with the numerical solutions of

the problem. It is obvious that by adding additional components to the decomposition series (4.6) and (4.9), the cumulative error can be diminished.

Analogously, we demonstrate the two-dimensional view of the change in fractional values of the order. We depict the response in Figure 2. It is remarkable that pairwise collisions of particle-like phenomena (including solitary waves and breathers) are fundamental mechanisms in the production of condensed soliton gas dynamics. Deep water waves, shallow groundwater waves, internally waves in a segmented sea, and fibre optics are all manifestations of these waves. \square




Figure 1. Three-dimensional illustration of the exact and approximate solution of Example 4.1 when $\lambda = 1$, and $\sigma = 0.05$.




Figure 2. Three-dimensional illustration of the absolute error plot and 2D-approximations of Example 4.1 when $\lambda = 1$, $\sigma = 0.05 = 0$, and $t = 0.5$.

Remark 3. It is remarkable that equivalent version of the KdV equation is presented as

$$\mathbf{D}_t^\lambda \mathcal{W}(x, t) + 6\mathcal{W}\mathcal{W}_x(x, t) + \mathcal{W}_{xxx}(x, t) = 0,$$

with IC:

$$\mathcal{W}_0(x, 0) = -2 \frac{\sigma^2 \exp(\sigma x)}{(1 + \exp(\sigma x))^2}.$$

has the solitary wave solution, when $\lambda = 1$, then

$$\mathcal{W}(x, t) = \frac{\sigma^2}{2} \operatorname{sech}^2 \frac{\sigma}{2} (x - \sigma^2 t).$$

4.2. Fractional modified KdV equation

Example 4.2. Assume that the time-fractional modified KdV equation

$$\mathbf{D}_t^\lambda \mathcal{W}(x, t) + 6\mathcal{W}^2 \mathcal{W}_x(x, t) + \mathcal{W}_{xxx}(x, t) = 0, \quad (4.10)$$

with IC:

$$\mathcal{W}_0(x, 0) = 2 \frac{\sigma \exp(\sigma x)}{1 + \exp(2\sigma x)}. \quad (4.11)$$

Proof. Foremost, we provide the solution of (4.10) in two general cases.

Case I. Firstly, we apply the Caputo fractional derivative operator coupled with the Jafari transform and Adomian decomposition method.

Applying the Jafari transform to (4.10).

$$\psi^\lambda(\mathfrak{s}) \mathcal{U}(x, \mathfrak{s}) - \phi(\mathfrak{s}) \sum_{\kappa=0}^{m-1} \psi^{\lambda-\kappa-1}(\mathfrak{s}) \mathcal{W}^{(\kappa)}(0) = -\mathbb{J} \left[6\mathcal{W}^2 \mathcal{W}_x(x, t) + \mathcal{W}_{xxx}(x, t) \right]. \quad (4.12)$$

Taking into consideration the IC given in (4.11), we have

$$\mathcal{U}(x, \mathfrak{s}) = \frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) - \frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[6\mathcal{W}^2 \mathcal{W}_x(x, t) + \mathcal{W}_{xxx}(x, t) \right].$$

Employing the inverse Jafari transform, we obtain

$$\mathcal{W}(x, t) = \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) - \frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[6\mathcal{W}^2 \mathcal{W}_x(x, t) + \mathcal{W}_{xxx}(x, t) \right] \right]. \quad (4.13)$$

Thanks to the JDM, we find

$$\begin{aligned} \mathcal{W}_0(x, t) &= \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) \right] = 2 \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \frac{\sigma \exp(\sigma x)}{1 + \exp(2\sigma x)} \right] \\ &= 2 \frac{\sigma \exp(\sigma x)}{1 + \exp(2\sigma x)}. \end{aligned}$$

Here, we surmise that the unknown function $\mathcal{W}(x, t)$ can be written by an infinite series of the form

$$\mathcal{W}(x, t) = \sum_{\ell=0}^{\infty} \mathcal{W}_{\ell}(x, t).$$

Also, the non-linearity $\mathcal{F}(\mathcal{W})$ can be decomposed by an infinite series of polynomials represented by

$$\mathcal{F}(\mathcal{W}) = \mathcal{W}^2 \mathcal{W}_x = \sum_{\ell=0}^{\infty} \mathcal{B}_{\ell},$$

where $\mathcal{W}_{\ell}(x, t)$ will be evaluated recurrently, and \mathcal{B}_{ℓ} is the so-called polynomial of $\mathcal{W}_0, \mathcal{W}_1, \dots, \mathcal{W}_{\ell}$ established by [65].

$$\sum_{\ell=0}^{\infty} \mathcal{W}_{\ell+1}(x, t) = -\mathbb{J}^{-1} \left[\frac{1}{\psi^{\lambda}(\xi)} \mathbb{J} \left[6 \sum_{\ell=0}^{\infty} (\mathcal{B})_{\ell} + \sum_{\ell=0}^{\infty} (\mathcal{W}_{xxx})_{\ell} \right] \right], \quad \ell = 0, 1, 2, \dots$$

The first few Adomian polynomials are presented as follows:

$$\mathcal{B}_{\ell}(\mathcal{W}^2 \mathcal{W}_x) = \begin{cases} \mathcal{W}_0^2 \mathcal{W}_{0x}, & \ell = 0, \\ \mathcal{W}_{0x}(2\mathcal{W}_0 \mathcal{W}_1) + \mathcal{W}_{1x} \mathcal{W}_0^2, & \ell = 1, \\ (2\mathcal{W}_2 \mathcal{W}_0 + \mathcal{W}_1^2) \mathcal{W}_{0x} + (2\mathcal{W}_0 \mathcal{W}_1) \mathcal{W}_{1x} + \mathcal{W}_0^2 \mathcal{W}_{2x}, & \ell = 2, \end{cases} \quad (4.14)$$

For $\ell = 0, 1, 2, 3, \dots$

$$\begin{aligned} \mathcal{W}_1(x, t) &= -\mathbb{J}^{-1} \left[\frac{1}{\psi^{\lambda}(\xi)} \mathbb{J} \left[6\mathcal{B}_0 + \mathcal{W}_{0xxx} \right] \right] \\ &= -2 \frac{\sigma^4 \exp(\sigma x) (1 - \exp(2\sigma x))}{(1 + \exp(2\sigma x_1))^2} \frac{t^{\lambda}}{\Gamma(\lambda + 1)}, \\ \mathcal{W}_2(x, t) &= -\mathbb{J}^{-1} \left[\frac{1}{\psi^{\lambda}(\xi)} \mathbb{J} \left[6\mathcal{B}_1 + \mathcal{W}_{1xxx} \right] \right] \\ &= 2 \frac{\sigma^7 \exp(\sigma x) (1 - 6 \exp(2\sigma x) + \exp(4\sigma x))}{(1 + \exp(2\sigma x_1))^3} \frac{t^{2\lambda}}{\Gamma(2\lambda + 1)}, \\ &\vdots \end{aligned}$$

The approximate solution for Example 4.2 is expressed as:

$$\begin{aligned} \mathcal{W}(x, t) &= \mathcal{W}_0(x, t) + \mathcal{W}_1(x, t) + \mathcal{W}_2(x, t) + \mathcal{W}_3(x, t) + \dots, \\ &= 2 \frac{\sigma \exp(\sigma x)}{1 + \exp(2\sigma x)} - 2 \frac{\sigma^4 \exp(\sigma x) (1 - \exp(2\sigma x))}{(1 + \exp(2\sigma x_1))^2} \frac{t^{\lambda}}{\Gamma(\lambda + 1)} \\ &\quad + 2 \frac{\sigma^7 \exp(\sigma x) (1 - 6 \exp(2\sigma x) + \exp(4\sigma x))}{(1 + \exp(2\sigma x_1))^3} \frac{t^{2\lambda}}{\Gamma(2\lambda + 1)} + \dots \end{aligned} \quad (4.15)$$

Case II. Here, we surmise ABC fractional derivative operator coupled with the Jafari transform and Adomian decomposition method.

Applying the Jafari transform on (4.10).

$$\begin{aligned} & \frac{\psi^\lambda(\mathfrak{s})\mathbb{A}(\lambda)}{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})} \mathcal{U}(x, \mathfrak{s}) - \phi(\mathfrak{s}) \sum_{\kappa=0}^{m-1} \psi^{\lambda-\kappa-1}(\mathfrak{s}) \mathcal{W}^{(\kappa)}(0) \\ &= -\mathbb{J} \left[6\mathcal{W}^2 \mathcal{W}_x(x, t) + \mathcal{W}_{xxx}(x, t) \right]. \end{aligned} \quad (4.16)$$

Taking into consideration the IC given in (4.11), we have

$$\mathcal{U}(x, \mathfrak{s}) = \frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) - \frac{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})}{\psi^\lambda(\mathfrak{s})\mathbb{A}(\lambda)} \mathbb{J} \left[6\mathcal{W}^2 \mathcal{W}_x(x, t) + \mathcal{W}_{xxx}(x, t) \right].$$

Employing the inverse Jafari transform, we obtain

$$\mathcal{W}(x, t) = \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) - \frac{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})}{\psi^\lambda(\mathfrak{s})\mathbb{A}(\lambda)} \mathbb{J} \left[6\mathcal{W}^2 \mathcal{W}_x(x, t) + \mathcal{W}_{xxx}(x, t) \right] \right]. \quad (4.17)$$

Thanks to the JDM, we find

$$\begin{aligned} \mathcal{W}_0(x, t) &= \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) \right] = 2\mathbb{J}^{-1} \left[\frac{\psi(s_1)}{\phi(\mathfrak{s})} \frac{\sigma \exp(\sigma x)}{1 + \exp(2\sigma x)} \right] \\ &= \frac{\sigma \exp(\sigma x)}{1 + \exp(2\sigma x)}. \end{aligned}$$

Here, we surmise that the unknown function $\mathcal{W}(x, t)$ can be written by an infinite series of the form

$$\mathcal{W}(x, t) = \sum_{\ell=0}^{\infty} \mathcal{W}_\ell(x, t).$$

Also, the non-linearity $\mathcal{F}(\mathcal{W})$, can be decomposed by an infinite series of polynomials represented by

$$\mathcal{F}_1(\mathcal{W}) = \mathcal{W}^2 \mathcal{W}_x = \sum_{\ell=0}^{\infty} \mathcal{B}_\ell,$$

where $\mathcal{W}_\ell(x, t)$ will be evaluated recurrently, and \mathcal{B}_ℓ is the so-called polynomial of $\mathcal{W}_0, \mathcal{W}_1, \dots, \mathcal{W}_\ell$ defined in (4.14).

For $\ell = 0, 1, 2, 3, \dots$

$$\begin{aligned} \mathcal{W}_1(x, t) &= -\mathbb{J}^{-1} \left[\frac{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})}{\psi^\lambda(\mathfrak{s})\mathbb{A}(\lambda)} \mathbb{J} \left[6\mathcal{B}_0 + \mathcal{W}_{0xxx} \right] \right] \\ &= -\frac{2}{\mathbb{A}(\lambda)} \frac{\sigma^4 \exp(\sigma x)(1 - \exp(2\sigma x))}{(1 + \exp(2\sigma x))^2} \left[\frac{\lambda t^\lambda}{\Gamma(\lambda + 1)} + (1 - \lambda) \right], \\ \mathcal{W}_2(x, t) &= \mathbb{J}^{-1} \left[\frac{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})}{\psi^\lambda(\mathfrak{s})\mathbb{A}(\lambda)} \mathbb{J} \left[6\mathcal{A}_1 + \mathcal{W}_{1xxx} \right] \right] \end{aligned}$$

$$\begin{aligned}
&= \frac{2}{\mathbb{A}^2(\lambda)} \frac{\sigma^7 \exp(\sigma x)(1 - 6 \exp(2\sigma x) + \exp(4\sigma x))}{(1 + \exp(2\sigma x_1))^3} \\
&\quad \times \left[\frac{\lambda^2 t^{2\lambda}}{\Gamma(2\lambda + 1)} + 2\lambda(1 - \lambda) \frac{t^\lambda}{\Gamma(\lambda + 1)} + (1 - \lambda)^2 \right], \\
&\quad \vdots
\end{aligned}$$

The approximate solution for Example 4.2 is expressed as:

$$\begin{aligned}
\mathcal{W}(x, t) &= \mathcal{W}_0(x, t) + \mathcal{W}_1(x, t) + \mathcal{W}_2(x, t) + \mathcal{W}_3(x, t) + \dots, \\
&= \frac{\sigma \exp(\sigma x)}{1 + \exp(2\sigma x)} - 2 \frac{\sigma^4 \exp(\sigma x)(1 - \exp(2\sigma x))}{\mathbb{A}(\lambda)(1 + \exp(2\sigma x_1))^2} \left[\frac{\lambda t^\lambda}{\Gamma(\lambda + 1)} + (1 - \lambda) \right] \\
&\quad + 2 \frac{\sigma^7 \exp(\sigma x)(1 - 6 \exp(2\sigma x) + \exp(4\sigma x))}{\mathbb{A}^2(\lambda)(1 + \exp(2\sigma x_1))^3} \\
&\quad \times \left[\frac{\lambda^2 t^{2\lambda}}{\Gamma(2\lambda + 1)} + 2\lambda(1 - \lambda) \frac{t^\lambda}{\Gamma(\lambda + 1)} + (1 - \lambda)^2 \right] + \dots . \tag{4.18}
\end{aligned}$$

For $\lambda = 1$, we obtained the exact solution of Example 4.2 as

$$\mathcal{W}(x, t) = \pm \sigma \sec h \sigma(x - \sigma^2 t).$$

Figure 3 shows the evolutionary outcomes for the explicit and approximate solutions of Example 4.2 for the particular instance $\lambda = 1$. The result generated by the proposed technique is remarkably similar to the exact solution, as shown in Figure 3.

Then, by considering only the first few elements of the nonlinear equations features are integrated, we can deduce that we have accomplished a reasonable estimation with the numerical solutions of the problem. It is obvious that by adding additional components to the decomposition series (4.15) and (4.18), the cumulative error can be diminished.

Analogously, we demonstrate the two-dimensional view of the change in fractional values of the order. Figure 4 depicts the response for exact CFD and AB fractional derivative operators. It is remarkable that pairwise collisions of particle-like phenomena (including solitary waves and breathers) are fundamental mechanisms in the production of condensed soliton gas dynamics. Deep water waves, shallow groundwater waves, internally waves in a segmented sea, and fibre optics are all manifestations of these waves. \square




Figure 3. Three-dimensional illustration of the exact and approximate solution of Example 4.2 when $\lambda = 1$, and $\sigma = 0.5$.




Figure 4. Two-dimensional illustration for change in fractional values of the order of Example 4.2 when $\sigma = 0.05$ and $t = 0.5$.

4.3. Fractional $K(2,2)$ equation

Example 4.3. Assume that the time-fractional $K(2,2)$ equation

$$\mathbf{D}_t^\lambda \mathcal{W}(x, t) + (\mathcal{W}^2)_x(x, t) + (\mathcal{W}^2)_{xxx}(x, t) = 0, \quad (4.19)$$

with IC:

$$\mathcal{W}_0(x, 0) = \frac{4}{3} \sigma \cos^2\left(\frac{x}{4}\right). \quad (4.20)$$

Proof. Foremost, we provide the solution of (4.19) in two general cases.

Case I. Firstly, we apply the Caputo fractional derivative operator coupled with the Jafari transform and Adomian decomposition method.

Applying the Jafari transform on (4.19).

$$\psi^\lambda(\mathfrak{s})\mathcal{U}(x, \mathfrak{s}) - \phi(\mathfrak{s}) \sum_{\kappa=0}^{m-1} \psi^{\lambda-\kappa-1}(\mathfrak{s})\mathcal{W}^{(\kappa)}(0) = -\mathbb{J} \left[(\mathcal{W}^2)_x(x, t) + (\mathcal{W}^2)_{xxx}(x, t) \right]. \quad (4.21)$$

Taking into consideration the IC given in (4.20), we have

$$\mathcal{U}(x, \mathfrak{s}) = \frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) - \frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[(\mathcal{W}^2)_x(x, t) + (\mathcal{W}^2)_{xxx}(x, t) \right].$$

Employing the inverse Jafari transform, we obtain

$$\mathcal{W}(x, t) = \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) - \frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[(\mathcal{W}^2)_x(x, t) + (\mathcal{W}^2)_{xxx}(x, t) \right] \right]. \quad (4.22)$$

Thanks to the JDM, we find

$$\begin{aligned} \mathcal{W}_0(x, t) &= \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \frac{4}{3} \sigma \cos^2 \left(\frac{x}{4} \right) \right] = 2\mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \frac{4}{3} \sigma \cos^2 \left(\frac{x}{4} \right) \right] \\ &= \frac{4}{3} \sigma \cos^2 \left(\frac{x}{4} \right). \end{aligned}$$

Here, we surmise that the unknown function $\mathcal{W}(x, t)$ can be written by an infinite series of the form

$$\mathcal{W}(x, t) = \sum_{\ell=0}^{\infty} \mathcal{W}_\ell(x, t).$$

Also, the non-linearities $\mathcal{F}_1(\mathcal{W})$ and $\mathcal{F}_2(\mathcal{W})$ can be decomposed by an infinite series of polynomials represented by

$$\mathcal{F}_1(\mathcal{W}) = (\mathcal{W}^2)_x = \sum_{\ell=0}^{\infty} \mathcal{D}_\ell, \quad \mathcal{F}_2(\mathcal{W}) = (\mathcal{W}^2)_{xxx} = \sum_{\ell=0}^{\infty} \mathcal{E}_\ell,$$

where $\mathcal{W}_\ell(x, t)$ will be evaluated recurrently, and \mathcal{D}_ℓ and \mathcal{E}_ℓ are the so-called polynomial of $\mathcal{W}_0, \mathcal{W}_1, \dots, \mathcal{W}_\ell$ established by [65].

$$\sum_{\ell=0}^{\infty} \mathcal{W}_{\ell+1}(x, t) = -\mathbb{J}^{-1} \left[\frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[\sum_{\ell=0}^{\infty} (\mathcal{D})_\ell + \sum_{\ell=0}^{\infty} (\mathcal{E})_\ell \right] \right], \quad \ell = 0, 1, 2, \dots$$

The first few Adomian polynomials are presented as follows:

$$\mathcal{D}_\ell((\mathcal{W}^2)_x) = \begin{cases} \mathcal{W}_{0x}^2, & \ell = 0, \\ (2\mathcal{W}_0\mathcal{W}_1)_x, & \ell = 1, \\ (2\mathcal{W}_2\mathcal{W}_0 + \mathcal{W}_1^2)_x, & \ell = 2, \end{cases}$$

$$\mathcal{E}_\ell((\mathcal{W}^2)_{xxx}) = \begin{cases} \mathcal{W}_{0xxx}^2, & \ell = 0, \\ (2\mathcal{W}_0\mathcal{W}_1)_{xxx}, & \ell = 1, \\ (2\mathcal{W}_2\mathcal{W}_0 + \mathcal{W}_1^2)_{xxx}, & \ell = 2, \end{cases} \quad (4.23)$$

For $\ell = 0, 1, 2, 3, \dots$

$$\begin{aligned} \mathcal{W}_1(x, t) &= -\mathbb{J}^{-1} \left[\frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[\mathcal{D}_0 + \mathcal{E}_0 \right] \right] = \frac{\sigma^2}{3} \sin\left(\frac{x}{2}\right) \frac{t^\lambda}{\Gamma(\lambda + 1)}, \\ \mathcal{W}_2(x, t) &= -\mathbb{J}^{-1} \left[\frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[\mathcal{D}_1 + \mathcal{E}_1 \right] \right] = -\frac{\sigma^3}{6} \sin\left(\frac{x}{2}\right) \frac{t^{2\lambda}}{\Gamma(2\lambda + 1)}, \\ \mathcal{W}_3(x, t) &= -\mathbb{J}^{-1} \left[\frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[\mathcal{D}_2 + \mathcal{E}_2 \right] \right] = -\frac{\sigma^4}{12} \sin\left(\frac{x}{2}\right) \frac{t^{3\lambda}}{\Gamma(3\lambda + 1)} \\ &\vdots \end{aligned}$$

The approximate solution for Example 4.3 is expressed as:

$$\begin{aligned} \mathcal{W}(x, t) &= \mathcal{W}_0(x, t) + \mathcal{W}_1(x, t) + \mathcal{W}_2(x, t) + \mathcal{W}_3(x, t) + \dots, \\ &= \frac{4}{3}\sigma \cos^2\left(\frac{x}{4}\right) + \frac{\sigma^2}{3} \sin\left(\frac{x}{2}\right) \frac{t^\lambda}{\Gamma(\lambda + 1)} \\ &\quad - \frac{\sigma^3}{6} \sin\left(\frac{x}{2}\right) \frac{t^{2\lambda}}{\Gamma(2\lambda + 1)} - \frac{\sigma^4}{12} \sin\left(\frac{x}{2}\right) \frac{t^{3\lambda}}{\Gamma(3\lambda + 1)} + \dots \end{aligned} \quad (4.24)$$

Case II. Here, we surmise ABC fractional derivative operator coupled with the Jafari transform and Adomian decomposition method. Applying the Jafari transform for Example 4.19.

$$\begin{aligned} &\frac{\psi^\lambda(\mathfrak{s})\mathbb{A}(\lambda)}{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})} \mathcal{U}(x, \mathfrak{s}) - \phi(\mathfrak{s}) \sum_{\kappa=0}^{m-1} \psi^{\lambda-\kappa-1}(\mathfrak{s}) \mathcal{W}^{(\kappa)}(0) \\ &= -\mathbb{J} \left[(\mathcal{W}^2)_x(x, t) + (\mathcal{W}^2)_{xxx}(x, t) \right]. \end{aligned} \quad (4.25)$$

Taking into consideration the IC given in (4.20), we have

$$\mathcal{U}(x, \mathfrak{s}) = \frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) - \frac{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})}{\psi^\lambda(\mathfrak{s})\mathbb{A}(\lambda)} \mathbb{J} \left[(\mathcal{W}^2)_x(x, t) + (\mathcal{W}^2)_{xxx}(x, t) \right].$$

Employing the inverse Jafari transform, we obtain

$$\mathcal{W}(x, t) = \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) - \frac{\lambda + (1 - \lambda)\psi^\lambda(\mathfrak{s})}{\psi^\lambda(\mathfrak{s})\mathbb{A}(\lambda)} \mathbb{J} \left[(\mathcal{W}^2)_x(x, t) + (\mathcal{W}^2)_{xxx}(x, t) \right] \right]. \quad (4.26)$$

Thanks to the JDM, we find

$$\mathcal{W}_0(x, t) = \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) \right] = 2\mathbb{J}^{-1} \left[\frac{\psi(\mathfrak{s}_1)}{\phi(\mathfrak{s})} \frac{4}{3} \sigma \cos^2\left(\frac{x}{4}\right) \right]$$

$$= \frac{4}{3}\sigma \cos^2\left(\frac{x}{4}\right).$$

Here, we surmise that the unknown function $\mathcal{W}(x, t)$ can be written by an infinite series of the form

$$\mathcal{W}(x, t) = \sum_{\ell=0}^{\infty} \mathcal{W}_{\ell}(x, t).$$

Also, the non-linearity $\mathcal{F}_j(\mathcal{W})$, $j = 1, 2$ can be decomposed by an infinite series of polynomials represented by

$$\mathcal{F}_1(\mathcal{W}) = (\mathcal{W}^2)_x = \sum_{\ell=0}^{\infty} \mathcal{D}_{\ell}, \quad \mathcal{F}_2(\mathcal{W}) = (\mathcal{W}^2)_{xxx} = \sum_{\ell=0}^{\infty} \mathcal{E}_{\ell},$$

where $\mathcal{W}_{\ell}(x, t)$ will be evaluated recurrently, and \mathcal{D}_{ℓ} and \mathcal{E}_{ℓ} are the so-called polynomial of $\mathcal{W}_0, \mathcal{W}_1, \dots, \mathcal{W}_{\ell}$ established defined in (4.23). Then, we have

For $\ell = 0, 1, 2, 3, \dots$

$$\begin{aligned} \mathcal{W}_1(x, t) &= -\mathbb{J}^{-1} \left[\frac{\lambda + (1 - \lambda)\psi^{\lambda}(\mathfrak{s})}{\psi^{\lambda}(\mathfrak{s})\mathbb{A}(\lambda)} \mathbb{J} \left[\mathcal{D}_0 + \mathcal{E}_0 \right] \right] \\ &= \frac{\sigma^2}{3\mathbb{A}(\lambda)} \sin\left(\frac{x}{2}\right) \left[\frac{\lambda t^{\lambda}}{\Gamma(\lambda + 1)} + (1 - \lambda) \right], \\ \mathcal{W}_2(x, t) &= \mathbb{J}^{-1} \left[\frac{\lambda + (1 - \lambda)\psi^{\lambda}(\mathfrak{s})}{\psi^{\lambda}(\mathfrak{s})\mathbb{A}(\lambda)} \mathbb{J} \left[\mathcal{D}_1 + \mathcal{E}_1 \right] \right] \\ &= -\frac{\sigma^3}{6\mathbb{A}^2(\lambda)} \sin\left(\frac{x}{2}\right) \left[\frac{\lambda^2 t^{2\lambda}}{\Gamma(2\lambda + 1)} + 2\lambda(1 - \lambda) \frac{t^{\lambda}}{\Gamma(\lambda + 1)} + (1 - \lambda)^2 \right], \\ &\vdots \end{aligned}$$

The approximate solution for Example 4.3 is expressed as:

$$\begin{aligned} \mathcal{W}(x, t) &= \mathcal{W}_0(x, t) + \mathcal{W}_1(x, t) + \mathcal{W}_2(x, t) + \mathcal{W}_3(x, t) + \dots, \\ &= \frac{4}{3}\sigma \cos^2\left(\frac{x}{4}\right) + \frac{\sigma^2}{3\mathbb{A}(\lambda)} \sin\left(\frac{x}{2}\right) \left[\frac{\lambda t^{\lambda}}{\Gamma(\lambda + 1)} + (1 - \lambda) \right] \\ &\quad - \frac{\sigma^3}{6\mathbb{A}^2(\lambda)} \sin\left(\frac{x}{2}\right) \left[\frac{\lambda^2 t^{2\lambda}}{\Gamma(2\lambda + 1)} + 2\lambda(1 - \lambda) \frac{t^{\lambda}}{\Gamma(\lambda + 1)} + (1 - \lambda)^2 \right] + \dots \quad (4.27) \end{aligned}$$

For $\lambda = 1$, we obtained the closed form solution of Example 4.3 as

$$\mathcal{W}(x, t) = \begin{cases} \frac{4}{3}\sigma \cos^2\left(\frac{x - \sigma t}{4}\right), & |x - \sigma t| \leq 2\pi, \\ 0, & \text{otherwise.} \end{cases}$$

Figure 5 shows the evolutionary outcomes for the explicit and approximate solutions of Example 4.3 for the particular instance $\lambda = 1$. The result generated by the proposed technique is remarkably similar to the exact solution, as shown in Figure 5.

Then, by considering only the first few elements of the nonlinear equations features are integrated, we can deduce that we have accomplished a reasonable estimation with the numerical solutions of the problem. It is obvious that by adding additional components to the decomposition series (4.24) and (4.27), the cumulative error can be diminished.

Analogously, we demonstrate the two-dimensional view of the change in fractional values of the order. Figure 6 depicts the response for exact CFD and AB fractional derivative operators. For a variation of the K(2,2) equation, constrained traveling-wave solutions are achieved. We acquire hump-shaped and valley-shaped solitary-wave solutions, as well as some periodic solutions, for the focusing branch. It is worth noting that optimal focusing provides the aggregate of the frequency and amplitude of the originating waves in the engaging phase, as illustrated in reference [66]. \square




Figure 5. Three-dimensional illustration of the exact and approximate solution of Example 4.3 when $\lambda = 1$ and $\sigma = 0.05$.




Figure 6. Two-dimensional illustration of the exact and approximate solution for the change in fractional values of the order of Example 4.3 when $\lambda = 1$, $\sigma = 0.05$ and $t = 0.5$.

4.4. Fractional K(3,3) equation

Example 4.4. Assume that the time-fractional K(3,3) equation

$$\mathbf{D}_t^\lambda \mathcal{W}(x, t) + (\mathcal{W}^3)_x(x, t) + (\mathcal{W}^3)_{xxx}(x, t) = 0, \quad (4.28)$$

with IC:

$$\mathcal{W}_0(x, 0) = \sqrt{\frac{3\sigma}{2}} \cos\left(\frac{x}{3}\right). \quad (4.29)$$

Proof. Foremost, we provide the solution of (4.28) in two general cases.

Case I. Firstly, we apply the Caputo fractional derivative operator coupled with the Jafari transform and Adomian decomposition method.

Applying the Jafari transform on (4.28).

$$\psi^\lambda(\mathfrak{s}) \mathcal{U}(x, \mathfrak{s}) - \phi(\mathfrak{s}) \sum_{\kappa=0}^{m-1} \psi^{\lambda-\kappa-1}(\mathfrak{s}) \mathcal{W}^{(\kappa)}(0) = -\mathbb{J} \left[(\mathcal{W}^3)_x(x, t) + (\mathcal{W}^3)_{xxx}(x, t) \right]. \quad (4.30)$$

Taking into consideration the IC given in (4.29), we have

$$\mathcal{U}(x, \mathfrak{s}) = \frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) - \frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[(\mathcal{W}^3)_x(x, t) + (\mathcal{W}^3)_{xxx}(x, t) \right].$$

Employing the inverse Jafari transform, we obtain

$$\mathcal{W}(x, t) = \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \mathcal{W}(x, 0) - \frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[(\mathcal{W}^3)_x(x, t) + (\mathcal{W}^3)_{xxx}(x, t) \right] \right]. \quad (4.31)$$

Thanks to the JDM, we find

$$\begin{aligned} \mathcal{W}_0(x, t) &= \mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \sqrt{\frac{3\sigma}{2}} \cos\left(\frac{x}{3}\right) \right] = 2\mathbb{J}^{-1} \left[\frac{\phi(\mathfrak{s})}{\psi(\mathfrak{s})} \sqrt{\frac{3\sigma}{2}} \cos\left(\frac{x}{3}\right) \right] \\ &= \sqrt{\frac{3\sigma}{2}} \cos\left(\frac{x}{3}\right). \end{aligned}$$

Here, we surmise that the unknown function $\mathcal{W}(x, t)$ can be written by an infinite series of the form

$$\mathcal{W}(x, t) = \sum_{\ell=0}^{\infty} \mathcal{W}_\ell(x, t).$$

Also, the non-linearities $\mathcal{F}_1(\mathcal{W})$ and $\mathcal{F}_2(\mathcal{W})$ can be decomposed by an infinite series of polynomials represented by

$$\mathcal{F}_1(\mathcal{W}) = (\mathcal{W}^3)_x = \sum_{\ell=0}^{\infty} \mathcal{G}_\ell, \quad \mathcal{F}_2(\mathcal{W}) = (\mathcal{W}^3)_{xxx} = \sum_{\ell=0}^{\infty} \mathcal{H}_\ell,$$

where $\mathcal{W}_\ell(x, t)$ will be evaluated recurrently, and \mathcal{G}_ℓ and \mathcal{H}_ℓ are the so-called polynomial of $\mathcal{W}_0, \mathcal{W}_1, \dots, \mathcal{W}_\ell$ established by [65].

$$\sum_{\ell=0}^{\infty} \mathcal{W}_{\ell+1}(x, t) = -\mathbb{J}^{-1} \left[\frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[\sum_{\ell=0}^{\infty} (\mathcal{G})_\ell + \sum_{\ell=0}^{\infty} (\mathcal{H})_\ell \right] \right], \quad \ell = 0, 1, 2, \dots$$

The first few Adomian polynomials are presented as follows:

$$\begin{aligned} \mathcal{G}_\ell((\mathcal{W}^3)_x) &= \begin{cases} \mathcal{W}_{0x}^3, & \ell = 0, \\ (3\mathcal{W}_0^2\mathcal{W}_1)_x, & \ell = 1, \\ (3\mathcal{W}_2\mathcal{W}_0^2 + 3\mathcal{W}_1^2\mathcal{W}_0)_x, & \ell = 2, \end{cases} \\ \mathcal{E}_\ell((\mathcal{W}^3)_{xxx}) &= \begin{cases} \mathcal{W}_{0xxx}^2, & \ell = 0, \\ (3\mathcal{W}_0^2\mathcal{W}_1)_{xxx}, & \ell = 1, \\ (3\mathcal{W}_2\mathcal{W}_0^2 + 3\mathcal{W}_1^2\mathcal{W}_0)_{xxx}, & \ell = 2, \end{cases} \end{aligned} \quad (4.32)$$

For $\ell = 0, 1, 2, 3, \dots$

$$\begin{aligned} \mathcal{W}_1(x, t) &= -\mathbb{J}^{-1} \left[\frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[\mathcal{G}_0 + \mathcal{H}_0 \right] \right] = \frac{\sqrt{6\sigma^3}}{6} \sin\left(\frac{x}{3}\right) \frac{t^\lambda}{\Gamma(\lambda+1)}, \\ \mathcal{W}_2(x, t) &= -\mathbb{J}^{-1} \left[\frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[\mathcal{G}_1 + \mathcal{H}_1 \right] \right] = -\frac{\sqrt{6\sigma^5}}{18} \sin\left(\frac{x}{3}\right) \frac{t^{2\lambda}}{\Gamma(2\lambda+1)}, \\ \mathcal{W}_3(x, t) &= -\mathbb{J}^{-1} \left[\frac{1}{\psi^\lambda(\mathfrak{s})} \mathbb{J} \left[\mathcal{G}_2 + \mathcal{H}_2 \right] \right] = -\frac{\sqrt{6\sigma^7}}{54} \sin\left(\frac{x}{3}\right) \frac{t^{3\lambda}}{\Gamma(3\lambda+1)} \\ &\vdots \end{aligned}$$

The approximate solution for Example 4.4 is expressed as:

$$\begin{aligned} \mathcal{W}(x, t) &= \mathcal{W}_0(x, t) + \mathcal{W}_1(x, t) + \mathcal{W}_2(x, t) + \mathcal{W}_3(x, t) + \dots, \\ &= \sqrt{\frac{3\sigma}{2}} \cos\left(\frac{x}{3}\right) + \frac{\sqrt{6\sigma^3}}{6} \sin\left(\frac{x}{3}\right) \frac{t^\lambda}{\Gamma(\lambda+1)} \\ &\quad - \frac{\sqrt{6\sigma^5}}{18} \sin\left(\frac{x}{3}\right) \frac{t^{2\lambda}}{\Gamma(2\lambda+1)} - \frac{\sqrt{6\sigma^7}}{54} \sin\left(\frac{x}{3}\right) \frac{t^{3\lambda}}{\Gamma(3\lambda+1)} + \dots \end{aligned} \quad (4.33)$$

Case II. Here, we surmise ABC fractional derivative operator coupled with the Jafari transform and Adomian decomposition method. Applying the Jafari transform for Example 4.28.

$$\begin{aligned} &\frac{\psi^\lambda(\mathfrak{s})\mathbb{A}(\lambda)}{\lambda + (1-\lambda)\psi^\lambda(\mathfrak{s})} \mathcal{U}(x, \mathfrak{s}) - \phi(\mathfrak{s}) \sum_{\kappa=0}^{m-1} \psi^{\lambda-\kappa-1}(\mathfrak{s}) \mathcal{W}^{(\kappa)}(0) \\ &= -\mathbb{J} \left[(\mathcal{W}^3)_x(x, t) + (\mathcal{W}^3)_{xxx}(x, t) \right]. \end{aligned} \quad (4.34)$$

Taking into consideration the IC given in (4.29), we have

$$\mathcal{U}(x, s) = \frac{\phi(s)}{\psi(s)} \mathcal{W}(x, 0) - \frac{\lambda + (1 - \lambda)\psi^\lambda(s)}{\psi^\lambda(s)\mathbb{A}(\lambda)} \mathbb{J} \left[(\mathcal{W}^3)_x(x, t) + (\mathcal{W}^3)_{xxx}(x, t) \right].$$

Employing the inverse Jafari transform, we obtain

$$\mathcal{W}(x, t) = \mathbb{J}^{-1} \left[\frac{\phi(s)}{\psi(s)} \mathcal{W}(x, 0) - \frac{\lambda + (1 - \lambda)\psi^\lambda(s)}{\psi^\lambda(s)\mathbb{A}(\lambda)} \mathbb{J} \left[(\mathcal{W}^3)_x(x, t) + (\mathcal{W}^3)_{xxx}(x, t) \right] \right]. \quad (4.35)$$

Thanks to the JDM, we find

$$\begin{aligned} \mathcal{W}_0(x, t) &= \mathbb{J}^{-1} \left[\frac{\phi(s)}{\psi(s)} \mathcal{W}(x, 0) \right] = 2\mathbb{J}^{-1} \left[\frac{\psi(s_1)}{\phi(s)} \sqrt{\frac{3\sigma}{2}} \cos\left(\frac{x}{3}\right) \right] \\ &= \sqrt{\frac{3\sigma}{2}} \cos\left(\frac{x}{3}\right). \end{aligned}$$

Here, we surmise that the unknown function $\mathcal{W}(x, t)$ can be written by an infinite series of the form

$$\mathcal{W}(x, t) = \sum_{\ell=0}^{\infty} \mathcal{W}_\ell(x, t).$$

Also, the non-linearity $\mathcal{F}_j(\mathcal{W})$, $j = 1, 2$ can be decomposed by an infinite series of polynomials represented by

$$\mathcal{F}_1(\mathcal{W}) = (\mathcal{W}^3)_x = \sum_{\ell=0}^{\infty} \mathcal{G}_\ell, \quad \mathcal{F}_2(\mathcal{W}) = (\mathcal{W}^3)_{xxx} = \sum_{\ell=0}^{\infty} \mathcal{H}_\ell,$$

where $\mathcal{W}_\ell(x, t)$ will be evaluated recurrently, and \mathcal{D}_ℓ and \mathcal{E}_ℓ are the so-called polynomial of $\mathcal{W}_0, \mathcal{W}_1, \dots, \mathcal{W}_\ell$ established defined in (4.32). Then, we have

For $\ell = 0, 1, 2, 3, \dots$

$$\begin{aligned} \mathcal{W}_1(x, t) &= -\mathbb{J}^{-1} \left[\frac{\lambda + (1 - \lambda)\psi^\lambda(s)}{\psi^\lambda(s)\mathbb{A}(\lambda)} \mathbb{J} \left[\mathcal{G}_0 + \mathcal{H}_0 \right] \right] \\ &= \frac{\sqrt{6\sigma^3}}{6\mathbb{A}(\lambda)} \sin\left(\frac{x}{3}\right) \left[\frac{\lambda t^\lambda}{\Gamma(\lambda + 1)} + (1 - \lambda) \right], \\ \mathcal{W}_2(x, t) &= \mathbb{J}^{-1} \left[\frac{\lambda + (1 - \lambda)\psi^\lambda(s)}{\psi^\lambda(s)\mathbb{A}(\lambda)} \mathbb{J} \left[\mathcal{G}_1 + \mathcal{H}_1 \right] \right] \\ &= -\frac{\sqrt{6\sigma^5}}{18\mathbb{A}^2(\lambda)} \sin\left(\frac{x}{3}\right) \left[\frac{\lambda^2 t^{2\lambda}}{\Gamma(2\lambda + 1)} + 2\lambda(1 - \lambda) \frac{t^\lambda}{\Gamma(\lambda + 1)} + (1 - \lambda)^2 \right], \\ &\vdots \end{aligned}$$

The approximate solution for Example 4.4 is expressed as:

$$\mathcal{W}(x, t) = \mathcal{W}_0(x, t) + \mathcal{W}_1(x, t) + \mathcal{W}_2(x, t) + \mathcal{W}_3(x, t) + \dots,$$

$$\begin{aligned}
&= \sqrt{\frac{3\sigma}{2}} \cos\left(\frac{x}{3}\right) + \frac{\sqrt{6\sigma^3}}{6A(\lambda)} \sin\left(\frac{x}{3}\right) \left[\frac{\lambda t^\lambda}{\Gamma(\lambda+1)} + (1-\lambda) \right] \\
&\quad - \frac{\sqrt{6\sigma^5}}{18A^2(\lambda)} \sin\left(\frac{x}{3}\right) \left[\frac{\lambda^2 t^{2\lambda}}{\Gamma(2\lambda+1)} + 2\lambda(1-\lambda) \frac{t^\lambda}{\Gamma(\lambda+1)} + (1-\lambda)^2 \right] + \dots \quad (4.36)
\end{aligned}$$

For $\lambda = 1$, we obtained the closed form solution of Example 4.4 as

$$\mathcal{W}(x, t) = \begin{cases} \frac{\sqrt{6\sigma}}{2} \sigma \cos\left(\frac{x-\sigma t}{3}\right), & |x - \sigma t| \leq \frac{3\pi}{2}, \\ 0, & \text{otherwise.} \end{cases}$$

Figure 7 shows the evolutionary outcomes for the explicit and approximate solutions of Example 4.4 for the particular instance $\lambda = 1$. The result generated by the proposed technique is remarkably similar to the exact solution, as shown in Figure 7.

Then, by considering only the first few elements of the nonlinear equations features are integrated, we can deduce that we have accomplished a reasonable estimation with the numerical solutions of the problem. It is obvious that by adding additional components to the decomposition series (4.33) and (4.36), the cumulative error can be diminished.

Analogously, we demonstrate the two-dimensional view of the change in fractional values of the order. Figure 8 depicts the response for exact CFD and AB fractional derivative operators.

For a variation of the K(3,3) equation, constrained traveling-wave solutions are achieved. We acquire hump-shaped and valley-shaped solitary-wave solutions, as well as some periodic solutions, for the focusing branch. It is worth noting that optimal focusing provides the aggregate of the frequency and amplitude of the originating waves in the engaging phase, as illustrated in reference [66]. \square




Figure 7. Three-dimensional illustration of the exact and approximate solution of Example 4.4 when $\lambda = 1$ and $\sigma = 0.05$.




Figure 8. Two-dimensional illustration of the exact and approximate solution of change in fractional values of the order of Example 4.4 when $\lambda = 1$, $\sigma = 0.5$ and $t = 0.5$.

5. Conclusions

In this paper, we conducted a novel algorithm based on the Jafari transform and Adomian decomposition method, known as the Jafari decomposition method. In the time-fractional technique, we investigated several models such as KdV, mKdV, K (2,2), and K (3,3). To comprehend their physical interpretation, we researched and examined several novel families of solutions and their simulation studies, presented in two-dimensional and three-dimensional plots. The new discoveries concerned the hyperbolic function, trigonometric function, exponential function, and constant function. These new solutions and results might be appreciated in the laser, plasma sciences and wave pattern. To summarise, the suggested method stated above was determined to solve this collection of challenges by utilizing successive fast converging approximations without any limiting requirements or manipulations that changed the physical attributes of the concerns. Also, increasing the recursive procedure leads to the closed form solution of the governing equation.

Conflict of interest

The authors declare that they have no competing interests.

References

1. K. Karthikeyan, P. Karthikeyan, H. M. Baskonus, K. Venkatachalam, Y. M. Chu, Almost sectorial operators on Ψ -Hilfer derivative fractional impulsive integro-differential equations, *Math. Method. Appl. Sci.*, 2021. <https://doi.org/10.1002/mma.7954>
2. S. Rashid, S. Sultana, Y. Karaca, A. Khalid, Y. M. Chu, Some further extensions considering discrete proportional fractional operators, *Fractals*, **30** (2022), 2240026. <https://doi.org/10.1142/S0218348X22400266>

3. Y. M. Chu, U. Nazir, M. Sohail, M. M. Selim, J. R. Lee, Enhancement in thermal energy and solute particles using hybrid nanoparticles by engaging activation energy and chemical reaction over a parabolic surface via finite element approach, *Fractal Fract.*, **5** (2021), 119. <https://doi.org/10.3390/fractalfract5030119>
4. T. H. Zhao, M. I. Khan, Y. M. Chu, Artificial neural networking (ANN) analysis for heat and entropy generation in flow of non-Newtonian fluid between two rotating disks, *Math. Method. Appl. Sci.*, 2021. <https://doi.org/10.1002/mma.7310>
5. Y. M. Chu, B. M. Shankaralingappa, B. J. Gireesha, F. Alzahrani, M. I. Khan, S. U. Khan, Combined impact of Cattaneo-Christov double diffusion and radiative heat flux on bio-convective flow of Maxwell liquid configured by a stretched nano-material surface, *Appl. Math. Comput.*, **419** (2022), 126883. <https://doi.org/10.1016/j.amc.2021.126883>
6. M. Nazeer, F. Hussain, M. I. Khan, A. ur Rehman, E. R. El-Zahar, Y. M. Chu, et al., Theoretical study of MHD electro-osmotically flow of third-grade fluid in micro channel, *Appl. Math. Comput.*, **420** (2022), 126868. <https://doi.org/10.1016/j.amc.2021.126868>
7. T. H. Zhao, O. Castillo, H. Jahanshahi, A. Yusuf, M. O. Alassafi, F. E. Alsaadi, et al., A fuzzy-based strategy to suppress the novel coronavirus (2019-NCOV) massive outbreak, *Appl. Comput. Math.*, **20** (2021), 160–176.
8. S. Rashid, F. Jarad, T. M. Jawa, A study of behaviour for fractional order diabetes model via the nonsingular kernel, *AIMS Mathematics*, **7** (2022), 5072–5092. <https://doi.org/10.3934/math.2022282>
9. S. Rashid, F. Jarad, F. S. Bayones, On new computations of the fractional epidemic childhood disease model pertaining to the generalized fractional derivative with nonsingular kernel, *AIMS Mathematics*, **7** (2022), 4552–4573. <https://doi.org/10.3934/math.2022254>
10. T. H. Zhao, M. K. Wang, Y. M. Chu, On the bounds of the perimeter of an ellipse, *Acta Math. Sci.*, **42** (2022), 491–501. <https://doi.org/10.1007/s10473-022-0204-y>
11. T. H. Zhao, M. K. Wang, G. J. Hai, Y. M. Chu, Landen inequalities for Gaussian hypergeometric function, *RACSAM*, **116** (2022), 53. <https://doi.org/10.1007/s13398-021-01197-y>
12. T. H. Zhao, Z. Y. He, Y. M. Chu, Sharp bounds for the weighted Hölder mean of the zero-balanced generalized complete elliptic integrals, *Comput. Methods Funct. Theory*, **21** (2021), 413–426. <https://doi.org/10.1007/s40315-020-00352-7>
13. T. H. Zhao, L. Shi, Y. M. Chu, Convexity and concavity of the modified Bessel functions of the first kind with respect to Hölder means, *RACSAM*, **114** (2020), 96. <https://doi.org/10.1007/s13398-020-00825-3>
14. T. H. Zhao, M. K. Wang, Y. M. Chu, Concavity and bounds involving generalized elliptic integral of the first kind, *J. Math. Inequal*, **15** (2021), 701–724. <https://doi.org/10.7153/jmi-2021-15-50>
15. H. H. Chu, T. H. Zhao, Y. M. Chu, Sharp bounds for the Toader mean of order 3 in terms of arithmetic, quadratic and contraharmonic means, *Math. Slovaca*, **70** (2020), 1097–1112. <https://doi.org/10.1515/ms-2017-0417>
16. T. H. Zhao, Z. Y. He, Y. M. Chu, On some refinements for inequalities involving zero-balanced hypergeometric function, *AIMS Mathematics*, **5** (2020), 6479–6495. <https://doi.org/10.3934/math.2020418>
17. M. Caputo, *Elasticita e dissipazione*, Bologna: Zanichelli, 1969.

18. A. Atangana, D. Baleanu, New fractional derivatives with non-local and non-singular kernel: Theory and application to heat transfer model, *Therm. Sci.*, **20** (2016), 763–769. <https://doi.org/10.2298/TSCI160111018A>
19. A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, *Theory and applications of fractional differential equations*, The Netherlands: Amsterdam, 2006.
20. W. X. Ma, N-soliton solutions and the Hirota conditions in (1+1)-dimensions, *Int. J. Nonlin. Sci. Num.*, **23** (2022), 123–133. <https://doi.org/10.1515/ijnsns-2020-0214>
21. W. X. Ma, N-soliton solution and the Hirota condition of a (2+1)-dimensional combined equation, *Math. Comput. Simulat.*, **190** (2021), 270–279. <https://doi.org/10.1016/j.matcom.2021.05.020>
22. W. X. Ma, N-soliton solution of a combined pKP-BKP equation, *J. Geom. Phys.*, **165** (2021), 104191. <https://doi.org/10.1016/j.geomphys.2021.104191>
23. W. X. Ma, X. L. Yong, X. Lü, Soliton solutions to the B-type Kadomtsev-Petviashvili equation under general dispersion relations, *Wave Motion*, **103** (2021), 102719. <https://doi.org/10.1016/j.wavemoti.2021.102719>
24. R. G. Zhang, L. G. Yang, Theoretical analysis of equatorial near-inertial solitary waves under complete Coriolis parameters, *Acta Oceanol. Sin.*, **40** (2021), 54–61. <https://doi.org/10.1007/s13131-020-1699-5>
25. J. Q. Zhang, R. G. Zhang, L. G. Yang, Q. S. Liu, L. G. Chen, Coherent structures of nonlinear barotropic-baroclinic interaction in unequal depth two-layer model, *Appl. Math. Comput.*, **408** (2021), 126347. <https://doi.org/10.1016/j.amc.2021.126347>
26. J. Wang, R. G. Zhang, L. G. Yang, A Gardner evolution equation for topographic Rossby waves and its mechanical analysis, *Appl. Math. Comput.*, **385** (2020), 125426. <https://doi.org/10.1016/j.amc.2020.125426>
27. J. Wang, R. G. Zhang, L. G. Yang, Solitary waves of nonlinear barotropic–baroclinic coherent structures, *Phys. Fluids*, **32** (2020), 096604. <https://doi.org/10.1063/5.0025167>
28. A. J. M. Jawad, New exact solutions of nonlinear partial differential equations using Tan-Cot function method, *Stud. Math. Sci.*, **5** (2012), 13–25.
29. W. J. Li, Y. N. Pang, Application of Adomian decomposition method to nonlinear systems, *Adv. Differ. Equ.*, **2020** (2020), 67. <https://doi.org/10.1186/s13662-020-2529-y>
30. J. H. He, Homotopy perturbation technique, *Comput. Method. Appl. Mech. Eng.*, **178** (1999), 257–262. [https://doi.org/10.1016/S0045-7825\(99\)00018-3](https://doi.org/10.1016/S0045-7825(99)00018-3)
31. M. Turkyilmazoglu, A note on the homotopy analysis method, *Appl. Math. Lett.*, **23** (2010), 1226–1230. <https://doi.org/10.1016/j.aml.2010.06.003>
32. A. H. A. AliaK, K. R. Raslan, Variational iteration method for solving partial differential equations with variable coefficients, *Chaos Soliton. Fract.*, **40** (2009), 1520–1529. <https://doi.org/10.1016/j.chaos.2007.09.031>
33. Q. M. Al-Mdallal, M. I. Syam, M. N. Anwar, A collocation-shooting method for solving fractional boundary value problems, *Commun. Nonlinear Sci.*, **15** (2010), 3814–3822. <https://doi.org/10.1016/j.cnsns.2010.01.020>
34. H. Naher, F. A. Abdullah, New approach of (G/G') -expansion method and new approach of generalized (G/G') -expansion method for nonlinear evolution equation, *AIP Adv.*, **3** (2013), 032116. <https://doi.org/10.1063/1.4794947>

35. J. Manafian, M. Foroutan, Application of $\tan(\phi(\tau)/2)$ -expansion method for the time-fractional Kuramoto–Sivashinsky equation, *Opt. Quant. Electron.*, **49** (2017), 272. <https://doi.org/10.1007/s11082-017-1107-3>
36. N. Maarouf, H. Maadan, K. Hilal, Lie symmetry analysis and explicit solutions for the time-fractional regularized long-wave equation, *Int. J. Differ. Equ.*, **2021** (2021), 6614231. <https://doi.org/10.1155/2021/6614231>
37. G. Hariharan, K. Kannan, Review of wavelet methods for the solution of reaction–diffusion problems in science and engineering, *Appl. Math. Model.*, **38** (2014), 799–813. <https://doi.org/10.1016/j.apm.2013.08.003>
38. S. Rashid, S. Sultana, R. Ashraf, M. k. A. Kaabar, On comparative analysis for the Black-Scholes model in the generalized fractional derivatives sense via Jafari transform, *J. Funct. Space*, **2021** (2021), 7767848. <https://doi.org/10.1155/2021/7767848>
39. P. E. Holloway, E. Pelinovsky, T. Talipova, A generalized Korteweg-de Vries model of internal tide transformation in the coastal zone, *J. Geophys Res.-Oceans*, **104** (1999), 18333–18350. <https://doi.org/10.1029/1999JC900144>
40. A. M. Wazwaz, The variational iteration method for solving linear and nonlinear ODEs and scientific models with variable coefficients, *Cent. Eur. J. Eng.*, **4** (2014), 64–71. <https://doi.org/10.2478/s13531-013-0141-6>
41. Y. A. Stepanyants, Nonlinear waves in a rotating ocean (the Ostrovsky equation and its generalizations and applications), *Izv. Atmos. Ocean. Phys.*, **56** (2020), 16–32. <https://doi.org/10.1134/S0001433820010077>
42. S. Rashid, A. Khalid, S. Sultana, Z. Hammouch, R. Shah, A. M. Alsharif, A novel analytical view of time-fractional Korteweg-De Vries equations via a new integral transform, *Symmetry*, **13** (2021), 1254. <https://doi.org/10.3390/sym13071254>
43. S. Rashid, Z. Hammouch, H. Aydi, A. G. Ahmad, A. M. Alsharif, Novel computations of the time-fractional Fisher’s model via generalized fractional integral operators by means of the Elzaki transform, *Fractal Fract.*, **5** (2021), 94. <https://doi.org/10.3390/fractalfract5030094>
44. S. Rashid, K. T. Kubra, J. L. G. Guirao, Construction of an approximate analytical solution for multi-dimensional fractional Zakharov-Kuznetsov equation via Aboodh Adomian decomposition method, *Symmetry*, **13** (2021), 1542. <https://doi.org/10.3390/sym13081542>
45. H. Jafari, A new general integral transform for solving integral equations, *J. Adv. Res.*, **32** (2021), 133–138. <https://doi.org/10.1016/j.jare.2020.08.016>
46. L. Debnath, D. Bhatta, *Integral transforms and their applications*, Boca Raton: CRC Press, 2014.
47. F. Jarad, T. Abdeljawad, A modified Laplace transform for certain generalized fractional operators, *Res. Nonlinear Anal.*, **1** (2018), 88–98.
48. G. K. Watugala, Sumudu transform: A new integral transform to solve differential equations and control engineering problems, *Int. J. Math. Edu. Sci. Technol.*, **24** (1993), 35–43. <https://doi.org/10.1080/0020739930240105>
49. K. S. Aboodh, The new integral transform Aboodh transform. *Glob. J. Pure Appl. Math.*, **9** (2013), 35–43.
50. S. A. P. Ahmadi, H. Hosseinzadeh, A. Y. Cherati, A new integral transform for solving higher order ordinary differential equations, *Nonlinear Dyn. Syst. Theory*, **19** (2019), 243–252.

51. S. A. P. Ahmadi, H. Hosseinzadeh, A. Y. Cherati, A new integral transform for solving higher order linear ordinary Laguerre and Hermite differential equations, *Int. J. Appl. Comput. Math.*, **5** (2019), 142. <https://doi.org/10.1007/s40819-019-0712-1>
52. T. M. Elzaki, The new integral transform Elzaki transform, *Global. J. Pure Appl. Math.*, **7** (2011), 57–64.
53. Z. H. Khan, W. A. Khan, N-transform properties and applications, *NUST. J. Eng. Sci.*, **1** (2008), 127–133.
54. M. M. A. Mahgoub, The new integral transform “Mohand Transform”, *Adv. Theor. Appl. Math.*, **12** (2017), 113–120.
55. M. M. A. Mahgoub, The new integral transform “Sawi Transform”, *Adv. Theor. Appl. Math.*, **14** (2019), 81–87.
56. A. Kamal, H. Sedeeg, The new integral transform “Kamal Transform”, *Adv. Theor. Appl. Math.* **11** (2016), 451–458.
57. H. Kim, On the form and properties of an integral transform with strength in integral transforms, *FJMS*, **102** (2017), 2831–2844. <http://doi.org/10.17654/MS102112831>
58. H. Kim, The intrinsic structure and properties of Laplace-typed integral transforms, *Math. Probl. Eng.*, **2017** (2017), 1762729. <https://doi.org/10.1155/2017/1762729>
59. M. Meddahi, H. Jafari, M. N. Ncube, New general integral transform via Atangana–Baleanu derivatives, *Adv. Differ. Equ.*, **2021** (2021), 385. <https://doi.org/10.1186/s13662-021-03540-4>
60. A. Atangana, D. Baleanu, New fractional derivatives with nonlocal and non-singular kernel: Theory and application to heat transfer model, 2016, arXiv: 1602.03408.
61. A. Atangana, I. Koca, Chaos in a simple nonlinear system with Atangana–Baleanu derivatives with fractional order, *Chaos Soliton Fract.*, **89** (2016), 447–454. <https://doi.org/10.1016/j.chaos.2016.02.012>
62. M. Yavuz, T. Abdeljawad, Nonlinear regularized long-wave models with a new integral transformation applied to the fractional derivative with power and Mittag-Leffler kernel, *Adv. Differ. Equ.*, **2020** (2020), 367. <https://doi.org/10.1186/s13662-020-02828-1>
63. A. Bokhari, D. Baleanu, R. Belgacema, Application of Shehu transform to Atangana–Baleanu derivatives, *J. Math. Comput. Sci.*, **20** (2020), 101–107. <https://doi.org/10.22436/jmcs.020.02.03>
64. M. G. Mittag-Leffler, *Sur la nouvelle fonction Ea(x)*, Paris: C. R. Academy of Science, 1903.
65. G. Adomian, R. Rach, Modified Adomian polynomial, *Math. Comput. Model.*, **24** (1996), 39–46. [https://doi.org/10.1016/S0895-7177\(96\)00171-9](https://doi.org/10.1016/S0895-7177(96)00171-9)
66. A. V. Slunyaev, E. N. Pelinovsky, Role of multiple soliton interactions in the generation of rogue waves: The modified Korteweg–de Vries framework, *Phys. Rev. Lett.*, **117** (2016), 214501. <https://doi.org/10.1103/PhysRevLett.117.214501>



AIMS Press

©2022 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)