

Research Article

Humaira Kalsoom, Muhammad Amer Latif, Saima Rashid, Dumitru Baleanu, and Yu-Ming Chu*

New (p, q) -estimates for different types of integral inequalities via (α, m) -convex mappings

<https://doi.org/10.1515/math-2020-0114>

received September 5, 2019; accepted November 12, 2020

Abstract: In the article, we present a new (p, q) -integral identity for the first-order (p, q) -differentiable functions and establish several new (p, q) -quantum error estimations for various integral inequalities via (α, m) -convexity. We also compare our results with the previously known results and provide two examples to show the superiority of our obtained results.

Keywords: (p, q) -quantum calculus, Hermite-Hadamard inequality, Simpson's type inequality, (α, m) -convex functions

MSC 2020: 26D15, 26D10, 26A51

1 Introduction

Integral inequalities are considered a fabulous tool for constructing the qualitative and quantitative properties in the field of pure and applied mathematics [1–20]. A continuous growth of interest has been occurring to meet the requirements for the wide applications of these inequalities. These applications are closely related to the convex functions and have been studied by many researchers using various techniques [21–32].

Now, we recall the definition of convex function as follows.

Let $K \subseteq \mathbb{R}$ be an interval. Then a real-valued function $g : K \rightarrow \mathbb{R}$ is said to be convex if the inequality

$$g(\lambda\phi + (1 - \lambda)\psi) \leq \lambda g(\phi) + (1 - \lambda)g(\psi)$$

holds for all $\phi, \psi \in K$ and $\lambda \in [0, 1]$.

For convex functions, many inequalities have been established by many authors, for example, Jensen inequality [33], Ostrowski inequality [34], hypergeometric function inequality [35], elliptic integral inequality

* **Corresponding author: Yu-Ming Chu**, Department of Mathematics, Huzhou University, Huzhou 313000, China; School of Mathematics and Statistics, Changsha University of Science & Technology, Changsha 410014, China, e-mail: chuyuming2005@126.com

Humaira Kalsoom: Department of Mathematics, Zhejiang Normal University, Jinhua 321004, P. R. China, e-mail: humaira87@zju.edu.cn, humairazju@yahoo.com

Muhammad Amer Latif: Department of Basic Sciences, Deanship of Preparatory Year, King Faisal University, Hofuf Al-Hasa 31982, Saudi Arabia, e-mail: m_amer_latif@hotmail.com

Saima Rashid: Department of Mathematics, Government College University, Faisalabad 38000, Pakistan, e-mail: saimarashid@gcuf.edu.pk

Dumitru Baleanu: Department of Mathematics, Çankaya University, Ankara 06530, Turkey, e-mail: dumitru@cankaya.edu.tr

ities [36–41] and so on. But the most celebrated and significant inequality is the Hermite-Hadamard inequality [42,43], which is stated as follows.

Let $\phi < \psi$ and $g : [\phi, \psi] \rightarrow \mathbb{R}$ be a convex function. Then the double inequality

$$g\left(\frac{\phi + \psi}{2}\right) \leq \frac{1}{\psi - \phi} \int_{\phi}^{\psi} g(\lambda) d\lambda \leq \frac{g(\phi) + g(\psi)}{2} \tag{1.1}$$

The Hermite-Hadamard inequality (1.1) has been extensively discussed because it is essential in developing a connection between the theory of convex functions and integral inequalities. A number of researchers have dedicated their efforts to extend, generalize and refine the Hermite-Hadamard inequality (1.1) for different classes of convex functions and mappings. Some recent results on inequality (1.1) can be found in the literature [44–46].

Let $g : [\phi, \psi] \rightarrow \mathbb{R}$ be a four times continuous and differentiable mapping on the interval $[\phi, \psi]$ such that $\|g^{(4)}\|_{\infty} = \sup_{z \in (\phi, \psi)} |g^{(4)}(z)| < \infty$. Then Simpson’s inequality [47]

$$\left| \frac{1}{3} \left[\frac{g(\phi) + g(\psi)}{2} + 2g\left(\frac{\phi + \psi}{2}\right) - \frac{1}{\phi - \psi} \int_{\phi}^{\psi} g(z) dz \right] \right| \leq \frac{(\psi - \phi)^4}{2,880} \|g^{(4)}\|_{\infty}$$

holds.

Quantum calculus is the study of calculus without limits and is also known as q -calculus [48]. In q -calculus, we obtain the initial mathematical formulas as q approaches 1. The commencement of the analysis of q -calculus can be dated back to the era of Euler (1707–1783), who first initiated the q -calculus in the tracks of Newton’s work on infinite series. Subsequently, Jackson [49] launched the concept of q -integrals and studied it in a systematic way. The aforementioned results lead to an intensive investigation on q -calculus in the twentieth century. The idea of q -calculus is used in numerous areas in mathematics and physics, especially in orthogonal polynomials, number theory, hypergeometric functions, mechanics and relativity theory. The concept of q -derivatives over the definite interval $[\phi, \psi]$ of \mathbb{R} is introduced by Tariboon et al. [50,51], and they addressed several problems on quantum analogs such as Hölder inequality, Ostrowski inequality, Cauchy-Schwarz inequality, Grüss-Chebyshev inequality, Grüss inequality and other integral inequalities by classical convexity.

From the last few years, q -calculus has become an interesting topic for many researchers and several new results have been established in the literature [52–58]. Furthermore, Tunç and Göv [59,60] derived the notion of (p, q) -calculus on the intervals $[\phi, \psi]$ of \mathbb{R} , found the formulae for (p, q) -derivative and (p, q) -integral and established their several fundamental properties. The results that depend on (p, q) -calculus are the Minkowski inequality, Hölder inequality, Grüss and Grüss-Chebyshev inequality and many others. Kunt et al. [61] gave the generalized (p, q) -Hermite-Hadamard inequalities on the finite interval and some important results which are connected with (p, q) -midpoint-type inequality. Recently, (p, q) -calculus has been the subject of intensive research, and its refinements and generalizations can be found in the literature [61,62].

Now, we recall the definitions and theorems for (p, q) -derivative and (p, q) -integral.

Definition 1.1. [59] Let $0 < q < p \leq 1$ and $g : [\phi, \psi] \rightarrow \mathbb{R}$ be a continuous function. Then the (p, q) -derivative of g at $\lambda \in [\phi, \psi]$ is defined by

$${}_{\phi}D_{p,q}g(\lambda) = \frac{g(p\lambda + (1-p)\phi) - g(q\lambda + (1-q)\phi)}{(p-q)(\lambda - \phi)} \quad (\lambda \neq \phi)$$

and

$${}_{\phi}D_{p,q}g(\phi) = \lim_{\lambda \rightarrow \phi} D_{p,q}g(\lambda).$$

Example 1.2. Define the function $g : [\phi, \psi] \rightarrow \mathbb{R}$ by $g(\lambda) = 2\lambda^2 + 1$ with $0 < q < p \leq 1$. Then for $\lambda \neq \phi$ we have

$$\begin{aligned} {}_{\phi}D_{p,q}(2\lambda^2 + 1) &= \frac{(2p\lambda + (1-p)\phi)^2 + 1 - (2q\lambda + (1-q)\phi)^2 + 1}{(p-q)(\lambda - \phi)} \\ &= \frac{2[2]_{p,q}\lambda^2 + 4\phi\lambda[1 - [2]_{p,q}] + 2\phi^2[[2]_{p,q} - 2]}{(\lambda - \phi)} \\ &= \frac{2\lambda[2]_{p,q}(\lambda - \phi) - 2\phi[2]_{p,q}(\lambda - \phi) + 4\phi(\lambda - \phi)}{(\lambda - \phi)} \\ &= 2[2]_{p,q}(\lambda - \phi) + 4\phi. \end{aligned}$$

Definition 1.3. [59] Let $0 < q < p \leq 1$ and $g : [\phi, \psi] \rightarrow \mathbb{R}$ be a continuous function. Then the (p, q) -integral on $[\phi, \psi]$ is defined by

$$\int_{\phi}^{\lambda} g(x)_{\phi}d_{p,q}x = (p-q)(\lambda - \phi) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} g\left(\frac{q^n}{p^{n+1}}\lambda + \left(1 - \frac{q^n}{p^{n+1}}\right)\phi\right)$$

for $\lambda \in [\phi, \psi]$.

If $c \in (\phi, \lambda)$, then the (p, q) -definite integral on $[c, \lambda]$ can be expressed as

$$\int_c^{\lambda} g(x)_{\phi}d_{p,q}x = \int_{\phi}^{\lambda} g(x)_{\phi}d_{p,q}x - \int_{\phi}^c g(x)_{\phi}d_{p,q}x.$$

Example 1.4. Define the function $g : [\phi, \psi] \rightarrow \mathbb{R}$ by $g(x) = 4x + 1$ with $0 < q < p \leq 1$. Then one has

$$\begin{aligned} \int_{\phi}^{\lambda} (4x + 1)_{\phi}d_{p,q}x &= (p-q)(\lambda - \phi) \left(4 \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} \left(\frac{q^n}{p^{n+1}}\lambda + \left(1 - \frac{q^n}{p^{n+1}}\right)\phi \right) + \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} \right) \\ &= \frac{(\lambda - \phi)[4(\lambda - \phi(1 - p - q)) + [2]_{p,q}]}{[2]_{p,q}}. \end{aligned}$$

Theorem 1.5. [61] Let $0 < q < p \leq 1$ and $g : [\phi, \psi] \rightarrow \mathbb{R}$ be a convex differentiable function on $[\phi, \psi]$. Then

$$g\left(\frac{q\phi + p\psi}{[2]_{p,q}}\right) \leq \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x)_{\phi}d_{p,q}x \leq \frac{qg(\phi) + pg(\psi)}{[2]_{p,q}}.$$

The definition of (α, m) -convex function was presented by Miheřan in [63] and is stated as follows.

Definition 1.6. Let $\alpha, m \in (0, 1]$ and $\psi^* > 0$. Then the function $g : [0, \psi^*] \rightarrow \mathbb{R}$ is said to be (α, m) -convex if the inequality

$$g(x\lambda + m(1 - \lambda)y) \leq \lambda^{\alpha}g(x) + m(1 - \lambda^{\alpha})g(y)$$

holds for all $x, y \in [0, \psi^*]$ and $\lambda \in [0, 1]$.

Zhang et al. [64] investigated some inequalities about q -differentiable convex and quasi-convex functions which are linked with the different types of inequalities in q -calculus.

Lemma 1.7. [64] Let $0 < q < 1$ and $g : [\phi, \psi] \rightarrow \mathbb{R}$ be a q -differentiable function on (ϕ, ψ) such that ${}_{\phi}D_qg$ is continuous and integrable on $[\phi, \psi]$. Then

$$\begin{aligned} & \gamma[\mu g(\psi) + (1 - \mu)g(\phi)] + (1 - \gamma)g(\mu\psi + (1 - \mu)\phi) - \frac{1}{\psi - \phi} \int_{\phi}^{\psi} g(x)_{\phi} d_q x \\ &= (\psi - \phi) \left[\int_0^{\mu} (q\lambda + \gamma\mu - \gamma)_{\phi} D_q g(\lambda\psi + (1 - \lambda)\phi)_0 d_q \lambda + \int_{\mu}^1 (q\lambda + \gamma\mu - 1)_{\phi} D_q g(\lambda\psi + (1 - \lambda)\phi)_0 d_q \lambda \right] \end{aligned}$$

for all $\gamma, \mu \in [0, 1]$.

The main purpose of the article is to provide an identity, which is the generalization of an identity presented in Lemma 1.7, and establish the (p, q) -analogues of different types of integral inequalities via the (p, q) -differentiable (α, m) -convex functions. By using the new identity with distinct parameters we obtain some new (p, q) -quantum error estimations for different types of inequalities such as the midpoint-type, the Simpson-type, the average of midpoint-trapezoid-type and the trapezoid-type inequalities via (α, m) -convexity.

2 Auxiliary results

In order to obtain different types of integral inequalities through (p, q) -differentiable (α, m) -convex functions, we need several lemmas which we present in this section.

Lemma 2.1. *Let $0 < q < p \leq 1$ and $g : [\phi, \psi] \rightarrow \mathbb{R}$ be a (p, q) -differentiable function on (ϕ, ψ) such that ${}_{\phi}D_{p,q}g$ is continuous and integrable on $[\phi, \psi]$. Then*

$$\begin{aligned} & \gamma[p\mu g(\psi) + (1 - p\mu)g(\phi)] + (1 - \gamma)g(p\mu\psi + (1 - p\mu)\phi) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x)_{\phi} d_{p,q} x \\ &= (\psi - \phi) \left[\int_0^{p\mu} (q\lambda + \gamma p\mu - \gamma)_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q} \lambda + \int_{p\mu}^1 (q\lambda + \gamma p\mu - 1)_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q} \lambda \right] \end{aligned}$$

for all $\gamma, \mu \in [0, 1]$.

Proof. By an identical transformation, we get

$$\begin{aligned} & (\psi - \phi) \left[\int_0^{p\mu} (q\lambda + \gamma p\mu - \gamma)_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q} \lambda + \int_{p\mu}^1 (q\lambda + \gamma p\mu - 1)_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q} \lambda \right] \\ &= (\psi - \phi) \left[\int_0^1 (q\lambda + \gamma p\mu - 1)_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q} \lambda + (1 - \gamma) \int_0^{p\mu} {}_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q} \lambda \right]. \end{aligned} \tag{2.1}$$

Applying Definitions 1.1 and 1.3, we have

$$\begin{aligned} & \int_0^1 \lambda {}_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q} \lambda \\ &= \int_0^1 \frac{g(p\lambda\psi + (1 - p\lambda)\phi) - g(q\lambda\psi + (1 - q\lambda)\phi)}{(p - q)(\psi - \phi)} {}_0 d_{p,q} \lambda \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{\psi - \phi} \left[\sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} g \left(\frac{q^n}{p^n} \psi + \left(1 - \frac{q^n}{p^n} \right) \phi \right) - \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} g \left(\frac{q^{n+1}}{p^{n+1}} \psi + \left(1 - \frac{q^{n+1}}{p^{n+1}} \right) \phi \right) \right] \\
 &= \frac{1}{\psi - \phi} \left[\frac{1}{p} \sum_{n=0}^{\infty} \frac{q^n}{p^n} g \left(\frac{q^n}{p^n} \psi + \left(1 - \frac{q^n}{p^n} \right) \phi \right) - \frac{1}{q} \sum_{n=1}^{\infty} \frac{q^n}{p^n} g \left(\frac{q^n}{p^n} \psi + \left(1 - \frac{q^n}{p^n} \right) \phi \right) \right] \\
 &= \frac{1}{\psi - \phi} \left[\frac{1}{q} g(\psi) - \left(\frac{1}{q} - \frac{1}{p} \right) \times \sum_{n=0}^{\infty} \frac{q^n}{p^n} g \left(\frac{q^n}{p^n} \psi + \left(1 - \frac{q^n}{p^n} \right) \phi \right) \right] \\
 &= \frac{1}{q(\psi - \phi)} g(\psi) - \frac{1}{pq(\psi - \phi)^2} \int_{\phi}^{p\psi + (1-p)\phi} g(x)_{\phi} d_{p,q} x,
 \end{aligned} \tag{2.2}$$

$$\begin{aligned}
 &\int_{\phi}^1 D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_{\phi} d_{p,q} \lambda \\
 &= \int_0^1 \frac{g(p\lambda\psi + (1 - p\lambda)\phi) - g(q\lambda\psi + (1 - q\lambda)\phi)}{\lambda(p - q)(\psi - \phi)} d_{p,q} \lambda \\
 &= \frac{1}{\psi - \phi} \left[\sum_{n=0}^{\infty} g \left(\frac{q^n}{p^n} \psi + \left(1 - \frac{q^n}{p^n} \right) \phi \right) - \sum_{n=0}^{\infty} g \left(\frac{q^{n+1}}{p^{n+1}} \psi + \left(1 - \frac{q^{n+1}}{p^{n+1}} \right) \phi \right) \right] \\
 &= \frac{g(\psi) - g(\phi)}{\psi - \phi},
 \end{aligned} \tag{2.3}$$

$$\begin{aligned}
 &\int_0^{p\mu} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_{\phi} d_{p,q} \lambda \\
 &= \int_0^{p\mu} \frac{g(p\lambda\psi + (1 - p\lambda)\phi) - g(q\lambda\psi + (1 - q\lambda)\phi)}{\lambda(p - q)(\psi - \phi)} d_{p,q} \lambda \\
 &= \frac{1}{\psi - \phi} \left[\sum_{n=0}^{\infty} g \left(\frac{q^n}{p^n} p\mu\psi + \left(1 - \frac{q^n}{p^n} p\mu \right) \phi \right) - \sum_{n=0}^{\infty} g \left(\frac{q^{n+1}}{p^{n+1}} p\mu\psi + \left(1 - \frac{q^{n+1}}{p^{n+1}} p\mu \right) \phi \right) \right] \\
 &= \frac{g(p\mu\psi + (1 - p\mu)\phi) - g(\phi)}{\psi - \phi}.
 \end{aligned} \tag{2.4}$$

Substituting (2.2), (2.3) and (2.4) into (2.1), we obtain the desired result. □

Remark 2.1. The following statements are true under the conditions of Lemma 2.1.

(1) If $\mu = 0$, then we get

$$g(\phi) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x)_{\phi} d_{p,q} x = (\psi - \phi) \int_0^1 (q\lambda - 1)_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_{\phi} d_{p,q} \lambda.$$

(2) If $p = \mu = 1$, then we have

$$g(\psi) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x)_{\phi} d_{p,q} x = (\psi - \phi) \int_0^1 q\lambda_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_{\phi} d_{p,q} \lambda.$$

(3) If $\mu = 1/[2]_{p,q}$, then one has

$$y \frac{qg(\phi) + pg(\psi)}{[2]_{p,q}} + (1 - y) g \left(\frac{q\phi + p\psi}{[2]_{p,q}} \right) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x)_{\phi} d_{p,q} x$$

$$\begin{aligned}
 &= (\psi - \phi) \left[\int_0^{\frac{p}{[2]_{p,q}}} \left(q\lambda - \frac{\gamma q}{[2]_{p,q}} \right) {}_{\phi}D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q}\lambda \right. \\
 &\quad \left. + \int_{\frac{p}{[2]_{p,q}}}^1 \left(q\lambda + \frac{p(\gamma - 1) - q}{[2]_{p,q}} \right) {}_{\phi}D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q}\lambda \right].
 \end{aligned}$$

Remark 2.2. If all the conditions of Lemma 2.1 are satisfied, then the following four statements are true:

(1) If $\gamma = 0$, then we get

$$\begin{aligned}
 &g(p\mu\psi + (1 - p\mu)\phi) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x)_{\phi} d_{p,q}x \\
 &= (\psi - \phi) \left[\int_0^{p\mu} q\lambda {}_{\phi}D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q}\lambda + \int_{p\mu}^1 (q\lambda - 1) {}_{\phi}D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q}\lambda \right].
 \end{aligned} \tag{2.5}$$

Let $\mu = 1/[2]_{p,q}$ in (2.5). Then we acquire the midpoint-type integral identity

$$\begin{aligned}
 &g\left(\frac{q\phi + p\psi}{[2]_{p,q}}\right) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x)_{\phi} d_{p,q}x \\
 &= (\psi - \phi) \left[\int_0^{\frac{p}{[2]_{p,q}}} q\lambda {}_{\phi}D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q}\lambda + \int_{[2]_{p,q}}^1 (q\lambda - 1) {}_{\phi}D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q}\lambda \right],
 \end{aligned} \tag{2.6}$$

which was proposed by Kunt et al. in [61], and equation (2.6) leads to Lemma 11 of [65] if $p = 1$.

(2) If $\gamma = 1/3$, then we get

$$\begin{aligned}
 &\frac{1}{3} [p\mu g(\psi) + (1 - p\mu)g(\phi) + 2g(p\mu\psi + (1 - p\mu)\phi)] - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x)_{\phi} d_{p,q}x \\
 &= (\psi - \phi) \left[\int_0^{p\mu} \left(q\lambda + \frac{p\mu - 1}{3} \right) {}_{\phi}D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q}\lambda \right. \\
 &\quad \left. + \int_{p\mu}^1 \left(q\lambda + \frac{p\mu - 3}{3} \right) {}_{\phi}D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q}\lambda \right].
 \end{aligned} \tag{2.7}$$

In particular, if $\mu = 1/[2]_{p,q}$, then equation (2.7) leads to the Simpson-type integral identity

$$\begin{aligned}
 &\frac{1}{3} \left[\frac{qg(\phi) + pg(\psi)}{[2]_{p,q}} + 2g\left(\frac{q\phi + p\psi}{[2]_{p,q}}\right) \right] - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x)_{\phi} d_{p,q}x \\
 &= (\psi - \phi) \left\{ \int_0^{\frac{p}{[2]_{p,q}}} \left(q\lambda - \frac{q}{3[2]_{p,q}} \right) {}_{\phi}D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q}\lambda \right. \\
 &\quad \left. + \int_{\frac{p}{[2]_{p,q}}}^1 \left(q\lambda - \frac{2p + 3q}{3[2]_{p,q}} \right) {}_{\phi}D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q}\lambda \right\}.
 \end{aligned}$$

(3) If $\gamma = 1/2$, then one has

$$\begin{aligned} & \frac{1}{2}[p\mu g(\psi) + (1 - p\mu)g(\phi) + g(p\mu\psi + (1 - p\mu)\phi)] - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi+(1-p)\phi} g(x)_{\phi} d_{p,q}x \\ &= (\psi - \phi) \left\{ \int_0^{p\mu} \left(q\lambda + \frac{p\mu - 1}{2} \right)_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_{\phi} d_{p,q}\lambda \right. \\ & \quad \left. + \int_{\frac{p}{2}}^1 \left(q\lambda + \frac{p\mu - 2}{2} \right)_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_{\phi} d_{p,q}\lambda \right\}. \end{aligned} \tag{2.8}$$

In particular, if $\mu = 1/[2]_{p,q}$, then equation (2.8) gives the average of midpoint-trapezoid-type integral identity

$$\begin{aligned} & \frac{1}{2} \left[\frac{qg(\phi) + pg(\psi)}{[2]_{p,q}} + g \left(\frac{q\phi + p\psi}{[2]_{p,q}} \right) \right] - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi+(1-p)\phi} g(x)_{\phi} d_{p,q}x \\ &= (\psi - \phi) \left\{ \int_0^{\frac{p}{[2]_{p,q}}} \left(q\lambda - \frac{q}{2[2]_{p,q}} \right)_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_{\phi} d_{p,q}\lambda \right. \\ & \quad \left. + \int_{\frac{p}{[2]_{p,q}}}^1 \left(q\lambda - \frac{p + 2q}{2[2]_{p,q}} \right)_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_{\phi} d_{p,q}\lambda \right\}. \end{aligned}$$

(4) Let $\gamma = 1$. Then we get

$$\begin{aligned} & p\mu g(\psi) + (1 - p\mu)g(\phi) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi+(1-p)\phi} g(x)_{\phi} d_{p,q}x \\ &= (\psi - \phi) \int_0^1 (q\lambda + p\mu - 1)_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_{\phi} d_{p,q}\lambda. \end{aligned} \tag{2.9}$$

Let $\mu = 1/[2]_{p,q}$. Then equation (2.9) leads to the trapezoid-type integral identity

$$\begin{aligned} & \frac{qg(\phi) + pg(\psi)}{[2]_{p,q}} - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi+(1-p)\phi} g(x)_{\phi} d_{p,q}x \\ &= (\psi - \phi) \int_0^1 \left(q\lambda - \frac{q}{[2]_{p,q}} \right)_{\phi} D_{p,q} g(\lambda\psi + (1 - \lambda)\phi)_{\phi} d_{p,q}\lambda, \end{aligned} \tag{2.10}$$

which was proposed by Latif et al. in [66].

In particular, Lemma 3.1 of [55] can be derived from equation (2.10) if we take $p = 1$.

The following Lemma 2.2 can be obtained immediately from Definition 1.3.

Lemma 2.2. Let $0 < q < p \leq 1$, $0 \leq \mu \leq 1$ and $\xi \in [0, \infty)$. Then we have

$$\int_0^{p\mu} \lambda^{\xi} d_{p,q}\lambda = (p - q)\mu^{\xi+1} \sum_{n=0}^{\infty} \left(\frac{q}{p} \right)^{(\xi+1)n} = \frac{\mu^{\xi+1}(p - q)p^{\xi+1}}{p^{\xi+1} - q^{\xi+1}}$$

and

$$\int_0^{p\mu} (1 - \lambda)^\xi {}_0d_{p,q}\lambda = (p - q)\mu \sum_{n=0}^\infty \frac{q^n}{p^n} \left(1 - \frac{q^n}{p^n}\mu\right)^\xi.$$

Lemma 2.3. Let $0 < q < p \leq 1$, $\gamma, \mu \in [0, 1]$ and $\xi \in [0, \infty)$. Then we get

$$\begin{aligned} v_1(\gamma, p\mu, \xi) &= \int_0^{p\mu} \lambda^\xi |q\lambda - (\gamma - \gamma p\mu)| {}_0d_{p,q}\lambda \\ &= \begin{cases} \frac{\mu^{\xi+1}(p - q)p^{\xi+1}(\gamma - \gamma p\mu)}{p^{\xi+1} - q^{\xi+1}} - \frac{q\mu^{\xi+2}(p - q)p^{\xi+2}}{p^{\xi+2} - q^{\xi+2}}, & (\gamma + q)p\mu \leq \gamma, \\ \left[\frac{2(p - q)(\gamma - \gamma p\mu)^{\xi+2}(p^{\xi+2} - q^{\xi+2} - p^{\xi+1} + q^{\xi+1})}{q^{\xi+1}(p^{\xi+1} - q^{\xi+1})(p^{\xi+2} - q^{\xi+2})} \right. \\ \left. + \frac{q\mu^{\xi+2}(p - q)p^{\xi+2}}{p^{\xi+2} - q^{\xi+2}} - \frac{\mu^{\xi+1}(p - q)p^{\xi+1}(\gamma - \gamma p\mu)}{p^{\xi+1} - q^{\xi+1}} \right], & (\gamma + q)p\mu > \gamma \end{cases} \end{aligned}$$

and

$$\begin{aligned} v_2(\gamma, p\mu, \xi) &= \int_0^{p\mu} (1 - \lambda)^\xi |(q\lambda - (\gamma - \gamma p\mu))| {}_0d_{p,q}\lambda \\ &= \begin{cases} (p - q)\mu \sum_{n=0}^\infty \frac{q^n}{p^n} \left(\gamma - \gamma p\mu - \frac{q^{n+1}}{p^{n+1}}\mu\right) \left(1 - \frac{q^n}{p^n}\mu\right)^\xi, & (\gamma + q)p\mu \leq \gamma, \\ \left[2(p - q)(\gamma - \gamma p\mu)^2 \sum_{n=0}^\infty \frac{q^{n-1}}{p^{n+1}} \left(1 - \frac{q^n}{p^{n+1}}\right) \left(1 - \frac{q^{n-1}}{p^{n+1}}(\gamma - \gamma p\mu)\right)^\xi \right. \\ \left. - (p - q)\mu \sum_{n=0}^\infty \frac{q^n}{p^n} \left(\gamma - \gamma p\mu - \frac{q^{n+1}}{p^{n+1}}\mu\right) \left(1 - \frac{q^n}{p^n}\mu\right)^\xi \right], & (\gamma + q)p\mu > \gamma. \end{cases} \end{aligned}$$

Proof. If $(\gamma + q)p\mu \leq \gamma$, then it follows from Lemma 2.1 that

$$\begin{aligned} \int_0^{p\mu} \lambda^\xi |q\lambda - (\gamma - \gamma p\mu)| {}_0d_{p,q}\lambda &= \int_0^{p\mu} [(\gamma - \gamma p\mu)\lambda^\xi - q\lambda^{\xi+1}] {}_0d_{p,q}\lambda \\ &= \frac{\mu^{\xi+1}(p - q)p^{\xi+1}(\gamma - \gamma p\mu)}{p^{\xi+1} - q^{\xi+1}} - \frac{q\mu^{\xi+2}(p - q)p^{\xi+2}}{p^{\xi+2} - q^{\xi+2}}. \end{aligned}$$

If $(\gamma + q)p\mu > \gamma$, then from Lemma 2.1 we get

$$\begin{aligned} \int_0^{p\mu} \lambda^\xi |q\lambda - (\gamma - \gamma p\mu)| {}_0d_{p,q}\lambda &= \int_0^{\frac{\gamma - \gamma p\mu}{q}} [(\gamma - \gamma p\mu)\lambda^\xi - q\lambda^{\xi+1}] {}_0d_{p,q}\lambda + \int_{\frac{\gamma - \gamma p\mu}{q}}^{p\mu} [q\lambda^{\xi+1} - (\gamma - \gamma p\mu)\lambda^\xi] {}_0d_{p,q}\lambda \\ &= 2 \int_0^{\frac{\gamma - \gamma p\mu}{q}} [(\gamma - \gamma p\mu)\lambda^\xi - q\lambda^{\xi+1}] {}_0d_{p,q}\lambda + \int_0^{p\mu} [q\lambda^{\xi+1} - (\gamma - \gamma p\mu)\lambda^\xi] {}_0d_{p,q}\lambda \\ &= \frac{2(p - q)(\gamma - \gamma p\mu)^{\xi+2}(p^{\xi+2} - q^{\xi+2} - p^{\xi+1} + q^{\xi+1})}{q^{\xi+1}(p^{\xi+1} - q^{\xi+1})(p^{\xi+2} - q^{\xi+2})} \\ &\quad + \frac{q\mu^{\xi+2}(p - q)p^{\xi+2}}{p^{\xi+2} - q^{\xi+2}} - \frac{\mu^{\xi+1}(p - q)p^{\xi+1}(\gamma - \gamma p\mu)}{p^{\xi+1} - q^{\xi+1}}. \end{aligned}$$

Similarly, we also get

$$\int_0^{p\mu} (1-\lambda)^\xi |q\lambda - (\gamma - \gamma p\mu)|_0 d_{p,q}\lambda = \begin{cases} (p-q)\mu \sum_{n=0}^\infty \frac{q^n}{p^n} \left(\gamma - \gamma p\mu - \frac{q^{n+1}}{p^n}\mu\right) \left(1 - \frac{q^n}{p^n}\mu\right)^\xi, & (\gamma + q)p\mu \leq \gamma, \\ \left[2(p-q)(\gamma - \gamma p\mu)^2 \sum_{n=0}^\infty \frac{q^{n-1}}{p^{n+1}} \left(1 - \frac{q^n}{p^{n+1}}\right) \left(1 - \frac{q^{n-1}}{p^{n+1}}(\gamma - \gamma p\mu)\right)^\xi \right. \\ \left. - (p-q)\mu \sum_{n=0}^\infty \frac{q^n}{p^n} \left(\gamma - \gamma p\mu - \frac{q^{n+1}}{p^n}\mu\right) \left(1 - \frac{q^n}{p^n}\mu\right)^\xi \right], & (\gamma + q)p\mu > \gamma, \end{cases}$$

which completes the proof of Lemma 2.3. □

The following Lemmas 2.4–2.9 can be obtained by using the definition of q -integrals, we omit the details of their proofs.

Lemma 2.4. *Let $0 < q < p \leq 1$, $\gamma, \mu \in [0, 1]$ and $\xi \in [0, \infty)$. Then we have*

$$v_3(\gamma, p\mu, \xi) = \int_0^1 \lambda^\xi |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda = \begin{cases} \frac{(p-q)p^{\xi+1}(1-\gamma p\mu)}{p^{\xi+1} - q^{\xi+1}} - \frac{q(p-q)p^{\xi+2}}{p^{\xi+2} - q^{\xi+2}}, & \gamma p\mu + q \leq 1, \\ \left[\frac{2(p-q)(1-\gamma p\mu)^{\xi+2}(p^{\xi+2} - q^{\xi+2} - p^{\xi+1} + q^{\xi+1})}{q^{\xi+1}(p^{\xi+1} - q^{\xi+1})(p^{\xi+2} - q^{\xi+2})} \right. \\ \left. + \frac{q(p-q)p^{\xi+2}}{p^{\xi+2} - q^{\xi+2}} - \frac{(p-q)p^{\xi+1}(1-\gamma p\mu)}{p^{\xi+1} - q^{\xi+1}} \right], & \gamma p\mu + q > 1 \end{cases}$$

and

$$v_4(\gamma, p\mu, \xi) = \int_0^1 (1-\lambda)^\xi |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda = \begin{cases} (p-q) \sum_{n=0}^\infty \frac{q^n}{p^{n+1}} \left(1 - \gamma p\mu - \frac{q^{n+1}}{p^{n+1}}\right) \left(1 - \frac{q^n}{p^{n+1}}\right)^\xi, & \gamma p\mu + q \leq 1, \\ \left[2(p-q)(1-\gamma p\mu)^2 \sum_{n=0}^\infty \frac{q^{n-1}}{p^{n+1}} \left(1 - \frac{q^n}{p^{n+1}}\right) \left(1 - \frac{q^{n-1}}{p^{n+1}}(1-\gamma p\mu)\right)^\xi \right. \\ \left. - (p-q) \sum_{n=0}^\infty \frac{q^n}{p^{n+1}} \left(1 - \gamma p\mu - \frac{q^{n+1}}{p^{n+1}}\right) \left(1 - \frac{q^n}{p^{n+1}}\right)^\xi \right], & \gamma p\mu + q > 1. \end{cases}$$

Lemma 2.5. *Let $0 < q < p \leq 1$, $\gamma, \mu \in [0, 1]$ and $\xi \in [0, \infty)$. Then one has*

$$v_5(\gamma, p\mu, \xi) = \int_0^{p\mu} \lambda^\xi |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda = \begin{cases} \frac{\mu^{\xi+1}(p-q)p^{\xi+1}(1-\gamma p\mu)}{p^{\xi+1} - q^{\xi+1}} - \frac{q\mu^{\xi+2}(p-q)p^{\xi+2}}{p^{\xi+2} - q^{\xi+2}}, & (\gamma + q)p\mu \leq 1, \\ \left[\frac{2(p-q)(1-\gamma p\mu)^{\xi+2}(p^{\xi+2} - q^{\xi+2} - p^{\xi+1} + q^{\xi+1})}{q^{\xi+1}(p^{\xi+1} - q^{\xi+1})(p^{\xi+2} - q^{\xi+2})} \right. \\ \left. + \frac{q\mu^{\xi+2}(p-q)p^{\xi+2}}{p^{\xi+2} - q^{\xi+2}} - \frac{\mu^{\xi+1}(p-q)p^{\xi+1}(1-\gamma p\mu)}{p^{\xi+1} - q^{\xi+1}} \right], & (\gamma + q)p\mu > 1 \end{cases}$$

and

$$\begin{aligned}
 v_6(\gamma, p\mu, \xi) &= \int_0^{p\mu} (1-\lambda)^\xi |q\lambda - (1-\gamma\mu)|_0 d_{p,q}\lambda \\
 &= \begin{cases} (p-q)\mu \sum_{n=0}^{\infty} \frac{q^n}{p^n} \left(1 - \gamma p\mu - \frac{q^{n+1}}{p^n} \mu\right) \left(1 - \frac{q^n}{p^{n+1}}\right)^\xi, & (\gamma+q)p\mu \leq 1, \\ \left[2(p-q)(1-\gamma\mu)^2 \sum_{n=0}^{\infty} \frac{q^{n-1}}{p^{n+1}} \left(1 - \frac{q^n}{p^{n+1}}\right) \left(1 - \frac{q^{n-1}}{p^{n+1}}(1-\gamma p\mu)\right)^\xi \right. \\ \left. - (p-q)\mu \sum_{n=0}^{\infty} \frac{q^n}{p^n} \left(1 - \gamma p\mu - \frac{q^{n+1}}{p^n} \mu\right) \left(1 - \frac{q^n}{p^{n+1}}\right)^\xi \right], & (\gamma+q)p\mu > 1. \end{cases}
 \end{aligned}$$

Lemma 2.6. Let $0 < q < p \leq 1$ and $\gamma, \mu \in [0, 1]$. Then we have

$$\begin{aligned}
 v_7(\gamma, p\mu) &= \int_0^{p\mu} |q\lambda - (\gamma - \gamma p\mu)|_0 d_{p,q}\lambda \\
 &= \begin{cases} \gamma p\mu(1-p\mu) - \frac{q\mu^2 p^2}{[2]_{p,q}}, & (\gamma+q)p\mu \leq \gamma, \\ \frac{2(\gamma - \gamma p\mu)^2([2]_{p,q} - 1)}{q[2]_{p,q}} + \frac{q\mu^2 p^2}{[2]_{p,q}} - \gamma p\mu(1-p\mu), & (\gamma+q)p\mu > \gamma. \end{cases}
 \end{aligned}$$

Lemma 2.7. Let $0 < q < p \leq 1$ and $\gamma, \mu \in [0, 1]$. Then we get

$$\begin{aligned}
 v_8(\gamma, p\mu) &= \int_0^1 |q\lambda - (1-\gamma p\mu)|_0 d_{p,q}\lambda \\
 &= \begin{cases} \frac{p}{[2]_{p,q}} - \gamma p\mu, & \gamma p\mu + q \leq 1, \\ \frac{2(1-\gamma p\mu)^2([2]_{p,q} - 1)}{q[2]_{p,q}} + \gamma p\mu - \frac{p}{[2]_{p,q}}, & \gamma p\mu + q > 1. \end{cases}
 \end{aligned}$$

Lemma 2.8. Let $0 < q < p \leq 1$ and $\gamma, \mu \in [0, 1]$. Then one has

$$v_9(\gamma, p\mu) = \int_0^{p\mu} |q\lambda - (1-\gamma p\mu)|_0 d_{p,q}\lambda = \begin{cases} p\mu(1-\gamma p\mu) - \frac{q\mu^2 p^2}{[2]_{p,q}}, & (\gamma+q)p\mu \leq 1, \\ \frac{2(1-\gamma p\mu)^2([2]_{p,q} - 1)}{q[2]_{p,q}} + \frac{q\mu^2 p^2}{[2]_{p,q}} - p\mu(1-\gamma p\mu), & (\gamma+q)p\mu > 1. \end{cases}$$

Lemma 2.9. Let $0 < q < p \leq 1$, $\gamma, \mu \in [0, 1]$ and $\sigma \in [1, \infty)$. Then we get

$$v_{10}(\gamma, p\mu) = \int_0^1 |q\lambda - (1-\gamma p\mu)|^\sigma_0 d_{p,q}\lambda = \begin{cases} (p-q) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} \left(1 - \gamma p\mu - \frac{q^{n+1}}{p^{n+1}}\right)^\sigma, & 0 \leq \gamma p\mu \leq 1-q, \\ \left[(p-q)(1-\gamma p\mu)^{\sigma+1} \sum_{n=0}^{\infty} \frac{q^{n-1}}{p^{n+1}} \left(1 - \frac{q^n}{p^{n+1}}\right)^\sigma \right. \\ \left. + (p-q) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} \left(\frac{q^{n+1}}{p^{n+1}} - 1 + \gamma p\mu\right)^\sigma \right. \\ \left. - (p-q)(1-\gamma p\mu)^{\sigma+1} \sum_{n=0}^{\infty} \frac{q^{n-1}}{p^{n+1}} \left(\frac{q^n}{p^{n+1}} - 1\right)^\sigma \right], & 1-q < \gamma p\mu \leq 1. \end{cases}$$

3 Main results

Theorem 3.1. Let $0 \leq \phi < \psi < \infty$, $0 < q < p \leq 1$, $\alpha, m \in (0, 1]$ and $g : J \supset (0, \infty) \rightarrow \mathbb{R}$ be a (p, q) -differentiable function on J° (the interior of J) such that ${}_{\phi}D_{p,q}g$ is continuous and integrable on $\left[0, \frac{\psi}{m}\right]$ and $|{}_{\phi}D_{p,q}g|$ is (α, m) -convex on $\left[0, \frac{\psi}{m}\right]$. Then the inequality

$$\left| \gamma[p\mu g(\psi) + (1 - p\mu)g(\phi)] + (1 - \gamma)g(p\mu\psi + (1 - p\mu)\phi) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x) {}_{\phi}d_{p,q}x \right| \leq \min[H_1(\gamma, p\mu, \alpha, m), H_2(\gamma, p\mu, \alpha, m)]$$

holds for all $\gamma, \mu \in [0, 1]$, where

$$\begin{aligned} H_1(\gamma, p\mu, \alpha, m) &= (\psi - \phi) \left\{ [v_1(\gamma, p\mu, \alpha) + v_3(\gamma, p\mu, \alpha) - v_5(\gamma, p\mu, \alpha)] |{}_{\phi}D_{p,q}g(\psi)| + m[v_7(\gamma, p\mu) \right. \\ &\quad \left. + v_8(\gamma, p\mu) - v_9(\gamma, p\mu) - v_1(\gamma, p\mu, \alpha) - v_3(\gamma, p\mu, \alpha) + v_5(\gamma, p\mu, \alpha)] \left| {}_{\phi}D_{p,q}g\left(\frac{\phi}{m}\right) \right| \right\}, \\ H_2(\gamma, p\mu, \alpha, m) &= (\psi - \phi) \left\{ [v_2(\gamma, p\mu, \alpha) + v_4(\gamma, p\mu, \alpha) - v_6(\gamma, p\mu, \alpha)] |{}_{\phi}D_{p,q}g(\phi)| + m[v_7(\gamma, p\mu) \right. \\ &\quad \left. + v_8(\gamma, p\mu) - v_9(\gamma, p\mu) - v_2(\gamma, p\mu, \alpha) - v_4(\gamma, p\mu, \alpha) + v_6(\gamma, p\mu, \alpha)] \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right| \right\}. \end{aligned}$$

Proof. From Lemma 2.1, the property of the modulus and the (α, m) -convexity of $|{}_{\phi}D_{p,q}g|$ we have

$$\begin{aligned} &\left| \gamma[p\mu g(\psi) + (1 - p\mu)g(\phi)] + (1 - \gamma)g(p\mu\psi + (1 - p\mu)\phi) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x) {}_{\phi}d_{p,q}x \right| \\ &\leq (\psi - \phi) \left[\int_0^{p\mu} |q\lambda + \gamma p\mu - \gamma| |{}_{\phi}D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)| {}_{\phi}d_{p,q}\lambda \right. \\ &\quad \left. + \int_{p\mu}^1 |q\lambda + \gamma p\mu - 1| |{}_{\phi}D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)| {}_{\phi}d_{p,q}\lambda \right] \\ &\leq (\psi - \phi) \left[\int_0^{p\mu} |q\lambda - (\gamma - \gamma p\mu)| \left[\lambda^\alpha |{}_{\phi}D_{p,q}g(\psi)| + m(1 - \lambda^\alpha) \left| {}_{\phi}D_{p,q}g\left(\frac{\phi}{m}\right) \right| \right] {}_{\phi}d_{p,q}\lambda \right. \\ &\quad \left. + \int_{p\mu}^1 |q\lambda - (1 - \gamma p\mu)| \left[\lambda^\alpha |{}_{\phi}D_{p,q}g(\psi)| + m(1 - \lambda^\alpha) \left| {}_{\phi}D_{p,q}g\left(\frac{\phi}{m}\right) \right| \right] {}_{\phi}d_{p,q}\lambda \right] \\ &= (\psi - \phi) \left[\int_0^{p\mu} \lambda^\alpha |q\lambda - (\gamma - \gamma p\mu)| {}_{\phi}d_{p,q}\lambda + \int_0^1 \lambda^\alpha |q\lambda - (1 - \gamma p\mu)| {}_{\phi}d_{p,q}\lambda \right. \\ &\quad \left. - \int_0^{p\mu} \lambda^\alpha |q\lambda - (1 - \gamma p\mu)| {}_{\phi}d_{p,q}\lambda \right] |{}_{\phi}D_{p,q}g(\psi)| + m \left[\int_0^{p\mu} |q\lambda - (\gamma - \gamma p\mu)| {}_{\phi}d_{p,q}\lambda \right. \end{aligned}$$

$$\begin{aligned}
 & + \int_0^1 |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda - \int_0^{p\mu} |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda \\
 & - \int_0^{p\mu} \lambda^\alpha |q\lambda - (y - \gamma p\mu)|_0 d_{p,q}\lambda - \int_0^1 \lambda^\alpha |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda \\
 & + \int_0^{p\mu} \lambda^\alpha |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda \left| \left| {}_\phi D_{p,q} \mathcal{G} \left(\frac{\phi}{m} \right) \right| \right\} \\
 \leq & (\psi - \phi) \left\{ \left[\int_0^{p\mu} (1 - \lambda)^\alpha |q\lambda - (y - \gamma p\mu)|_0 d_{p,q}\lambda + \int_0^1 (1 - \lambda)^\alpha \right. \right. \\
 & \times \left. \left. |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda - \int_0^{p\mu} (1 - \lambda)^\alpha |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda \right] \right. \\
 & \times \left. |{}_\phi D_{p,q} \mathcal{G}(\phi)| + m \left[\int_0^{p\mu} |q\lambda - (y - \gamma p\mu)|_0 d_{p,q}\lambda + \int_0^1 |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda \right. \right. \\
 & - \left. \left. \int_0^{p\mu} |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda - \int_0^{p\mu} (1 - \lambda)^\alpha |q\lambda - (y - \gamma p\mu)|_0 d_{p,q}\lambda - \int_0^1 (1 - \lambda)^\alpha \right. \right. \\
 & \times \left. \left. |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda + \int_0^{p\mu} (1 - \lambda)^\alpha |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda \right] \left| \left| {}_\phi D_{p,q} \mathcal{G} \left(\frac{\psi}{m} \right) \right| \right\}.
 \end{aligned}$$

Using Lemmas 2.3–2.8, we get the desired result. □

Corollary 3.1. *Let $\mu = 1/[2]_{p,q}$. Then Theorem 3.1 leads to*

$$\begin{aligned}
 & \left| y \frac{q\mathcal{G}(\phi) + p\mathcal{G}(\psi)}{[2]_{p,q}} + (1 - \gamma) \mathcal{G} \left(\frac{q\phi + p\psi}{[2]_{p,q}} \right) - \frac{1}{p(\psi - \phi)} \int_\phi^{p\psi + (1-p)\phi} \mathcal{G}(x)_\phi d_{p,q}x \right| \\
 & \leq \min \left[H_1 \left(\gamma, \frac{p}{[2]_{p,q}}, \alpha, m \right), H_2 \left(\gamma, \frac{p}{[2]_{p,q}}, \alpha, m \right) \right].
 \end{aligned}$$

Remark 3.1.

(1) Let $\gamma = 0$. Then Corollary 3.1 gives the midpoint-type integral inequality

$$\left| \mathcal{G} \left(\frac{q\phi + p\psi}{[2]_{p,q}} \right) - \frac{1}{p(\psi - \phi)} \int_\phi^{p\psi + (1-p)\phi} \mathcal{G}(x)_\phi d_{p,q}x \right| \leq \min \left[H_1 \left(0, \frac{p}{[2]_{p,q}}, \alpha, m \right), H_2 \left(0, \frac{p}{[2]_{p,q}}, \alpha, m \right) \right],$$

where

$$\begin{aligned}
 H_1 \left(0, \frac{p}{[2]_{p,q}}, \alpha, m \right) & = (\phi - \psi) \left[\frac{[([2]_{p,q})^{\alpha+2} - (p^{\alpha+2} + q^{\alpha+2})](p - q)^2}{([2]_{p,q})^{\alpha+2}(p^{\alpha+1} - q^{\alpha+1})(p^{\alpha+2} - q^{\alpha+2})} |{}_\phi D_{p,q} \mathcal{G}(\psi)| \right. \\
 & \left. + m \left[\frac{2qp^2}{([2]_{p,q})^3} - \frac{[([2]_{p,q})^{\alpha+2} - (p^{\alpha+2} + q^{\alpha+2})](p - q)^2}{([2]_{p,q})^{\alpha+2}(p^{\alpha+1} - q^{\alpha+1})(p^{\alpha+2} - q^{\alpha+2})} \right] \left| \left| {}_\phi D_{p,q} \mathcal{G} \left(\frac{\phi}{m} \right) \right| \right],
 \end{aligned}$$

$$H_2\left(0, \frac{p}{[2]_{p,q}}, \alpha, m\right) = (\psi - \phi) \left\{ \left[v_2\left(0, \frac{p}{[2]_{p,q}}, \alpha\right) + v_4\left(0, \frac{p}{[2]_{p,q}}, \alpha\right) - v_6\left(0, \frac{p}{[2]_{p,q}}, \alpha\right) \right] |{}_{\phi}D_{p,q}g(\phi)| \right. \\ \left. + m \left[\frac{2qp^2}{([2]_{p,q})^3} - v_2\left(0, \frac{p}{[2]_{p,q}}, \alpha\right) - v_4\left(0, \frac{p}{[2]_{p,q}}, \alpha\right) + v_6\left(0, \frac{p}{[2]_{p,q}}, \alpha\right) \right] \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right| \right\}.$$

In particular, if $\alpha = 1 = m$, then we obtain

$$\left| g\left(\frac{q\phi + p\psi}{[2]_{p,q}}\right) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x) {}_{\phi}d_{p,q}x \right| \tag{3.1} \\ \leq (\psi - \phi)q \left[\frac{3p^3}{([2]_{p,q})^3(p^2 + pq + q^2)} |{}_{\phi}D_{p,q}g(\psi)| + \frac{2p^4 + 2p^2q^2 + 2p^3q - 3p^3}{([2]_{p,q})^3(p^2 + pq + q^2)} |{}_{\phi}D_{p,q}g(\phi)| \right],$$

which was proposed by Kunt et al. in [61].

Let $p = 1$. Then equation (3.1) becomes Theorem 13 of [65].

(2) Taking $\gamma = 1/3$ and $\alpha = 1 = m$ in Corollary 3.1, we get the Simpson-type integral inequality

$$\left| \frac{1}{3} \left[\frac{qg(\phi) + pg(\psi)}{[2]_{p,q}} + 2g\left(\frac{q\phi + p\psi}{[2]_{p,q}}\right) \right] - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x) {}_{\phi}d_{p,q}x \right| \\ \leq \min \left[H_1\left(\frac{1}{3}, \frac{p}{[2]_{p,q}}, 1, 1\right), H_2\left(\frac{1}{3}, \frac{p}{[2]_{p,q}}, 1, 1\right) \right].$$

If $q \rightarrow 1^-$ and $p = 1$, then we obtain

$$\left| \frac{1}{3} \left[\frac{g(\phi) + g(\psi)}{2} + 2g\left(\frac{\phi + \psi}{2}\right) \right] - \frac{1}{(\psi - \phi)} \int_a^{\psi} g(x) dx \right| \leq \frac{5(\psi - \phi)}{72} [|g'(\psi)| + |g'(\phi)|],$$

which was proposed by Alomari et al. in [67].

(3) Let $\gamma = 1/2$ and $\alpha = 1 = m$. Then we get the average of midpoint and trapezoid-type integral inequality

$$\left| \frac{1}{2} \left[\frac{qg(\phi) + pg(\psi)}{[2]_{p,q}} + g\left(\frac{q\phi + p\psi}{[2]_{p,q}}\right) \right] - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x) {}_{\phi}d_{p,q}x \right| \\ \leq \min \left[H_1\left(\frac{1}{2}, \frac{p}{[2]_{p,q}}, 1, 1\right), H_2\left(\frac{1}{2}, \frac{p}{[2]_{p,q}}, 1, 1\right) \right].$$

If $q \rightarrow 1^-$ and $p = 1$, then we obtain

$$\left| \frac{1}{2} \left[\frac{g(\phi) + g(\psi)}{2} + g\left(\frac{\phi + \psi}{2}\right) \right] - \frac{1}{(\psi - \phi)} \int_{\phi}^{\psi} g(x) dx \right| \leq \frac{\psi - \phi}{16} [|g'(\psi)| + |g'(\phi)|],$$

which was proposed by Xi et al. in [68].

(4) Let $\gamma = 1$. Then we get the trapezoid-type integral inequality

$$\left| \frac{qg(\phi) + pg(\psi)}{[2]_{p,q}} - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x) {}_{\phi}d_{p,q}x \right| \leq \min \left[H_1\left(1, \frac{p}{[2]_{p,q}}, \alpha, m\right), H_2\left(1, \frac{p}{[2]_{p,q}}, \alpha, m\right) \right].$$

In particular, if $\alpha = p = m = 1$, then we obtain

$$\left| \frac{qg(\phi) + g(\psi)}{[2]_q} - \frac{1}{\psi - \phi} \int_{\phi}^{\psi} g(x) {}_{\phi}d_q x \right| \leq (\psi - \phi) q^2 \left[\frac{1 + 4q + q^2}{([2]_q)^4 [3]_q} |{}_{\phi}D_q g(\psi)| + \frac{1 + 3q^2 + 2q^3}{([2]_q)^4 [3]_q} |{}_{\phi}D_q g(\phi)| \right],$$

which was proposed by Sudsutad et al. in [55].

Theorem 3.2. Let $0 \leq \phi < \psi < \infty, r > 1, \alpha, m \in (0, 1], 0 < q < p \leq 1$ and $g : J \supset [0, \infty) \rightarrow \mathbb{R}$ be a (p, q) -differentiable function on J° (the interior of J) such that ${}_{\phi}D_{p,q}g$ is continuous and integrable on $\left[0, \frac{\psi}{m}\right]$ and $|{}_{\phi}D_{p,q}g|^r$ is (α, m) -convex on $\left[0, \frac{\psi}{m}\right]$. Then the inequality

$$\left| \gamma[p\mu g(\psi) + (1 - p\mu)g(\phi)] + (1 - \gamma)g(p\mu\psi + (1 - p\mu)\phi) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x) {}_{\phi}d_{p,q} x \right| \leq (\psi - \phi) \min[T_1(\gamma, p\mu, \alpha, m, r), T_2(\gamma, p\mu, \alpha, m, r)]$$

holds for all $\gamma, \mu \in [0, 1]$, where

$$T_1(\gamma, p\mu, \alpha, m, r) = v_8^{1-\frac{1}{r}}(\gamma, p\mu) \left[v_3(\gamma, p\mu, \alpha) |{}_{\phi}D_{p,q}g(\psi)|^r + m(v_8(\gamma, p\mu) - v_3(\gamma, p\mu, \alpha)) \left| {}_{\phi}D_{p,q}g\left(\frac{\phi}{m}\right) \right|^r \right]^{\frac{1}{r}} + (1 - \gamma)(p\mu)^{1-\frac{1}{r}} \left[\Gamma_1(p\mu, \alpha) |{}_{\phi}D_{p,q}g(\psi)|^r + m\Gamma_2(p\mu, \alpha) \left| {}_{\phi}D_{p,q}g\left(\frac{\phi}{m}\right) \right|^r \right]^{\frac{1}{r}},$$

$$T_2(\gamma, p\mu, \alpha, m, r) = v_8^{1-\frac{1}{r}}(\gamma, p\mu) [v_4(\gamma, p\mu, \alpha) |{}_{\phi}D_{p,q}g(\phi)|^r + m(v_8(\gamma, p\mu) - v_4(\gamma, p\mu, \alpha)) \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r]^{\frac{1}{r}} + (p\mu)^{1-\frac{1}{r}}(1 - \gamma) \left[\Gamma_3(p\mu, \alpha) |{}_{\phi}D_{p,q}g(\phi)|^r + m\Gamma_4(p\mu, \alpha) \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r \right]^{\frac{1}{r}}$$

and

$$\Gamma_1(p\mu, \alpha) = \int_0^{p\mu} \lambda^\alpha {}_0d_{p,q}\lambda = \frac{\mu^{\alpha+1}(p - q)p^{\alpha+1}}{p^{\alpha+1} - q^{\alpha+1}}, \tag{3.2}$$

$$\Gamma_2(p\mu, \alpha) = \int_0^{p\mu} (1 - \lambda^\alpha) {}_0d_{p,q}\lambda = p\mu - \frac{\mu^{\alpha+1}(p - q)p^{\alpha+1}}{p^{\alpha+1} - q^{\alpha+1}}, \tag{3.3}$$

$$\Gamma_3(p\mu, \alpha) = \int_0^{p\mu} (1 - \lambda)^\alpha {}_0d_{p,q}\lambda = (p - q)\mu \sum_{n=0}^{\infty} \frac{q^n}{p^n} \left(1 - \frac{q^n}{p^n}\mu\right)^\alpha, \tag{3.4}$$

$$\Gamma_4(p\mu, \alpha) = \int_0^{p\mu} (1 - (1 - \lambda)^\alpha) {}_0d_{p,q}\lambda = p\mu - (p - q)\mu \sum_{n=0}^{\infty} \frac{q^n}{p^n} \left(1 - \frac{q^n}{p^n}\mu\right)^\alpha \tag{3.5}$$

and $v_3(\gamma, p\mu, \alpha), v_4(\gamma, p\mu, \alpha)$ and $v_8(\gamma, p\mu)$ are defined in Lemmas 2.4 and 2.7, respectively.

Proof. Using Lemma 2.1 and the power mean inequality, we have

$$\begin{aligned} & \left| \gamma[p\mu g(\psi) + (1 - p\mu)g(\phi)] + (1 - \gamma)g(p\mu\psi + (1 - p\mu)\phi) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi+(1-p)\phi} g(x)_{\phi} d_{p,q}x \right| \\ & \leq (\psi - \phi) \left\{ \left[\int_0^1 |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda \right]^{1-\frac{1}{r}} \times \left[\int_0^1 |q\lambda - (1 - \gamma p\mu)|_{\phi} D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)^r {}_0 d_{p,q}\lambda \right]^{\frac{1}{r}} \right. \\ & \quad \left. + (1 - \gamma) \left[\int_0^{p\mu} 1_0 d_{p,q}\lambda \right]^{1-\frac{1}{r}} \left[\int_0^{p\mu} |_{\phi} D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)^r {}_0 d_{p,q}\lambda \right]^{\frac{1}{r}} \right\}. \end{aligned} \tag{3.6}$$

Utilizing the (α, m) -convexity of $|_{\phi} D_{p,q}g|^r$, we get

$$\begin{aligned} & \int_0^1 |q\lambda - (1 - \gamma p\mu)|_{\phi} D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)^r {}_0 d_{p,q}\lambda \\ & \leq \int_0^1 |q\lambda - (1 - \gamma p\mu)| \left[\lambda^{\alpha} |_{\phi} D_{p,q}g(\psi)^r + m(1 - \lambda^{\alpha}) \left| |_{\phi} D_{p,q}g\left(\frac{\phi}{m}\right)^r \right| \right] {}_0 d_{p,q}\lambda \\ & = \left(\int_0^1 \lambda^{\alpha} |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda \right) |_{\phi} D_{p,q}g(\psi)^r \end{aligned} \tag{3.7}$$

$$\begin{aligned} & + m \left(\int_0^1 |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda - \int_0^1 \lambda^{\alpha} |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda \right) \left| |_{\phi} D_{p,q}g\left(\frac{\phi}{m}\right)^r \right|, \\ & \int_0^{p\mu} |_{\phi} D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)^r {}_0 d_{p,q}\lambda \\ & \leq \int_0^{p\mu} \left[\lambda^{\alpha} |_{\phi} D_{p,q}g(\psi)^r + m(1 - \lambda^{\alpha}) \left| |_{\phi} D_{p,q}g\left(\frac{\phi}{m}\right)^r \right| \right] {}_0 d_{p,q}\lambda \\ & = \left(\int_0^{p\mu} \lambda^{\alpha} {}_0 d_{p,q}\lambda \right) |_{\phi} D_{p,q}g(\psi)^r + m \left(\int_0^{p\mu} (1 - \lambda^{\alpha}) {}_0 d_{p,q}\lambda \right) \left| |_{\phi} D_{p,q}g\left(\frac{\phi}{m}\right)^r \right|. \end{aligned} \tag{3.8}$$

Using (3.7) and (3.8) in (3.6), we get

$$\begin{aligned} & \left| \gamma[p\mu g(\psi) + (1 - p\mu)g(\phi)] + (1 - \gamma)g(p\mu\psi + (1 - p\mu)\phi) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi+(1-p)\phi} g(x)_{\phi} d_{p,q}x \right| \\ & \leq (\psi - \phi) \left\{ \left[\int_0^1 |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda \right]^{1-\frac{1}{r}} \left[\left(\int_0^1 \lambda^{\alpha} |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda \right) |_{\phi} D_{p,q}g(\psi)^r \right. \right. \\ & \quad \left. \left. + m \left(\int_0^1 |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda - \int_0^1 \lambda^{\alpha} |q\lambda - (1 - \gamma p\mu)|_0 d_{p,q}\lambda \right) \left| |_{\phi} D_{p,q}g\left(\frac{\phi}{m}\right)^r \right| \right]^{\frac{1}{r}} \right. \\ & \quad \left. + (1 - \gamma)(p\mu)^{1-\frac{1}{r}} \left[\left(\int_0^{p\mu} \lambda^{\alpha} {}_0 d_{p,q}\lambda \right) |_{\phi} D_{p,q}g(\psi)^r + m \left(\int_0^{p\mu} (1 - \lambda^{\alpha}) {}_0 d_{p,q}\lambda \right) \left| |_{\phi} D_{p,q}g\left(\frac{\phi}{m}\right)^r \right| \right]^{\frac{1}{r}} \right\}. \end{aligned} \tag{3.9}$$

Similarly, we get

$$\begin{aligned}
 & \int_0^1 |q\lambda - (1 - \gamma p\mu)| |{}_{\phi}D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)|^r {}_0d_{p,q}\lambda \\
 & \leq \int_0^1 |q\lambda - (1 - \gamma p\mu)| \left[(1 - \lambda)^\alpha |{}_{\phi}D_{p,q}g(\phi)|^r + m(1 - (1 - \lambda)^\alpha) \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r \right] {}_0d_{p,q}\lambda \\
 & = \left(\int_0^1 (1 - \lambda)^\alpha |q\lambda - (1 - \gamma p\mu)| {}_0d_{p,q}\lambda \right) |{}_{\phi}D_{p,q}g(\phi)|^r \\
 & \quad + m \left(\int_0^1 |q\lambda - (1 - \gamma p\mu)| {}_0d_{p,q}\lambda - \int_0^1 (1 - \lambda)^\alpha |q\lambda - (1 - \gamma p\mu)| {}_0d_{p,q}\lambda \right) \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r.
 \end{aligned} \tag{3.10}$$

$$\begin{aligned}
 & \int_0^{p\mu} |{}_{\phi}D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)|^r {}_0d_{p,q}\lambda \\
 & \leq \int_0^{p\mu} \left[(1 - \lambda)^\alpha |{}_{\phi}D_{p,q}g(\phi)|^r + m(1 - (1 - \lambda)^\alpha) \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r \right] {}_0d_{p,q}\lambda \\
 & = \left(\int_0^{p\mu} (1 - \lambda)^\alpha {}_0d_{p,q}\lambda \right) |{}_{\phi}D_{p,q}g(\phi)|^r + m \left(\int_0^{p\mu} (1 - (1 - \lambda)^\alpha) {}_0d_{p,q}\lambda \right) \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r.
 \end{aligned} \tag{3.11}$$

Using (3.10) and (3.11) in (3.6), we get

$$\begin{aligned}
 & \left| \gamma[p\mu g(\psi) + (1 - p\mu)g(\phi)] + (1 - \gamma)g(p\mu b + (1 - p\mu)a) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x) {}_{\phi}d_{p,q}x \right| \\
 & \leq (\psi - \phi) \left\{ \left[\int_0^1 |q\lambda - (1 - \gamma p\mu)| {}_0d_{p,q}\lambda \right]^{1-\frac{1}{r}} \left[\int_0^1 (1 - \lambda)^\alpha |q\lambda - (1 - \gamma p\mu)| {}_0d_{p,q}\lambda \right] |{}_{\phi}D_{p,q}g(\phi)|^r \right. \\
 & \quad + m \left(\int_0^1 |q\lambda - (1 - \gamma p\mu)| {}_0d_{p,q}\lambda - \int_0^1 (1 - \lambda)^\alpha |q\lambda - (1 - \gamma p\mu)| {}_0d_{p,q}\lambda \right) \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r \left. \right\} \\
 & \quad + (1 - \gamma)(p\mu)^{1-\frac{1}{r}} \left\{ \left[\int_0^{p\mu} (1 - \lambda)^\alpha {}_0d_{p,q}\lambda \right] |{}_{\phi}D_{p,q}g(\phi)|^r + m \left(\int_0^{p\mu} (1 - (1 - \lambda)^\alpha) {}_0d_{p,q}\lambda \right) \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r \right\}.
 \end{aligned} \tag{3.12}$$

Therefore, the desired result follows from (3.9) and (3.12) together with Lemmas 2.4 and 2.7. □

Theorem 3.3. Let $0 \leq \phi < \psi < \infty$, $0 < q < p \leq 1$, $r, s > 1$ with $r^{-1} + s^{-1} = 1$, $\alpha, m \in (0, 1]$ and $g : J \supset [0, \infty) \rightarrow \mathbb{R}$ be a (p, q) -differentiable function on J° (the interior of J) such that ${}_{\phi}D_{p,q}g$ is continuous and integrable on $[0, \frac{\psi}{m}]$ and $|{}_{\phi}D_{p,q}g|^r$ is (α, m) -convex on $[0, \frac{\psi}{m}]$. Then the inequality

$$\begin{aligned}
 & \left| \gamma[p\mu g(\psi) + (1 - p\mu)g(\phi)] + (1 - \gamma)g(p\mu\psi + (1 - p\mu)\phi) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x) {}_{\phi}d_{p,q}x \right| \\
 & \leq (\psi - \phi) \min[K_1(\gamma, p\mu, \alpha, m), K_2(\gamma, p\mu, \alpha, m)]
 \end{aligned}$$

holds for all $\gamma, \mu \in [0, 1]$, where

$$\begin{aligned}
 K_1(\gamma, p\mu, \alpha, m) &= \nu_{i_0}^{\frac{1}{s}}(\gamma, p\mu) \left[\eta_2(\alpha) |\phi D_{p,q}g(\psi)|^r + m(1 - \eta_2(\alpha)) \left| \phi D_{p,q}g\left(\frac{\phi}{m}\right) \right|^r \right]^{\frac{1}{r}} \\
 &\quad + (1 - \gamma)(p\mu)^{\frac{1}{s}} \left[\Gamma_1(p\mu, \alpha) |\phi D_{p,q}g(\psi)|^r + m\Gamma_2(p\mu, \alpha) \left| \phi D_{p,q}g\left(\frac{\phi}{m}\right) \right|^r \right]^{\frac{1}{r}}, \\
 K_2(\gamma, p\mu, \alpha, m) &= \nu_{i_0}^{\frac{1}{s}}(\gamma, p\mu) \left[\eta_3(\alpha) |\phi D_{p,q}g(\phi)|^r + m(1 - \eta_3(\alpha)) \left| \phi D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r \right]^{\frac{1}{r}} \\
 &\quad + (1 - \gamma)(p\mu)^{\frac{1}{s}} \left[\Gamma_3(p\mu, \alpha) |\phi D_{p,q}g(\phi)|^r + m\Gamma_4(p\mu, \alpha) \left| \phi D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r \right]^{\frac{1}{r}}, \\
 \eta_2(\alpha) &= \int_0^1 \lambda^\alpha d_{p,q}\lambda = \frac{p - q}{p^{1+\alpha} - q^{1+\alpha}}, \\
 \eta_3(\alpha) &= \int_0^1 (1 - \lambda)^\alpha d_{p,q}\lambda = (p - q) \sum_{n=0}^\infty \frac{q^n}{p^{n+1}} \left(1 - \frac{q^n}{p^{n+1}}\right)^\alpha
 \end{aligned}$$

and $\Gamma_1(p\mu, \alpha)$, $\Gamma_2(p\mu, \alpha)$, $\Gamma_3(p\mu, \alpha)$ and $\Gamma_4(p\mu, \alpha)$ are defined in Theorem 3.2.

Proof. Using Lemma 2.1 and the Hölder inequality, we have

$$\begin{aligned}
 &\left| \gamma[p\mu g(\psi) + (1 - p\mu)g(\phi)] + (1 - \gamma)g(p\mu\psi + (1 - p\mu)\phi) - \frac{1}{p(\psi - \phi)} \int_\phi^{p\psi+(1-p)\phi} g(x) \phi d_{p,q}x \right| \\
 &\leq (\psi - \phi) \left\{ \left(\int_0^1 |q\lambda - (1 - \gamma p\mu)|^s d_{p,q}\lambda \right)^{\frac{1}{s}} \left(\int_0^1 |\phi D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)|^r d_{p,q}\lambda \right)^{\frac{1}{r}} \right. \\
 &\quad \left. + (1 - \gamma) \left(\int_0^{p\mu} 1^s d_{p,q}\lambda \right)^{\frac{1}{s}} \left(\int_0^{p\mu} |\phi D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)|^r d_{p,q}\lambda \right)^{\frac{1}{r}} \right\}.
 \end{aligned} \tag{3.13}$$

Utilizing the (α, m) -convexity of $|\phi D_{p,q}g|^r$, we get

$$\begin{aligned}
 \int_0^1 |\phi D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)|^r d_{p,q}\lambda &\leq \int_0^1 \left[\lambda^\alpha |\phi D_{p,q}g(\psi)|^r + m(1 - \lambda^\alpha) \left| \phi D_{p,q}g\left(\frac{\phi}{m}\right) \right|^r \right] d_{p,q}\lambda \\
 &= \left(\int_0^1 \lambda^\alpha d_{p,q}\lambda \right) |\phi D_{p,q}g(\psi)|^r + m \left(\int_0^1 (1 - \lambda^\alpha) d_{p,q}\lambda \right) \left| \phi D_{p,q}g\left(\frac{\phi}{m}\right) \right|^r
 \end{aligned} \tag{3.14}$$

and

$$\begin{aligned}
 \int_0^{p\mu} |\phi D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)|^r d_{p,q}\lambda &\leq \int_0^{p\mu} \left[\lambda^\alpha |\phi D_{p,q}g(\psi)|^r + m(1 - \lambda^\alpha) \left| \phi D_{p,q}g\left(\frac{\phi}{m}\right) \right|^r \right] d_{p,q}\lambda \\
 &= \left(\int_0^{p\mu} \lambda^\alpha d_{p,q}\lambda \right) |\phi D_{p,q}g(\psi)|^r + m \left(\int_0^{p\mu} (1 - \lambda^\alpha) d_{p,q}\lambda \right) \left| \phi D_{p,q}g\left(\frac{\phi}{m}\right) \right|^r.
 \end{aligned} \tag{3.15}$$

Using (3.14) and (3.15) in (3.13), we get

$$\begin{aligned}
 & \left| \gamma[p\mu g(\psi) + (1 - p\mu)g(\phi)] + (1 - \gamma)g(p\mu\psi + (1 - p\mu)\phi) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x)_{\phi} d_{p,q}x \right| \\
 & \leq (\psi - \phi) \left\{ \left[\int_0^1 |q\lambda - (1 - \gamma p\mu)|^s {}_0d_{p,q}\lambda \right]^{\frac{1}{s}} \left[\int_0^1 \lambda^{\alpha} {}_0d_{p,q}\lambda \right] |{}_{\phi}D_{p,q}g(\psi)|^r + m \left(\int_0^1 (1 - \lambda^{\alpha}) {}_0d_{p,q}\lambda \right) \left| {}_{\phi}D_{p,q}g\left(\frac{\phi}{m}\right) \right|^r \right]^{\frac{1}{r}} \\
 & + (1 - \gamma)(p\mu)^{\frac{1}{s}} \left\{ \left[\int_0^{p\mu} \lambda^{\alpha} {}_0d_{p,q}\lambda \right] |{}_{\phi}D_{p,q}g(\psi)|^r + m \left(\int_0^{p\mu} (1 - \lambda^{\alpha}) {}_0d_{p,q}\lambda \right) \left| {}_{\phi}D_{p,q}g\left(\frac{\phi}{m}\right) \right|^r \right\}^{\frac{1}{r}}.
 \end{aligned} \tag{3.16}$$

Similarly, we get

$$\begin{aligned}
 & \int_0^1 |\phi D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)|^r {}_0d_{p,q}\lambda \\
 & \leq \int_0^1 \left[(1 - \lambda)^{\alpha} |{}_{\phi}D_{p,q}g(\phi)|^r + m(1 - (1 - \lambda)^{\alpha}) \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r \right] {}_0d_{p,q}\lambda \\
 & = \left(\int_0^1 (1 - \lambda)^{\alpha} {}_0d_{p,q}\lambda \right) |{}_{\phi}D_{p,q}g(\phi)|^r + m \left(\int_0^1 (1 - (1 - \lambda)^{\alpha}) {}_0d_{p,q}\lambda \right) \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r
 \end{aligned} \tag{3.17}$$

and

$$\begin{aligned}
 & \int_0^{p\mu} |\phi D_{p,q}g(\lambda\psi + (1 - \lambda)\phi)|^r {}_0d_{p,q}\lambda \\
 & \leq \int_0^{p\mu} \left[(1 - \lambda)^{\alpha} |{}_{\phi}D_{p,q}g(\phi)|^r + m(1 - (1 - \lambda)^{\alpha}) \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r \right] {}_0d_{p,q}\lambda \\
 & = \left(\int_0^{p\mu} (1 - \lambda)^{\alpha} {}_0d_{p,q}\lambda \right) |{}_{\phi}D_{p,q}g(\phi)|^r + m \left(\int_0^{p\mu} (1 - (1 - \lambda)^{\alpha}) {}_0d_{p,q}\lambda \right) \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r.
 \end{aligned} \tag{3.18}$$

Using (3.17) and (3.18) in (3.13), we get

$$\begin{aligned}
 & \left| \gamma[\mu f(\psi) + (1 - \mu)g(\phi)] + (1 - \gamma)g(p\mu\psi + (1 - p\mu)\phi) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x)_{\phi} d_{p,q}x \right| \\
 & \leq (\psi - \phi) \left\{ \left[\int_0^1 |q\lambda - (1 - \gamma p\mu)|^s {}_0d_{p,q}\lambda \right]^{\frac{1}{s}} \left[\int_0^1 (1 - \lambda)^{\alpha} {}_0d_{p,q}\lambda \right] |{}_{\phi}D_{p,q}g(\phi)|^r \right. \\
 & + m \left(\int_0^1 (1 - (1 - \lambda)^{\alpha}) {}_0d_{p,q}\lambda \right) \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r \right]^{\frac{1}{r}} \\
 & + (1 - \gamma)(p\mu)^{\frac{1}{s}} \left\{ \left[\int_0^{p\mu} (1 - \lambda)^{\alpha} {}_0d_{p,q}\lambda \right] |{}_{\phi}D_{p,q}g(\phi)|^r + m \left(\int_0^{p\mu} (1 - (1 - \lambda)^{\alpha}) {}_0d_{p,q}\lambda \right) \left| {}_{\phi}D_{p,q}g\left(\frac{\psi}{m}\right) \right|^r \right\}^{\frac{1}{r}}.
 \end{aligned} \tag{3.19}$$

Therefore, Theorem 3.3 follows from (3.2)–(3.5), (3.16) and (3.19) together with Lemmas 2.2 and 2.9. \square

Remark 3.2. If we put $\gamma = 0, \frac{1}{3}, \frac{1}{2}, 1$ and $\mu = \frac{1}{[2]_{p,q}}$ in Theorems 3.2 and 3.3, then we can get the midpoint-type integral inequality, the Simpson-type integral inequality, average of midpoint and trapezoid-type integral inequality and the trapezoid-type integral inequality, respectively.

Next, we establish the (p, q) -integral inequalities involving the product of two (α, m) -convex functions.

Theorem 3.4. *Let $0 \leq \phi < \psi < \infty, 0 < q < p \leq 1, \alpha_1, \alpha_2, m \in (0, 1]$ and $f, g : J \supset [0, \infty) \rightarrow \mathbb{R}$ be continuous and integrable on $\left[0, \frac{\psi}{m}\right]$ such that f and g are (α_1, m) -convex and (α_2, m) -convex on $\left[0, \frac{\psi}{m}\right]$, respectively. Then the inequality*

$$\frac{1}{\psi - \phi} \int_{\phi}^{\psi} f(x)g(x) {}_{\phi}d_{p,q}x \leq \min\{L_1(\alpha_1, \alpha_2, m), L_2(\alpha_1, \alpha_2, m)\}$$

holds, where

$$\begin{aligned} L_1(\alpha_1, \alpha_2, m) &= m^2 \left[\frac{p - q}{p^{\alpha_1 + \alpha_2 + 1} - q^{\alpha_1 + \alpha_2 + 1}} - \frac{p - q}{p^{\alpha_1 + 1} - q^{\alpha_1 + 1}} - \frac{p - q}{p^{\alpha_2 + 1} - q^{\alpha_2 + 1}} + 1 \right] f\left(\frac{\phi}{m}\right)g\left(\frac{\phi}{m}\right) \\ &+ \frac{p - q}{p^{\alpha_1 + \alpha_2 + 1} - q^{\alpha_1 + \alpha_2 + 1}} f(\psi)g(\psi) + m \left[\frac{p - q}{p^{\alpha_2 + 1} - q^{\alpha_2 + 1}} - \frac{p - q}{p^{\alpha_1 + \alpha_2 + 1} - q^{\alpha_1 + \alpha_2 + 1}} \right] f\left(\frac{\phi}{m}\right)g(\psi) \\ &+ m \left[\frac{p - q}{p^{\alpha_1 + 1} - q^{\alpha_1 + 1}} - \frac{p - q}{p^{\alpha_1 + \alpha_2 + 1} - q^{\alpha_1 + \alpha_2 + 1}} \right] f(\psi)g\left(\frac{\phi}{m}\right), \end{aligned}$$

$$\begin{aligned} L_2(\alpha_1, \alpha_2, m) &= m^2 [\Lambda(\alpha_1, \alpha_2) - \Lambda(\alpha_1) - \Lambda(\alpha_2) + 1] f\left(\frac{\psi}{m}\right)g\left(\frac{\psi}{m}\right) \\ &+ \Lambda(\alpha_1, \alpha_2) f(\phi)g(\phi) + m [\Lambda(\alpha_1) - \Lambda(\alpha_1, \alpha_2)] f(\phi)g\left(\frac{\psi}{m}\right) \\ &+ m [\Lambda(\alpha_2) - \Lambda(\alpha_1, \alpha_2)] f\left(\frac{\psi}{m}\right)g(\phi), \end{aligned}$$

$$\Lambda(\alpha_1, \alpha_2) = \int_0^1 (1 - \lambda)^{\alpha_1 + \alpha_2} {}_0d_{p,q}\lambda = (p - q) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} \left(1 - \frac{q^n}{p^{n+1}}\right)^{\alpha_1 + \alpha_2}$$

and

$$\Lambda(\alpha_i) = \int_0^1 (1 - \lambda)^{\alpha_i} {}_0d_{p,q}\lambda = (p - q) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} \left(1 - \frac{q^n}{p^{n+1}}\right)^{\alpha_i} \quad (i = 1, 2).$$

Proof. Let $\lambda \in [0, 1]$. Then it follows from the (α_1, m) -convexity of f and the (α_2, m) -convexity of g that

$$f(\lambda\psi + (1 - \lambda)\phi) \leq \lambda^{\alpha_1} f(\psi) + m(1 - \lambda^{\alpha_1}) f\left(\frac{\phi}{m}\right) \tag{3.20}$$

and

$$g(\lambda\psi + (1 - \lambda)\phi) \leq \lambda^{\alpha_2} g(\psi) + m(1 - \lambda^{\alpha_2}) g\left(\frac{\phi}{m}\right). \tag{3.21}$$

Multiplying (3.20) with (3.21), we get

$$\begin{aligned} f(\lambda\psi + (1 - \lambda)\phi)g(\lambda\psi + (1 - \lambda)\phi) &\leq \lambda^{\alpha_1 + \alpha_2} f(\psi)g(\psi) + m^2(1 - \lambda^{\alpha_1})(1 - \lambda^{\alpha_2}) f\left(\frac{\phi}{m}\right)g\left(\frac{\phi}{m}\right) \\ &+ m\lambda^{\alpha_2}(1 - \lambda^{\alpha_1}) f\left(\frac{\phi}{m}\right)g(\psi) + m\lambda^{\alpha_1}(1 - \lambda^{\alpha_2}) f(\psi)g\left(\frac{\phi}{m}\right). \end{aligned} \tag{3.22}$$

Taking the (p, q) -integral for (3.22) with respect to λ on $(0, 1)$ and by using Lemma 2.2, we get

$$\begin{aligned} & \int_0^1 f(\lambda\psi + (1 - \lambda)\phi)g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q}\lambda \\ & \leq m^2 \left[\frac{p - q}{p^{\alpha_1 + \alpha_2 + 1} - q^{\alpha_1 + \alpha_2 + 1}} - \frac{p - q}{p^{\alpha_1 + 1} - q^{\alpha_1 + 1}} - \frac{p - q}{p^{\alpha_2 + 1} - q^{\alpha_2 + 1}} + 1 \right] f\left(\frac{\phi}{m}\right)g\left(\frac{\phi}{m}\right) + \frac{p - q}{p^{\alpha_1 + \alpha_2 + 1} - q^{\alpha_1 + \alpha_2 + 1}} f(\psi)g(\psi) \quad (3.23) \\ & + m \left[\frac{p - q}{p^{\alpha_2 + 1} - q^{\alpha_2 + 1}} - \frac{p - q}{p^{\alpha_1 + \alpha_2 + 1} - q^{\alpha_1 + \alpha_2 + 1}} \right] f\left(\frac{\phi}{m}\right)g(\psi) + m \left[\frac{p - q}{p^{\alpha_1 + 1} - q^{\alpha_1 + 1}} - \frac{p - q}{p^{\alpha_1 + \alpha_2 + 1} - q^{\alpha_1 + \alpha_2 + 1}} \right] f(\psi)g\left(\frac{\phi}{m}\right). \end{aligned}$$

Similarly, we have

$$\begin{aligned} & \int_0^1 f(\lambda\psi + (1 - \lambda)\phi)g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q}\lambda \\ & \leq m^2 \left(\int_0^1 (1 - \lambda)^{\alpha_1 + \alpha_2} d_{p,q}\lambda - \int_0^1 (1 - \lambda)^{\alpha_1} d_{p,q}\lambda - \int_0^1 (1 - \lambda)^{\alpha_2} d_{p,q}\lambda + 1 \right) f\left(\frac{\psi}{m}\right)g\left(\frac{\psi}{m}\right) \quad (3.24) \\ & + \left(\int_0^1 (1 - \lambda)^{\alpha_1 + \alpha_2} d_{p,q}\lambda \right) f(\phi)g(\phi) + m \left(\int_0^1 (1 - \lambda)^{\alpha_1} d_{p,q}\lambda - \int_0^1 (1 - \lambda)^{\alpha_1 + \alpha_2} d_{p,q}\lambda \right) f(\phi)g\left(\frac{\psi}{m}\right) \\ & + m \left(\int_0^1 (1 - \lambda)^{\alpha_2} d_{p,q}\lambda - \int_0^1 (1 - \lambda)^{\alpha_1 + \alpha_2} d_{p,q}\lambda \right) f\left(\frac{\psi}{m}\right)g(\phi). \end{aligned}$$

Some simple calculations lead to

$$\int_0^1 f(\lambda\psi + (1 - \lambda)\phi)g(\lambda\psi + (1 - \lambda)\phi)_0 d_{p,q}\lambda = \frac{1}{\psi - \phi} \int_{\phi}^{\psi} f(x)g(x)_{\phi} d_{p,q}x. \quad (3.25)$$

Therefore, the desired result follows easily from (3.23) to (3.25). □

Corollary 3.2. *If we choose $\alpha_1 = \alpha_2 = \alpha$ in Theorem 3.4, then we obtain*

$$\frac{1}{\psi - \phi} \int_{\phi}^{\psi} f(x)g(x)_{\phi} d_{p,q}x \leq \min\{L_1(\alpha, m), L_2(\alpha, m)\},$$

where

$$\begin{aligned} L_1(\alpha, m) &= m^2 \left[\frac{p - q}{p^{2\alpha + 1} - q^{2\alpha + 1}} - \frac{2(p - q)}{p^{\alpha + 1} - q^{\alpha + 1}} + 1 \right] f\left(\frac{\phi}{m}\right)g\left(\frac{\phi}{m}\right) \\ &+ \frac{p - q}{p^{2\alpha + 1} - q^{2\alpha + 1}} f(\psi)g(\psi) + m \left[\frac{q^{\alpha + 1}(p - q)(p^{\alpha} - q^{\alpha})}{(p^{\alpha + 1} - q^{\alpha + 1})(p^{2\alpha + 1} - q^{2\alpha + 1})} \right] \left[f\left(\frac{\phi}{m}\right)g(\psi) + f(\psi)g\left(\frac{\phi}{m}\right) \right] \end{aligned}$$

and

$$\begin{aligned} L_2(\alpha, m) &= m^2 \left[(p - q) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} \left(1 - \frac{q^n}{p^{n+1}}\right)^{2\alpha} - 2(p - q) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} \left(1 - \frac{q^n}{p^{n+1}}\right)^{\alpha} + 1 \right] f\left(\frac{\psi}{m}\right)g\left(\frac{\psi}{m}\right) \\ &+ (p - q) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} \left(1 - \frac{q^n}{p^{n+1}}\right)^{2\alpha} f(\phi)g(\phi) + m \left[(p - q) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} \left(1 - \frac{q^n}{p^{n+1}}\right)^{\alpha} \right. \\ &\left. - (p - q) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} \left(1 - \frac{q^n}{p^{n+1}}\right)^{2\alpha} \right] \left[f(\phi)g\left(\frac{\psi}{m}\right) + f\left(\frac{\psi}{m}\right)g(\phi) \right]. \end{aligned}$$

In particular, the special case of $p = \alpha = m = 1$ for Corollary 3.2 was proved by Sudsutad et al. in [55].

4 Examples

Example 4.1. Let $g : J \supset [0, \infty) \rightarrow \mathbb{R}$ be defined by $g(x) = 4x + 1$. Then it is $(1, \frac{1}{3})$ -differentiable function on J° (the interior of J) and ${}_1D_{1, \frac{1}{3}}g$ is continuous and integrable on $[0, 10]$, $0 \leq 1 < 5 < \infty$ and $0 < \frac{1}{3} < 1 \leq 1$. If $|\phi D_{1, \frac{1}{3}}g|$ is $(1, \frac{1}{2})$ -convex on $[0, 10]$ with $\gamma = 0$ and $\mu = \frac{3}{4}$, then all the assumptions of Theorem 3.1 are satisfied.

We clearly see that

$$\begin{aligned} & \left| \gamma[p\mu g(\psi) + (1 - p\mu)g(\phi)] + (1 - \gamma)g(p\mu\psi + (1 - p\mu)\phi) - \frac{1}{p(\psi - \phi)} \int_{\phi}^{p\psi + (1-p)\phi} g(x)_{\phi}d_{p,q}x \right| \\ &= \left| g(4) - \frac{1}{4} \int_1^5 (4x + 1)_{1}d_{1, \frac{1}{3}}dx \right| = 17 - \frac{68}{4} = 0, \end{aligned} \tag{4.1}$$

where

$$\int_1^5 (4x + 1)_{1}d_{1, \frac{1}{3}}dx = 68.$$

On the other hand,

$$\begin{aligned} v_1(\gamma, p\mu, \alpha) &= v_1\left(0, \frac{3}{4}, \frac{1}{2}\right) \approx 0.1157, & (\gamma + q)p\mu > \gamma, \\ v_2(\gamma, p\mu, \alpha) &= v_2\left(0, \frac{3}{4}, \frac{1}{2}\right) \approx 0.0625, & (\gamma + q)p\mu > \gamma, \\ v_3(\gamma, p\mu, \alpha) &= v_3\left(0, \frac{3}{4}, \frac{1}{2}\right) \approx 0.5881, & \gamma p\mu + q \leq 1, \\ v_4(\gamma, p\mu, \alpha) &= v_4\left(0, \frac{3}{4}, \frac{1}{2}\right) = 0, & \gamma p\mu + q \leq 1, \\ v_5(\gamma, p\mu, \alpha) &= v_5\left(0, \frac{3}{4}, \frac{1}{2}\right) \approx 0.4205, & (\gamma + q)p\mu \leq 1, \\ v_6(\gamma, p\mu, \alpha) &= v_6\left(0, \frac{3}{4}, \frac{1}{2}\right) \approx 0.1875, & (\gamma + q)p\mu \leq 1, \\ v_7(\gamma, p\mu) &= v_7\left(0, \frac{3}{4}\right) \approx 0.1406, & (\gamma + q)p\mu \geq \gamma, \\ v_8(\gamma, p\mu) &= v_8\left(0, \frac{3}{4}\right) \approx 0.75, & \gamma p\mu + q \leq 1, \\ v_9(\gamma, p\mu) &= v_9\left(0, \frac{3}{4}\right) \approx 0.6094, & (\gamma + q)p\mu \leq 1, \\ v_{10}(\gamma, p\mu) &= v_{10}\left(0, \frac{3}{4}\right) \approx 0.2963, & 0 \leq \gamma p\mu \leq 1 - q. \end{aligned} \tag{4.2}$$

Also, we have

$$\begin{aligned} |\phi D_{p,q}g(\psi)| &= |{}_0D_{1, \frac{1}{3}}(4\psi + 1)| = 4, \\ \left| \phi D_{p,q}g\left(\frac{\phi}{m}\right) \right| &= |{}_0D_{1, \frac{1}{3}}(8\phi + 1)| = 3, \\ |\phi D_{p,q}g(\phi)| &= |{}_0D_{1, \frac{1}{3}}(4\phi + 1)| = 0, \\ \left| \phi D_{p,q}g\left(\frac{\psi}{m}\right) \right| &= |{}_0D_{1, \frac{1}{3}}(8\psi + 1)| = 8. \end{aligned} \tag{4.3}$$

Observe that

$$\begin{aligned}
 H_1(\gamma, p\mu, \alpha, m) &= (\psi - \phi) \left\{ [v_1(\gamma, p\mu, \alpha) + v_3(\gamma, p\mu, \alpha) - v_5(\gamma, p\mu, \alpha)] |_{\phi} D_{p,q} g(\psi) \right. \\
 &\quad + m [v_7(\gamma, p\mu) + v_8(\gamma, p\mu) - v_9(\gamma, p\mu) - v_1(\gamma, p\mu, \alpha) \\
 &\quad \left. - v_3(\gamma, p\mu, \alpha) + v_5(\gamma, p\mu, \alpha)] \left|_{\phi} D_{p,q} g\left(\frac{\phi}{m}\right) \right. \right\}. \tag{4.4}
 \end{aligned}$$

Substituting (4.2) and (4.3) in (4.4), and simple computations yield

$$H_1\left(0, \frac{3}{4}, \frac{1}{2}, \frac{1}{2}\right) \approx 4.3167. \tag{4.5}$$

Analogously, we have

$$\begin{aligned}
 H_2(\gamma, p\mu, \alpha, m) &= (\psi - \phi) \left\{ [v_2(\gamma, p\mu, \alpha) + v_4(\gamma, p\mu, \alpha) - v_6(\gamma, p\mu, \alpha)] |_{\phi} D_{p,q} g(\phi) \right. \\
 &\quad + m [v_7(\gamma, p\mu) + v_8(\gamma, p\mu) - v_9(\gamma, p\mu) - v_2(\gamma, p\mu, \alpha) \\
 &\quad \left. - v_4(\gamma, p\mu, \alpha) + v_6(\gamma, p\mu, \alpha)] \left|_{\phi} D_{p,q} g\left(\frac{\psi}{m}\right) \right. \right\}.
 \end{aligned}$$

After simplification, we have

$$H_2\left(0, \frac{3}{4}, \frac{1}{2}, \frac{1}{2}\right) \approx 6.5. \tag{4.6}$$

From (4.5) and (4.6), we get

$$\min[H_1(\gamma, p\mu, \alpha, m), H_2(\gamma, p\mu, \alpha, m)] = \min\{4.3167, 6.5\} \approx 4.3167, \tag{4.7}$$

which shows that the following implications hold in (4.1) and (4.7)

$$0 < 4.3167.$$

Example 4.2. Let $f, g : J \supset [0, \infty) \rightarrow \mathbb{R}$ be defined by $f(x) = g(x) = x$. Then these functions are (p, q) -differentiable functions on J° (the interior of J) and continuous and integrable on $[0, 10]$ with $0 \leq 1 < 5 < \infty$. If f and g are $(1, \frac{1}{2})$ -convex on $[0, 10]$ with $\alpha = 1$ and $m = \frac{1}{2}$, then all assumptions of Corollary 3.2 are satisfied.

Clearly,

$$\frac{1}{\psi - \phi} \int_{\phi}^{\psi} f(x)g(x) |_{\phi} d_{p,q} x = \frac{1}{4} \int_1^5 x^2 |_{p,q} x \tag{4.8}$$

follows from Definition 1.3.

On the other hand,

$$L_1\left(1, \frac{1}{2}\right) = \frac{[2]_{p,q} + (p^2 + pq + q^2)[[2]_{p,q} - 2] + 25([2]_{p,q}) + 10q^2}{[2]_{p,q}(p^2 + pq + q^2)} \tag{4.9}$$

and

$$L_2\left(1, \frac{1}{2}\right) = \frac{25([2]_{p,q} + (p^2 + pq + q^2)[[2]_{p,q} - 2]) + ([2]_{p,q}) + 10q^2}{[2]_{p,q}(p^2 + pq + q^2)}. \tag{4.10}$$

From (4.8) and (4.10), we get

$$\min\left[L_1\left(1, \frac{1}{2}\right), L_2\left(1, \frac{1}{2}\right)\right] = L_1\left(1, \frac{1}{2}\right), \tag{4.11}$$

which shows that the following implications hold in (4.8) and (4.11)

$$\frac{1}{4} \int_1^5 x^2 d_{p,q} x < L_1 \left(1, \frac{1}{2} \right)$$

for every $0 < q < p \leq 1$.

Remark 4.1. Similar technique can be applied to Theorems 3.2 and 3.3 to get the immediate consequences.

Acknowledgments: The authors express their gratitude to the referees for very helpful and detailed comments and suggestions, which have significantly improved the presentation of this paper. The research was supported by the Natural Science Foundation of China under grant numbers 11701176, 61673169, 11301127, 11626101 and 11601485.

Conflict of interest: The authors declare that they have no competing interests.

Author contributions: All authors contributed equally to the manuscript, and they read and approved the final manuscript.

References

- [1] M. Adil Khan, N. Mohammad, E. R. Nwaeze, and Y.-M. Chu, *Quantum Hermite-Hadamard inequality by means of a Green function*, Adv. Differ. Equ. **2020** (2020), 99.
- [2] M. Adil Khan, J. Pečarić and Y.-M. Chu, *Refinements of Jensen's and McShane's inequalities with applications*, AIMS Math. **5** (2020), no. 5, 4931–4945.
- [3] M. U. Awan, N. Akhtar, A. Kashuri, M. A. Noor, and Y.-M. Chu, *2D approximately reciprocal p -convex functions and associated integral inequalities*, AIMS Math. **5** (2020), no. 5, 4662–4680.
- [4] M. A. Latif, M. Kunt, S. S. Dragomir, and İ. İşcan, *Post-quantum trapezoid type inequalities*, AIMS Math. **5** (2020), no. 4, 4011–4026.
- [5] M. U. Awan, S. Talib, Y.-M. Chu, M. A. Noor, and K. I. Noor, *Some new refinements of Hermite-Hadamard-type inequalities involving Ψ_k -Riemann-Liouville fractional integrals and applications*, Math. Probl. Eng. **2020** (2020), 3051920.
- [6] A. Iqbal, M. Adil Khan, S. Ullah, and Y.-M. Chu, *Some new Hermite-Hadamard-type inequalities associated with conformable fractional integrals and their applications*, J. Funct. Spaces **2020** (2020), 9845407.
- [7] Y. Khurshid, M. Adil Khan and Y.-M. Chu, *Conformable fractional integral inequalities for GG- and GA-convex function*, AIMS Math. **5** (2020), no. 5, 5012–5030.
- [8] S. Rashid, R. Ashraf, M. A. Noor, K. I. Noor, and Y.-M. Chu, *New weighted generalizations for differentiable exponentially convex mapping with application*, AIMS Math. **5** (2020), no. 4, 3525–3546.
- [9] S. Rashid, İ. İşcan, D. Baleanu, and Y.-M. Chu, *Generation of new fractional inequalities via n polynomials s -type convexity with applications*, Adv. Differ. Equ. **2020** (2020), 264.
- [10] S. Rashid, F. Jarad and Y.-M. Chu, *A note on reverse Minkowski inequality via generalized proportional fractional integral operator with respect to another function*, Math. Probl. Eng. **2020** (2020), 7630260.
- [11] S. Rashid, F. Jarad, H. Kalsoom, and Y.-M. Chu, *On Pólya-Szegő and Čebyšev type inequalities via generalized k -fractional integrals*, Adv. Differ. Equ. **2020** (2020), 125.
- [12] S.-S. Zhou, S. Rashid, F. Jarad, H. Kalsoom, and Y.-M. Chu, *New estimates considering the generalized proportional Hadamard fractional integral operators*, Adv. Differ. Equ. **2020** (2020), 275.
- [13] T. Abdeljawad, S. Rashid, H. Khan, and Y.-M. Chu, *On new fractional integral inequalities for p -convexity within interval-valued functions*, Adv. Differ. Equ. **2020** (2020), 330.
- [14] S. Hussain, J. Khalid and Y.-M. Chu, *Some generalized fractional integral Simpson's type inequalities with applications*, AIMS Math. **5** (2020), no. 6, 5859–5883.
- [15] L. Xu, Y.-M. Chu, S. Rashid, A. A. El-Deeb, and K. S. Nisar, *On new unified bounds for a family of functions with fractional q -calculus theory*, J. Funct. Spaces **2020** (2020), 4984612.
- [16] S. Rashid, A. Khalid, G. Rahman, K. S. Nisar, and Y.-M. Chu, *On new modifications governed by quantum Hahn's integral operator pertaining to fractional calculus*, J. Funct. Spaces **2020** (2020), 8262860.

- [17] J.-M. Shen, S. Rashid, M. A. Noor, R. Ashraf, and Y.-M. Chu, *Certain novel estimates within fractional calculus theory on time scales*, AIMS Math. **5** (2020), no. 6, 6073–6086.
- [18] H.-X. Qi, M. Yussouf, S. Mehmood, Y.-M. Chu, and G. Farid, *Fractional integral versions of Hermite-Hadamard type inequality for generalized exponentially convexity*, AIMS Math. **5** (2020), no. 6, 6030–6042.
- [19] T. Abdeljawad, S. Rashid, Z. Hammouch, and Y.-M. Chu, *Some new local fractional inequalities associated with generalized (s, m) -convex functions and applications*, Adv. Differ. Equ. **2020** (2020), 406.
- [20] X.-Z. Yang, G. Farid, W. Nazeer, Y.-M. Chu, and C.-F. Dong, *Fractional generalized Hadamard and Fejér-Hadamard inequalities for m -convex function*, AIMS Math. **5** (2020), no. 6, 6325–6340.
- [21] M. Adil Khan, M. Hanif, Z. A. Khan, K. Ahmad, and Y.-M. Chu, *Association of Jensen's inequality for s -convex function with Csiszár divergence*, J. Inequal. Appl. **2019** (2019), 162.
- [22] P.-Y. Yan, Q. Li, Y.-M. Chu, S. Mukhtar, and S. Waheed, *On some fractional integral inequalities for generalized strongly modified h -convex function*, AIMS Math. **5** (2020), no. 6, 6620–6638.
- [23] P. Agarwal, M. Kadakal, İ. İşcan, and Y.-M. Chu, *Better approaches for n -times differentiable convex functions*, Mathematics **8** (2020), 950.
- [24] Y. Khurshid, M. Adil Khan and Y.-M. Chu, *Conformable integral version of Hermite-Hadamard-Fejér inequalities via η -convex functions*, AIMS Math. **5** (2020), no. 5, 5106–5120.
- [25] M. A. Latif, S. Rashid, S. S. Dragomir, and Y.-M. Chu, *Hermite-Hadamard type inequalities for co-ordinated convex and quasi-convex functions and their applications*, J. Inequal. Appl. **2019** (2019), 317.
- [26] H. Ge-JiLe, S. Rashid, M. A. Noor, A. Suhail, and Y.-M. Chu, *Some unified bounds for exponentially tgs -convex functions governed by conformable fractional operators*, AIMS Math. **5** (2020), no. 6, 6108–6123.
- [27] M.-B. Sun and Y.-M. Chu, *Inequalities for the generalized weighted mean values of g -convex functions with applications*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. **114** (2020), no. 4, 172.
- [28] S.-Y. Guo, Y.-M. Chu, G. Farid, S. Mehmood, and W. Nazeer, *Fractional Hadamard and Fejér-Hadamard inequalities associated with exponentially (s, m) -convex functions*, J. Funct. Spaces **2020** (2020), 2410385.
- [29] I. Abbas Baloch, A. A. Mughal, Y.-M. Chu, A. U. Haq, and M. De La Sen, *A variant of Jensen-type inequality and related results for harmonic convex functions*, AIMS Math. **5** (2020), no. 6, 6404–6418.
- [30] M. U. Awan, S. Talib, M. A. Noor, Y.-M. Chu, and K. I. Noor, *Some trapezium-like inequalities involving functions having strongly n -polynomial preinvexity property of higher order*, J. Funct. Spaces **2020** (2020), 9154139.
- [31] Y.-M. Chu, M. U. Awan, M. Z. Javad, and A. W. Khan, *Bounds for the remainder in Simpson's inequality via n -polynomial convex functions of higher order using Katugampola fractional integrals*, J. Math. **2020** (2020), 4189036.
- [32] S. Zaheer Ullah, M. Adil Khan and Y.-M. Chu, *A note on generalized convex functions*, J. Inequal. Appl. **2019** (2019), 291.
- [33] S. Khan, M. Adil Khan and Y.-M. Chu, *Converses of the Jensen inequality derived from the Green functions with applications in information theory*, Math. Methods Appl. Sci. **43** (2020), no. 5, 2577–2587.
- [34] H. Kalsoom, M. Idrees, D. Baleanu, and Y.-M. Chu, *New estimates of $q_1 q_2$ -Ostrowski-type inequalities within a class of n -polynomial preconvexity of function*, J. Funct. Spaces **2020** (2020), 3720798.
- [35] T.-H. Zhao, Z.-Y. He and Y.-M. Chu, *On some refinements for inequalities involving zero-balanced hypergeometric function*, AIMS Math. **5** (2020), no. 6, 6479–6495.
- [36] Z.-H. Yang, W.-M. Qian, W. Zhang, and Y.-M. Chu, *Notes on the complete elliptic integral of the first kind*, Math. Inequal. Appl. **23** (2020), no. 1, 77–93.
- [37] M.-K. Wang, Y.-M. Chu, Y.-M. Li, and W. Zhang, *Asymptotic expansion and bounds for complete elliptic integrals*, Math. Inequal. Appl. **23** (2020), no. 3, 821–841.
- [38] M.-K. Wang, H.-H. Chu and Y.-M. Chu, *Precise bounds for the weighted Hölder mean of the complete p -elliptic integrals*, J. Math. Anal. Appl. **480** (2019), no. 2, 123388.
- [39] T.-H. Zhao, M.-K. Wang and Y.-M. Chu, *A sharp double inequality involving generalized complete elliptic integral of the first kind*, AIMS Math. **5** (2020), no. 5, 4512–4528.
- [40] M.-K. Wang, H.-H. Chu, Y.-M. Li, and Y.-M. Chu, *Answers to three conjectures on convexity of three functions involving complete elliptic integrals of the first kind*, Appl. Anal. Discrete Math. **14** (2020), no. 1, 255–271.
- [41] M.-K. Wang, Z.-Y. He and Y.-M. Chu, *Sharp power mean inequalities for the generalized elliptic integral of the first kind*, Comput. Methods Funct. Theory **20** (2020), no. 1, 111–124.
- [42] S. Rashid, M. A. Noor, K. I. Noor, F. Safdar, and Y.-M. Chu, *Hermite-Hadamard type inequalities for the class of convex functions on time scale*, Math. **7** (2019), no. 10, 956, DOI: <https://doi.org/10.3390/math7100956>.
- [43] A. Iqbal, M. Adil Khan, N. Mohammad, E. R. Nwaeze, and Y.-M. Chu, *Revisiting the Hermite-Hadamard integral inequality via a Green function*, AIMS Math. **5** (2020), no. 6, 6087–6107.
- [44] M. A. Latif and M. Alomari, *Hadamard-type inequalities for product two convex functions on the co-ordinates*, Int. Math. Forum **4** (2009), no. 45-48, 2327–2338.
- [45] D. S. Mitrinović and I. B. Lacković, *Hermite and convexity*, Aequationes Math. **28** (1985), no. 3, 229–232.
- [46] F. Zafar, H. Kalsoom and N. Hussain, *Some inequalities of Hermite-Hadamard type for n -times differentiable (ρ, m) -geometrically convex functions*, J. Nonlinear Sci. Appl. **8** (2015), no. 3, 201–217.
- [47] S. S. Dragomir, R. P. Agarwal and P. Cerone, *On Simpson's inequality and applications*, J. Inequal. Appl. **5** (2000), no. 6, 533–579.

- [48] M. U. Awan, S. Talib, A. Kashuri, M. A. Noor, and Y.-M. Chu, *Estimates of quantum bounds pertaining to new q -integral identity with applications*, Adv. Differ. Equ. **2020** (2020), 424.
- [49] F. H. Jackson, *On a q -definite integrals*, Quart. J. Pure Appl. Math. **41** (1910), 193–203.
- [50] J. Tariboon and S. K. Ntouyas, *Quantum integral inequalities on finite intervals*, J. Inequal. Appl. **2014** (2014), 121.
- [51] J. Tariboon and S. K. Ntouyas, *Quantum calculus on finite intervals and applications to impulsive difference equations*, Adv. Differ. Equ. **2013** (2013), 282.
- [52] V. Kac and P. Cheung, *Quantum Calculus*, Springer-Verlag, New York, 2002.
- [53] H. Gauchman, *Integral inequalities in q -calculus*, Comput. Math. Appl. **47** (2004), no. 2–3, 281–300.
- [54] M. A. Noor, M. U. Awan and K. I. Noor, *Quantum Ostrowski inequalities for q -differentiable convex functions*, J. Math. Inequal. **10** (2016), no. 4, 1013–1018.
- [55] W. Sudsutad, S. K. Ntouyas and J. Tariboon, *Quantum integral inequalities for convex functions*, J. Math. Inequal. **9** (2015), no. 3, 781–793.
- [56] M. A. Noor, K. I. Noor and M. U. Awan, *Some quantum integral inequalities via preinvex functions*, Appl. Math. Comput. **269** (2015), 242–251.
- [57] M. A. Noor, K. I. Noor and M. U. Awan, *Some quantum estimates for Hermite-Hadamard inequalities*, Appl. Math. Comput. **251** (2015), 675–679.
- [58] W. J. Liu and H. F. Zhuang, *Some quantum estimates of Hermite-Hadamard inequalities for convex function*, J. Appl. Anal. Comput. **7** (2017), no. 2, 501–522.
- [59] M. Tunç and E. Göv, *(p,q) -integral inequalities*, RGMIA Res. Rep. Coll. **19** (2016), 97.
- [60] M. Tunç and E. Göv, *Some integral inequalities via (p,q) -calculus on finite intervals*, RGMIA Res. Rep. Coll. **19** (2016), 95.
- [61] M. Kunt, İ. İşcan, N. Alp, and M. Z. Sarikaya, *(p,q) -Hermite-Hadamard inequalities and (p,q) -estimates for midpoint type inequalities via convex and quasi-convex functions*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. **112** (2018), no. 4, 969–992.
- [62] M. Mursaleen, K. J. Ansari and A. Khan, *Some approximation results by (p,q) -analogue of Bernstein-Stancu operators*, Appl. Math. Comput. **264** (2015), 392–402.
- [63] V. G. Miheşan, *A generalization of the convexity*, in: Seminar on Functional Equations, Approximation and Convexity, Cluj-Napoca, Romania, 1993.
- [64] Y. Zhang, T.-S. Du, H. Wang, and Y.-J. Shen, *Different types of quantum integral inequalities via (α,m) -convexity*, J. Inequal. Appl. **2018** (2018), 264.
- [65] N. Alp, M. Z. Sarikaya, M. Kunt, and İ. İşcan, *q -Hermite Hadamard inequalities and quantum estimates for midpoint type inequalities via convex and quasi-convex functions*, J. King Saud Univ. Sci. **30** (2018), no. 2, 193–203.
- [66] M. A. Latif, S. S. Dragomir and E. Momoniat, *Some q -analogues of Hermite-Hadamard inequality of functions of two variables on finite rectangles in the plane*, J. King Saud Univ. Sci. **29** (2017), no. 3, 263–273.
- [67] M. Alomari, M. Darus and S. S. Dragomir, *New inequalities of Simpson's type for s -convex functions with applications*, RGMIA Res. Rep. Coll. **12** (2009), no. 4, 9.
- [68] B.-Y. Xi and F. Qi, *Some Hermite-Hadamard type inequalities for differentiable convex functions and applications*, Hacet. J. Math. Stat. **42** (2013), no. 3, 243–257.