



---

*Research article*

## Odd-order differential equations with deviating arguments: asymptomatic behavior and oscillation

A. Muhib<sup>1,2</sup>, I. Dassios<sup>3</sup>, D. Baleanu<sup>4,5,6</sup>, S. S. Santra<sup>7</sup> and O. Moaaz<sup>1,8,\*</sup>

<sup>1</sup> Department of Mathematics, Faculty of Science, Mansoura University, Mansoura 35516, Egypt

<sup>2</sup> Department of Mathematics, Faculty of Education – Al-Nadirah, Ibb University, Ibb, Yemen

<sup>3</sup> AMPSAS, University College Dublin, D4 Dublin, Ireland

<sup>4</sup> Department of Mathematics and Computer Science, Faculty of Arts and Sciences, Çankaya University Ankara, 06790 Etimesgut, Turkey

<sup>5</sup> Institute of Space Sciences, Magurele-Bucharest, 077125 Magurele, Romania; Department of Medical Research, China

<sup>6</sup> Medical University Hospital, China Medical University, Taichung, 40402, Taiwan, China

<sup>7</sup> Department of Mathematics, JIS College of Engineering, Kalyani, West Bengal - 741235, India

<sup>8</sup> Section of Mathematics, International Telematic University Uninettuno, Corso Vittorio Emanuele II, 39, 00186 Roma, Italy

\* **Correspondence:** Email: o\_moaaz@mans.edu.eg.

**Abstract:** Despite the growing interest in studying the oscillatory behavior of delay differential equations of even-order, odd-order equations have received less attention. In this work, we are interested in studying the oscillatory behavior of two classes of odd-order equations with deviating arguments. We get more than one criterion to check the oscillation in different methods. Our results are an extension and complement to some results published in the literature.

**Keywords:** differential equation with deviating argument; odd-order; neutral; oscillation; asymptotic behavior

---

### 1. Introduction

This study investigates the oscillatory and asymptotic behavior of delay differential equations (DDEs) of odd-order

$$(a(\eta) (\psi^{(n-1)}(\eta))^{\kappa})' + q(\eta) f(\psi(\phi(\eta))) = 0 \quad (1.1)$$

and

$$(a(\eta)(v^{(n-1)}(\eta))^\kappa)' + q(\eta)f(\psi(\phi(\eta))) = 0, \quad (1.2)$$

where  $n \geq 3$  is an odd integer and  $v(\eta) = \psi(\eta) + p(\eta)\psi(\tau(\eta))$ . Further, we assume that:

- (i)  $\kappa$  is a ratio of odd natural numbers;
- (ii)  $q, p \in C([\eta_0, \infty), (0, \infty))$  and  $0 \leq p(\eta) < 1$ ;
- (iii)  $a, \tau, \phi \in C^1([\eta_0, \infty))$ ,  $a(\eta) > 0$ ,  $a'(\eta) \geq 0$ ,  $\phi(\eta) \geq \eta \geq \tau(\eta)$ ,  $\lim_{\eta \rightarrow \infty} \tau(\eta) = \infty$ ;
- (iv)  $f \in C(\mathbb{R}, \mathbb{R})$ ,  $f(\psi) \geq k\psi^\kappa$  and

$$\pi(\eta) = \int_{\eta_0}^{\eta} \frac{1}{a^{1/\kappa}(s)} ds \rightarrow \infty \text{ as } \eta \rightarrow \infty. \quad (1.3)$$

If there exists a  $\eta_\psi \geq \eta_0$  with a continuous function  $\psi$  satisfies (1.1),  $a(\eta)(\psi^{(n-1)})^\kappa(\eta) \in C^1([\eta_\psi, \infty), \mathbb{R})$ , and  $\sup\{|\psi(\eta)| : \eta_1 \leq \eta\} > 0$  for every  $\eta_1, \eta \in [\eta_\psi, \infty)$ , then  $\psi$  is said to be a proper solution of (1.1). A solution  $\psi$  of (1.1) is said to be non-oscillatory if it is positive or negative, ultimately; otherwise, it is said to be oscillatory.

Lately, great attention has been devoted to the theory of oscillation in DDEs. The works [1–10] develop techniques and methods for studying the oscillations of second-order DDEs. This development was necessarily reflected in the study of the oscillation of even-order DDEs, and this can be seen through the works, for example [11–18]. On the other hand, odd-order DDEs have received less attention compared to even-order DDEs. The development of the study of such equations can be traced through papers, [19–29], and the references cited therein.

Baculikova and Dzurina [30] studied the asymptotic properties of neutral DDE

$$(a(\eta)((\psi(\eta) \pm p(\eta)\psi(\delta(\eta)))''^\kappa)' + q(\eta)\psi^\kappa(\phi(\eta)) = 0.$$

Li and Rogovchenko [31] investigated the oscillation of neutral DDE

$$(a(\eta)(v''(\eta))^\kappa)' + q(\eta)\psi^\kappa(\phi(\eta)) = 0,$$

where  $v(\eta) = \psi(\eta) + p_0\psi(\eta - \delta_0)$  and  $\delta_0 \geq 0$  (delayed argument) or  $\delta_0 \leq 0$  (advanced argument). Lackova [32] deduced oscillatory and asymptotic behavior of neutral DDE

$$(\psi(\eta) + p(\eta)\psi(\tau(\eta)))^{(n)} + q(\eta)f(\psi(\phi(\eta))) = 0,$$

where  $\phi(\eta)$  is a delayed argument and  $n \geq 2$ ,  $f(\varrho) \operatorname{sgn} \varrho \geq k|\varrho|^\kappa$ ,  $\kappa \geq 1$ ,  $k > 0$ .

The half-linear differential equations arise in the study of  $p$  Laplace equations, porous medium problems, chemotaxis models, and so forth; see, for instance, the papers [33–35] for more details. In this study, we investigate the oscillatory and asymptotic properties of solutions for half-linear differential equations (1.1) and (1.2) under the conditions mentioned above and present some new results which are complementary and extend to [30–32]. We will support the results obtained with two examples.

## 2. Auxiliary lemmas

We start with some lemmas that we will need to use later. The next result is a well-known result; see [36, Lemma 2], also see [37, Lemma 2.2.1].

**Lemma 2.1.** *If  $\psi$  is a solution of (1.1) and positive eventually, then  $\psi^{(k)}(\eta)$ ,  $1 \leq k \leq n-1$ , are of constant signs,  $a(\eta) \left(\psi^{(n-1)}(\eta)\right)^k$  is decreasing. Moreover,  $\psi$  satisfies either*

$$\psi'(\eta) > 0, \psi''(\eta) > 0, \psi^{(n-1)}(\eta) > 0, \psi^{(n)}(\eta) < 0 \quad (2.1)$$

or

$$(-1)^m \psi^{(m)} > 0, m = 1, 2, \dots, n. \quad (2.2)$$

**Lemma 2.2.** [36] *Let  $\psi \in C^n([\eta_0, \infty), (0, \infty))$ ,  $\psi^{(n-1)}(\eta)\psi^{(n)}(\eta) \leq 0$  for  $\eta \geq \eta_\psi$  and assume that  $\lim_{\eta \rightarrow \infty} \psi(\eta) \neq 0$ , then there exists an  $\eta_\theta \in [\eta_\psi, \infty)$  with*

$$\psi(\eta) \geq \frac{\theta}{(n-1)!} \eta^{n-1} |\psi^{(n-1)}(\eta)| \text{ for all } \eta \in [\eta_\theta, \infty) \text{ and } \theta \in (0, 1).$$

**Lemma 2.3.** *Assume that  $\psi^{(i)}(\eta) > 0$  for  $i = 0, 1, 2$ , eventually. Then, for all  $\delta_0 \in (0, 1)$ ,*

$$\psi'(\eta) \geq \frac{\delta_0}{\eta} \psi(\eta)$$

and

$$\psi(\phi(\eta)) \geq \left(\frac{\phi(\eta)}{\eta}\right)^{\delta_0} \psi(\eta). \quad (2.3)$$

*Proof.* Assume that  $\psi^{(i)}(\eta) > 0$  for  $i = 0, 1, 2$  and for all  $\eta \geq \eta_1 \geq \eta_0$ ,  $\eta_1$  large enough. Then, we get

$$\psi(\phi(\eta)) - \psi(\eta) = \int_{\eta}^{\phi(\eta)} \psi'(s) ds \geq \psi'(\eta)(\phi(\eta) - \eta). \quad (2.4)$$

It is easy to notice that  $\lim_{\eta \rightarrow \infty} \psi(\eta) = \infty$ . Hence, there exists  $\eta_2 \geq \eta_1$  large enough such that

$$\delta_0 \psi(\eta) \leq \psi(\eta) - \psi(\eta_2) = \int_{\eta_2}^{\eta} \psi'(s) ds \leq \psi'(\eta)(\eta - \eta_2) \leq \eta \psi'(\eta), \quad (2.5)$$

for all  $\delta_0 \in (0, 1)$ . By integrating this inequality from  $\eta$  to  $\phi(\eta)$ , we find

$$\psi(\phi(\eta)) \geq \left(\frac{\phi(\eta)}{\eta}\right)^{\delta_0} \psi(\eta).$$

This completes the proof of this Lemma.

**Lemma 2.4.** *Suppose that  $\psi$  is a positive solution of (1.2). Then,  $(a(\eta) \left(v^{(n-1)}(\eta)\right)^k)' < 0$ ,  $v^{(i)}(\eta)$ ,  $1 \leq i \leq n-1$ , are of constant signs, and  $v(\eta)$  satisfies either*

$$\text{Case (1): } v(\eta) > 0, v'(\eta) > 0, v''(\eta) > 0, v^{(n-1)}(\eta) > 0 \text{ and } v^{(n)}(\eta) \leq 0 \quad (2.6)$$

or

$$\text{Case (2): } (-1)^k v^{(k)}(\eta) > 0, \text{ for } k = 1, 2, \dots, n. \quad (2.7)$$

*Proof.* Suppose that  $\psi$  is a solution of (1.2) and positive eventually. Produces directly from (1.2) that

$$\left(a(\eta)\left(v^{(n-1)}(\eta)\right)^k\right)' \leq -kq(\eta)\psi^k(\phi(\eta)) < 0. \quad (2.8)$$

Now, from the above inequality we find either  $v^{(n-1)}(\eta) > 0$  or  $v^{(n-1)}(\eta) < 0$ .

If  $v^{(n-1)}(\eta) < 0$ , then

$$a(\eta)\left(v^{(n-1)}(\eta)\right)^k < -c < 0,$$

integration from  $\eta_1$  to  $\eta$ , we have

$$v^{(n-2)}(\eta) < v^{(n-2)}(\eta_1) - c^{1/k} \int_{\eta_1}^{\eta} \frac{1}{a^{1/k}(s)} ds,$$

by using (1.3) we have  $v^{(n-2)}(\eta) \rightarrow -\infty$  at  $\eta \rightarrow \infty$ , and by doing this process several times we get  $v(\eta) \rightarrow -\infty$ . This contradicts the positive  $v(\eta)$ , then  $v^{(n-1)}(\eta) > 0$ . Since  $v^{(n-1)}(\eta) > 0$ , we have that either  $v^{(n-2)}(\eta) > 0$  or  $v^{(n-2)}(\eta) < 0$ . But,  $v^{(n-2)}(\eta) > 0$  leads to  $v^{(i)}(\eta) > 0$  for  $0 \leq i \leq n-2$ . Repeating these considerations, we verify that  $v(\eta)$  satisfies either (2.6) or (2.7).

Now since  $v^{(n-1)}(\eta) > 0$  and  $a' \geq 0$ . Then we have

$$0 > \left(a(\eta)\left(v^{(n-1)}(\eta)\right)^k\right)' = a'(\eta)\left(v^{(n-1)}(\eta)\right)^k + \kappa a(\eta)\left(v^{(n-1)}(\eta)\right)^{k-1} v^{(n)}(\eta),$$

which shows us that  $v^{(n)}(\eta) < 0$ . This completes the proof of this lemma.

**Lemma 2.5.** *Let Case (1) hold. Then*

$$\frac{v(\phi(\eta))}{v(\eta)} \geq \left(\frac{\phi(\eta)}{\eta}\right)^{\delta_0}, \quad (2.9)$$

for all  $\delta_0 \in (0, 1)$ .

*Proof.* The proof of the above lemma is similar to that of Lemma 2.3 and so it is omitted.

Next, we will present the basic definitions and notations that we will use in our results.  $\{h_m(\eta)\}_{m=0}^{\infty}$  is a sequence of continuous functions defined as follows

$$h_0(\eta) = k\Psi(\eta), \quad k \in (0, 1) \text{ fixed},$$

$$h_{m+1}(\eta) = h_0(\eta) + \frac{\kappa k}{(n-2)!} \int_{\eta}^{\infty} h_m^{(\kappa+1)/\kappa}(s) \frac{s^{n-2}}{a^{1/\kappa}(s)} ds, \quad m = 0, 1, \dots, \quad (2.10)$$

$$\Psi(\eta) = \int_{\eta}^{\infty} q(s) \left(\frac{\phi(s)}{s}\right)^{k\delta_0} (1-p(\phi(s)))^k ds$$

and

$$\Theta(\eta) = \frac{k^{1/\kappa}}{a^{1/\kappa}(\eta)} \left(\int_{\eta}^{\infty} q(s) ds\right)^{1/\kappa},$$

where  $\delta_0 \in (0, 1)$ .

### 3. Main results

Now, we present our results for (1.1) and (1.2).

**Theorem 3.1.** Assume that

$$\liminf_{\eta \rightarrow \infty} \frac{1}{\Upsilon(\eta)} \int_{\eta}^{\infty} \frac{s^{n-2} \Upsilon^{(\kappa+1)/\kappa}(s)}{a^{1/\kappa}(s)} ds > \frac{(n-2)!}{(\kappa+1)^{(\kappa+1)/\kappa}} \quad (3.1)$$

and

$$\int_{\eta_0}^{\infty} s^{n-2} \frac{1}{a^{1/\kappa}(s)} \left( \int_s^{\infty} q(\varrho) d\varrho \right)^{1/\kappa} ds = \infty, \quad (3.2)$$

where

$$\Upsilon(\eta) = \int_{\eta}^{\infty} q(s) \left( \frac{\phi(s)}{s} \right)^{\kappa \delta_0} ds,$$

then  $\lim_{\eta \rightarrow \infty} \psi(\eta) = 0$ , for every nonoscillatory solution  $\psi(\eta)$  of (1.1).

*Proof.* Assume that  $\psi$  is a solution of (1.1), positive eventually, and satisfies (2.1). By (3.1), we have

$$\liminf_{\eta \rightarrow \infty} \frac{k^{(\kappa+1)/\kappa}}{\Upsilon(\eta)} \int_{\eta}^{\infty} \frac{s^{n-2} \Upsilon^{(\kappa+1)/\kappa}(s)}{a^{1/\kappa}(s)} ds > \frac{(n-2)!}{(\kappa+1)^{1+1/\kappa}}, \quad (3.3)$$

for some  $k \in (0, 1)$ . From Lemma 2.3 and (1.1), we have

$$\left( a(\eta) \left( \psi^{(n-1)}(\eta) \right)^{\kappa} \right)' + kq(\eta) \left( \frac{\phi(\eta)}{\eta} \right)^{\kappa \delta_0} \psi^{\kappa}(\eta) \leq 0. \quad (3.4)$$

Now we define  $\varpi(\eta)$  as follows

$$\varpi(\eta) = \frac{a(\eta) \left( \psi^{(n-1)}(\eta) \right)^{\kappa}}{\psi^{\kappa}(\eta)}, \quad (3.5)$$

then

$$\varpi'(\eta) = \frac{\left( a(\eta) \left( \psi^{(n-1)}(\eta) \right)^{\kappa} \right)'}{\psi^{\kappa}(\eta)} - \kappa \frac{a(\eta) \left( \psi^{(n-1)}(\eta) \right)^{\kappa} \psi'(\eta)}{\psi^{\kappa+1}(\eta)}.$$

By using (3.4) and (3.5), we get

$$\varpi'(\eta) \leq -kq(\eta) \left( \frac{\phi(\eta)}{\eta} \right)^{\kappa \delta_0} - \kappa \varpi(\eta) \frac{\psi'(\eta)}{\psi(\eta)}.$$

By using Lemma 2.2, we have

$$\varpi'(\eta) \leq -kq(\eta) \left( \frac{\phi(\eta)}{\eta} \right)^{\kappa \delta_0} - \kappa \varpi(\eta) \frac{k\eta^{n-2} \psi^{(n-1)}(\eta)}{(n-2)! \psi(\eta)},$$

from (3.5), we get

$$\varpi'(\eta) \leq -kq(\eta) \left( \frac{\phi(\eta)}{\eta} \right)^{\kappa \delta_0} - \kappa \frac{k\eta^{n-2}}{a^{1/\kappa}(\eta) (n-2)!} \varpi^{1+1/\kappa}(\eta). \quad (3.6)$$

Integrating (3.6) from  $\eta$  to  $\infty$ , we have

$$\varpi(\eta) \geq k\Upsilon(\eta) + \frac{\kappa k}{(n-2)!} \int_{\eta}^{\infty} \frac{s^{n-2}}{a^{1/\kappa}(s)} \varpi^{1+1/\kappa}(s) ds \quad (3.7)$$

or

$$\frac{\varpi(\eta)}{k\Upsilon(\eta)} \geq 1 + \frac{\kappa k^{(\kappa+1)/\kappa}}{(n-2)! \Upsilon(\eta)} \int_{\eta}^{\infty} \frac{s^{n-2} \Upsilon^{(\kappa+1)/\kappa}(s)}{a^{1/\kappa}(s)} \left( \frac{\varpi(s)}{k\Upsilon(s)} \right)^{(\kappa+1)/\kappa} ds,$$

eventually, let us say  $\eta \geq \eta_1$ . Since

$$\varpi(\eta) > k\Upsilon(\eta),$$

then

$$\inf_{\eta \geq \eta_1} \frac{\varpi(\eta)}{k\Upsilon(\eta)} = \zeta \geq 1,$$

thus

$$\frac{\varpi(\eta)}{k\Upsilon(\eta)} \geq 1 + \frac{\kappa (k\zeta)^{(\kappa+1)/\kappa}}{(n-2)! \Upsilon(\eta)} \int_{\eta}^{\infty} \frac{s^{n-2} \Upsilon^{(\kappa+1)/\kappa}(s)}{a^{1/\kappa}(s)} ds, \quad (3.8)$$

from (3.3), we have

$$\frac{k^{(\kappa+1)/\kappa}}{(n-2)! \Upsilon(\eta)} \int_{\eta}^{\infty} \frac{s^{n-2} \Upsilon^{(\kappa+1)/\kappa}(s)}{a^{1/\kappa}(s)} ds > \alpha > (\kappa+1)^{-(\kappa+1)/\kappa}, \quad (3.9)$$

for some positive  $\alpha$ . From (3.8) and (3.9), we have

$$\frac{\varpi(\eta)}{k\Upsilon(\eta)} \geq 1 + \kappa \zeta^{(\kappa+1)/\kappa} \alpha, \quad (3.10)$$

therefore

$$\zeta \geq 1 + \kappa \zeta^{(\kappa+1)/\kappa} \alpha > 1 + \kappa \zeta^{(\kappa+1)/\kappa} (\kappa+1)^{-(\kappa+1)/\kappa},$$

that is,

$$0 \geq \frac{1}{\kappa+1} + \frac{\kappa}{\kappa+1} \left( \frac{\zeta}{\kappa+1} \right)^{(\kappa+1)/\kappa} - \frac{\zeta}{\kappa+1}.$$

But, we have

$$f(\vartheta) = \frac{1}{\kappa+1} + \frac{\kappa}{\kappa+1} \vartheta^{(\kappa+1)/\kappa} - \vartheta$$

is a non-negative function for every  $\vartheta > 0$ . Thus, we obtain that  $\psi$  cannot satisfy (2.1).

Next, we suppose that (2.2) holds. Then there exists a finite  $\lim_{\eta \rightarrow \infty} \psi(\eta) = D$ . Suppose the contrary that  $D > 0$ . By integrating (1.1) over  $[\eta, \infty)$ , we have

$$a(\eta) (\psi^{(n-1)}(\eta))^{\kappa} \geq k \int_{\eta}^{\infty} q(s) \psi^{\kappa}(\phi(s)) ds \geq kD^{\kappa} \int_{\eta}^{\infty} q(s) ds,$$

that is,

$$\psi^{(n-1)}(\eta) \geq \frac{k^{1/\kappa} D}{a^{1/\kappa}(\eta)} \left( \int_{\eta}^{\infty} q(s) ds \right)^{1/\kappa}. \quad (3.11)$$

Integrating (3.11)  $n - 2$  times, we obtain

$$-\psi'(\eta) \geq \frac{k^{1/\kappa} D}{(n-3)!} \int_{\eta}^{\infty} \left( (s-\eta)^{n-3} \frac{1}{a^{1/\kappa}(s)} \left( \int_s^{\infty} q(\varrho) d\varrho \right)^{1/\kappa} \right) ds, \tag{3.12}$$

integrating (3.12) from  $\eta_1$  to  $\infty$ , we get

$$\begin{aligned} \psi(\eta_1) &\geq \frac{k^{1/\kappa} D}{(n-2)!} \int_{\eta_1}^{\infty} \left( (s-\eta_1)^{n-2} \frac{1}{a^{1/\kappa}(s)} \left( \int_s^{\infty} q(\varrho) d\varrho \right)^{1/\kappa} \right) ds \\ &\geq \frac{k^{1/\kappa} D}{2^{n-2} (n-2)!} \int_{2\eta_1}^{\infty} \left( s^{n-2} \frac{1}{a^{1/\kappa}(s)} \left( \int_s^{\infty} q(\varrho) d\varrho \right)^{1/\kappa} \right) ds, \end{aligned}$$

which contradicts (3.2). Then,  $\lim_{\eta \rightarrow \infty} \psi(\eta) = 0$ . The proof is complete.

**Theorem 3.2.** *If*

$$\liminf_{\eta \rightarrow \infty} \frac{1}{\Psi(\eta)} \int_{\eta}^{\infty} \frac{s^{n-2} \Psi^{(\kappa+1)/\kappa}(s)}{a^{1/\kappa}(s)} ds > \frac{(n-2)!}{(\kappa+1)^{(\kappa+1)/\kappa}} \tag{3.13}$$

and

$$\int_{\eta_0}^{\infty} s^{n-2} \Theta(s) ds = \infty, \tag{3.14}$$

then every nonoscillatory solution  $\psi(\eta)$  of (1.2) satisfies  $\lim_{\eta \rightarrow \infty} \psi(\eta) = 0$ .

*Proof.* Suppose that  $\psi$  is a solution of (1.2) and positive eventually. Assume that Case (1) holds. Since

$$\psi(\eta) = v(\eta) - p(\eta)\psi(\tau(\eta)) \geq v(\eta) - p(\eta)v(\tau(\eta)),$$

From  $\eta \geq \tau(\eta)$ , we have

$$\psi(\eta) \geq v(\eta) - p(\eta)v(\tau(\eta)) \geq v(\eta) - p(\eta)v(\eta) = v(\eta)(1 - p(\eta))$$

and so

$$\psi(\phi(\eta)) \geq v(\phi(\eta))(1 - p(\phi(\eta))), \tag{3.15}$$

from Lemma 2.2, we have

$$v'(\eta) \geq \frac{k\eta^{n-2}}{(n-2)!} v^{(n-1)}(\eta). \tag{3.16}$$

From (2.9), (3.15) and (2.8), we have

$$\left( a(\eta) \left( v^{(n-1)}(\eta) \right)^{\kappa} \right)' \leq -kq(\eta) \left( \frac{\phi(\eta)}{\eta} \right)^{\kappa \delta_0} (1 - p(\phi(\eta)))^{\kappa} v^{\kappa}(\eta). \tag{3.17}$$

Now, we define the function  $\varphi(\eta)$  as follows

$$\varphi(\eta) = \frac{a(\eta) \left( v^{(n-1)}(\eta) \right)^{\kappa}}{v^{\kappa}(\eta)}. \tag{3.18}$$

Differentiating (3.18) and using (3.17), we get

$$\varphi'(\eta) \leq -kq(\eta) \left( \frac{\phi(\eta)}{\eta} \right)^{\kappa\delta_0} (1 - p(\phi(\eta)))^\kappa - \frac{\kappa a(\eta) \left( v^{(n-1)}(\eta) \right)^\kappa v'(\eta)}{v^{\kappa+1}(\eta)}, \quad (3.19)$$

by using (3.16) and (3.18), we have

$$\varphi'(\eta) \leq -kq(\eta) \left( \frac{\phi(\eta)}{\eta} \right)^{\kappa\delta_0} (1 - p(\phi(\eta)))^\kappa - \frac{\kappa k \eta^{n-2}}{(n-2)! a^{1/\kappa}(\eta)} \varphi^{(\kappa+1)/\kappa}(\eta). \quad (3.20)$$

Integrating (3.20) from  $\eta$  to  $\infty$ , we have

$$\varphi(\eta) \geq k\Psi(\eta) + \frac{\kappa k}{(n-2)!} \int_{\eta}^{\infty} \frac{s^{n-2}}{a^{1/\kappa}(s)} \varphi^{(\kappa+1)/\kappa}(s) ds \quad (3.21)$$

or equivalently,

$$\frac{\varphi(\eta)}{k\Psi(\eta)} \geq 1 + \frac{\kappa k^{(1+\kappa)/\kappa}}{(n-2)! \Psi(\eta)} \int_{\eta}^{\infty} \frac{s^{n-2} \Psi^{(\kappa+1)/\kappa}(s)}{a^{1/\kappa}(s)} \frac{\varphi^{(\kappa+1)/\kappa}(s)}{(k\Psi(s))^{(\kappa+1)/\kappa}} ds, \quad (3.22)$$

eventually, let us say  $\eta \geq \eta_1$ . From (3.22), we have

$$\frac{\varphi(\eta)}{k\Psi(\eta)} \geq 1,$$

then

$$\inf_{\eta \geq \eta_1} \frac{\varphi(\eta)}{k\Psi(\eta)} = \varrho \geq 1, \quad (3.23)$$

by using (3.22) and (3.23), we have

$$\frac{\varphi(\eta)}{k\Psi(\eta)} \geq 1 + \frac{\kappa (\varrho k)^{(\kappa+1)/\kappa}}{(n-2)! \Psi(\eta)} \int_{\eta}^{\infty} \frac{s^{n-2} \Psi^{(\kappa+1)/\kappa}(s)}{a^{1/\kappa}(s)} ds, \quad (3.24)$$

by using (3.13), we have

$$\liminf_{\eta \rightarrow \infty} \frac{k^{(1+\kappa)/\kappa}}{\Psi(\eta)} \int_{\eta}^{\infty} \frac{s^{n-2} \Psi^{(1+\kappa)/\kappa}(s)}{a^{1/\kappa}(s)} ds > \frac{(n-2)!}{(\kappa+1)^{(1+\kappa)/\kappa}}, \quad (3.25)$$

for some  $k \in (0, 1)$ , from above inequality there exists some positive  $\xi$  such that

$$\frac{k^{(1+\kappa)/\kappa}}{(n-2)! \Psi(\eta)} \int_{\eta}^{\infty} \frac{s^{n-2} \Psi^{(1+\kappa)/\kappa}(s)}{a^{1/\kappa}(s)} ds > \xi > \frac{1}{(\kappa+1)^{(1+\kappa)/\kappa}}. \quad (3.26)$$

From (3.26) and (3.24), we get

$$\frac{\varphi(\eta)}{k\Psi(\eta)} \geq 1 + \kappa \xi \varrho^{(1+\kappa)/\kappa}, \quad (3.27)$$

therefore, from (3.23), we have

$$\varrho \geq 1 + \kappa \xi \varrho^{(1+\kappa)/\kappa} > 1 + \kappa \left( \frac{\varrho}{\kappa+1} \right)^{(1+\kappa)/\kappa},$$



that is,

$$\frac{1}{\kappa+1} + \frac{\kappa}{\kappa+1} \left( \frac{\varrho}{\kappa+1} \right)^{(1+\kappa)/\kappa} - \frac{\varrho}{\kappa+1} < 0.$$

This contradicts the fact that the function

$$f(\varsigma) = \frac{1}{\kappa+1} + \frac{\kappa}{\kappa+1} \varsigma^{(1+\kappa)/\kappa} - \varsigma \geq 0,$$

for all  $\varsigma > 0$ . Thus  $v(\eta)$  cannot satisfy Case (1).

Assume that Case (2) holds. Then there exists a constant  $c \geq 0$  such that  $\lim_{\eta \rightarrow \infty} \psi(\eta) = c$ . Suppose that  $c > 0$ . Integrating (1.2), we see that

$$a(\eta) \left( v^{(n-1)}(\eta) \right)^k \geq k \int_{\eta}^{\infty} q(s) \psi^k(\phi(s)) ds \geq kc^k \int_{\eta}^{\infty} q(s) ds, \quad (3.28)$$

that is,

$$v^{(n-1)}(\eta) \geq c\Theta(\eta). \quad (3.29)$$

Integrating (3.29) twice, we obtain

$$v^{(n-3)}(\eta) \geq c \int_{\eta}^{\infty} \left( \int_{\varrho}^{\infty} \Theta(s) ds \right) d\varrho = c \int_{\eta}^{\infty} \Theta(s) (s - \eta) ds, \quad (3.30)$$

integrating (3.30)  $n - 4$  times, we get

$$-v'(\eta) \geq \frac{c}{(n-3)!} \int_{\eta}^{\infty} \Theta(s) (s - \eta)^{n-3} ds, \quad (3.31)$$

integrating (3.31) from  $\eta_1$  to  $\infty$ , we get

$$v(\eta_1) \geq \frac{c}{(n-2)!} \int_{\eta_1}^{\infty} \Theta(s) (s - \eta_1)^{n-2} ds \geq \frac{c}{2^{n-2} (n-2)!} \int_{2\eta_1}^{\infty} s^{n-2} \Theta(s) ds,$$

which contradicts (3.14), and so we have verified that  $\lim_{\eta \rightarrow \infty} \psi(\eta) = 0$ . The proof is complete.

**Theorem 3.3.** *Let  $\psi$  be a nonoscillatory solution of (1.2), (3.14) hold, and*

$$\int_{\eta_0}^{\infty} q(\eta) \left( \frac{\phi(\eta)}{\eta} \right)^{k\delta_0} (1 - p(\phi(\eta)))^k \exp\left( \frac{\kappa k}{(n-2)!} \int_{\eta_0}^{\eta} h_m^{1/\kappa}(s) \frac{s^{n-2}}{a^{1/\kappa}(s)} ds \right) d\eta = \infty, \quad (3.32)$$

for some  $k \in (0, 1)$  and some  $m = 0, 1, \dots$ . Then  $\lim_{\eta \rightarrow \infty} \psi(\eta) = 0$ .

*Proof.* Suppose that  $\psi$  is a solution of (1.2) and positive eventually, we conclude from Lemma 2.4 that  $v(\eta)$  satisfies Case (1) or Case (2). If Case (1) holds, then from proof of Theorem 3.2, we find that (3.21) holds. By induction, using (3.21), we have that the sequence  $\{h_m(\eta)\}_{m=0}^{\infty}$  is nondecreasing and  $\varphi(\eta) \geq h_m(\eta)$ . Thus the sequence  $\{h_m(\eta)\}_{m=0}^{\infty}$  converges to  $h(\eta)$ . By the Lebesgue monotone convergence theorem and letting  $m \rightarrow \infty$  in (2.10), we have

$$h(\eta) = h_0(\eta) + \frac{\kappa k}{(n-2)!} \int_{\eta}^{\infty} h^{(k+1)/\kappa}(s) \frac{s^{n-2}}{a^{1/\kappa}(s)} ds,$$

since  $\varphi(\eta) \geq h_m(\eta)$ , gives

$$\begin{aligned} h'(\eta) &= -kq(\eta) \left( \frac{\phi(\eta)}{\eta} \right)^{\kappa\delta_0} (1 - p(\phi(\eta)))^\kappa - \frac{\kappa k}{(n-2)!} h^{(\kappa+1)/\kappa}(\eta) \frac{\eta^{n-2}}{a^{1/\kappa}(\eta)} \\ &\leq -kq(\eta) \left( \frac{\phi(\eta)}{\eta} \right)^{\kappa\delta_0} (1 - p(\phi(\eta)))^\kappa - \frac{\kappa k}{(n-2)!} h(\eta) h_m^{1/\kappa}(\eta) \frac{\eta^{n-2}}{a^{1/\kappa}(\eta)}. \end{aligned}$$

Therefore,

$$h'(\eta) + \frac{\kappa k}{(n-2)!} h(\eta) h_m^{1/\kappa}(\eta) \frac{\eta^{n-2}}{a^{1/\kappa}(\eta)} \leq -kq(\eta) \left( \frac{\phi(\eta)}{\eta} \right)^{\kappa\delta_0} (1 - p(\phi(\eta)))^\kappa,$$

that is,

$$\begin{aligned} &\left( h(\eta) \exp \left( \frac{\kappa k}{(n-2)!} \int_{\eta_1}^{\eta} h_m^{1/\kappa}(s) \frac{s^{n-2}}{a^{1/\kappa}(s)} ds \right) \right)' \\ &\leq -kq(\eta) \left( \frac{\phi(\eta)}{\eta} \right)^{\kappa\delta_0} (1 - p(\phi(\eta)))^\kappa \exp \left( \frac{\kappa k}{(n-2)!} \int_{\eta_1}^{\eta} h_m^{1/\kappa}(s) \frac{s^{n-2}}{a^{1/\kappa}(s)} ds \right). \end{aligned}$$

Integrating the above inequality from  $\eta_1$  to  $\eta$  we obtain

$$\begin{aligned} &h(\eta) \exp \left( \frac{\kappa k}{(n-2)!} \int_{\eta_1}^{\eta} h_m^{1/\kappa}(s) \frac{s^{n-2}}{a^{1/\kappa}(s)} ds \right) \\ &\leq h(\eta_1) - k \int_{\eta_1}^{\eta} q(\varrho) \left( \frac{\phi(\varrho)}{\varrho} \right)^{\kappa\delta_0} (1 - p(\phi(\varrho)))^\kappa \exp \left( \frac{\kappa k}{(n-2)!} \int_{\eta_1}^{\varrho} h_m^{1/\kappa}(s) \frac{s^{n-2}}{a^{1/\kappa}(s)} ds \right) d\varrho. \end{aligned} \quad (3.33)$$

Since,

$$h(\eta) \exp \left( \frac{\kappa k}{(n-2)!} \int_{\eta_1}^{\eta} h_m^{1/\kappa}(s) \frac{s^{n-2}}{a^{1/\kappa}(s)} ds \right) \geq 0, \quad (3.34)$$

letting  $\eta \rightarrow \infty$  in (3.33) and using (3.32), we obtain a contradiction with (3.34).

If Case (2) holds, then from proof of Theorem 3.2, condition (3.14) insures that  $\psi(\eta)$  tends to zero at  $\eta \rightarrow \infty$ . This completes the proof of the theorem.

**Theorem 3.4.** *Let  $\psi$  be a nonoscillatory solution of (1.2), (3.14) hold, and*

$$\limsup_{\eta \rightarrow \infty} k \frac{\eta^{\kappa(n-1)}}{a(\eta)} h_m(\eta) > ((n-1)!)^\kappa, \quad (3.35)$$

for some  $h_m(\eta)$  and for some  $k \in (0, 1)$ . Then  $\lim_{\eta \rightarrow \infty} \psi(\eta) = 0$ .

*Proof.* Suppose that  $\psi$  is an eventually positive solution of (1.2), we conclude from Lemma 2.4 that  $v(\eta)$  satisfies Case (1) or Case (2). Assume that Case (1) holds. From Lemma 2.2, we have

$$v(\eta) \geq \frac{k^{1/\kappa}}{(n-1)!} \eta^{n-1} v^{(n-1)}(\eta),$$

where  $k$  is the same as in  $h_m(\eta)$ . Then

$$\frac{1}{\varphi(\eta)} = \frac{1}{a(\eta)} \frac{\nu^\kappa(\eta)}{(\nu^{(n-1)}(\eta))^\kappa} \geq \frac{k}{a(\eta) ((n-1)!)^\kappa} \eta^{\kappa(n-1)},$$

that is,

$$((n-1)!)^\kappa \geq \frac{k}{a(\eta)} \eta^{\kappa(n-1)} \varphi(\eta).$$

Since  $\varphi(\eta) \geq h_m(\eta)$ , we have

$$((n-1)!)^\kappa \geq \frac{k}{a(\eta)} \eta^{\kappa(n-1)} h_m(\eta), \quad (3.36)$$

from (3.35) and (3.36) we get a contradiction. This completes the proof of the theorem.

**Corollary 1.** *Let (3.14) be satisfied, and*

$$\limsup_{\eta \rightarrow \infty} \frac{\eta^{\kappa(n-1)}}{a(\eta)} \int_{\eta}^{\infty} q(s) \left( \frac{\phi(s)}{s} \right)^{\kappa \delta_0} (1 - p(\phi(s)))^\kappa ds > ((n-1)!)^\kappa. \quad (3.37)$$

Then every nonoscillatory solution  $\psi(\eta)$  of (1.2) satisfies  $\lim_{\eta \rightarrow \infty} \psi(\eta) = 0$ .

*Proof.* From (3.37) there exists some  $k \in (0, 1)$  such that

$$\limsup_{\eta \rightarrow \infty} k^2 \frac{\eta^{\kappa(n-1)}}{a(\eta)} \int_{\eta}^{\infty} q(s) \left( \frac{\phi(s)}{s} \right)^{\kappa \delta_0} (1 - p(\phi(s)))^\kappa ds > ((n-1)!)^\kappa,$$

that is,

$$\limsup_{\eta \rightarrow \infty} k \frac{\eta^{\kappa(n-1)}}{a(\eta)} h_0(\eta) > ((n-1)!)^\kappa.$$

The assertion now follows from Theorem 3.4.

**Example 3.5.** Consider the differential equation of third order

$$\left( \eta (\psi''(\eta))^3 \right)' + \frac{q_0}{\eta^6} \psi^3(2\eta) = 0. \quad (3.38)$$

From (3.38), we find that  $n = 3$ ,  $\kappa = 3$ ,  $a(\eta) = \eta$ ,  $q(\eta) = q_0/\eta^6$ ,  $q_0 > 0$  and  $\phi(\eta) = 2\eta$ .

Now, from Theorem 3.1 we notice that condition (3.2) is satisfied and condition (3.1) is satisfied if  $q_0 > \frac{625}{256}$ . Thus, we obtain that all nonoscillatory solutions of (3.38) tend to zero at infinity when  $q_0 > \frac{625}{256}$ .

**Example 3.6.** Consider the third-order neutral differential equation

$$\left( \eta^{1/3} ((\psi(\eta) + p_0 \psi(\lambda\eta))'')^{1/3} \right)' + \frac{q_0}{\eta^{4/3}} \psi^{1/3}(\varpi\eta) = 0, \quad (3.39)$$

where  $p_0$  and  $q_0$  are constants,  $\kappa = 1/3$ ,  $n = 3$ ,  $a(\eta) = \eta^{1/3}$ ,  $\tau(\eta) = \lambda\eta$ ,  $\lambda \in (0, 1)$ ,  $\varpi \geq 1$ ,  $\phi(\eta) = \varpi\eta$ ,  $p(\eta) = p_0$ ,  $q(\eta) = q_0/\eta^{4/3}$  and  $q_0 > 0$ .

It is easy to get

$$\Psi(\eta) = \int_{\eta}^{\infty} q(s) \left( \frac{\phi(s)}{s} \right)^{\kappa \delta_0} (1 - p(\phi(s)))^\kappa ds = q_0 \varpi^{\delta_0/3} (1 - p_0)^{1/3} \frac{3}{\eta^{1/3}}$$

and

$$\Theta(\eta) = \frac{k^{1/\kappa}}{a^{1/\kappa}(\eta)} \left( \int_{\eta}^{\infty} q(s) ds \right)^{1/\kappa} = k^3 q_0^3 \frac{3^3}{\eta^2}$$

and condition (3.14) holds

$$\int_{\eta_0}^{\infty} s^{n-2} \Theta(s) ds = \int_{\eta_0}^{\infty} s k^3 q_0^3 \frac{3^3}{s^2} ds = 3^3 k^3 q_0^3 \int_{\eta_0}^{\infty} \frac{1}{s} ds = \infty$$

and

$$\liminf_{\eta \rightarrow \infty} \frac{1}{\Psi(\eta)} \int_{\eta}^{\infty} \frac{s^{n-2} \Psi^{(\kappa+1)/\kappa}(s)}{a^{1/\kappa}(s)} ds = 3^4 q_0^3 \varpi^{\delta_0} (1 - p_0),$$

then condition (3.13) reduces to

$$3^4 q_0^3 \varpi^{\delta_0} (1 - p_0) > \frac{1}{(4/3)^4},$$

which, by Theorem 3.2, guarantees that all nonoscillatory solutions of (3.39) tend to zero at infinity.

#### 4. Conclusions

In this paper, several new results for (1.1) and (1.2) have been presented which complement and expand some results introduced in the cited papers in the introduction. We obtain the conditions by using Riccati transformation and some analytical skill. In fact, our results are applicable in the case where  $\kappa$  is a ratio of odd positive integers. We supported the results obtained in this paper with two examples. An interesting problem for further research is to study the problem of nonoscillation for  $(a(\eta)(v^{(n-1)}(\eta))^\kappa)' + q(\eta)f(\psi(\phi(\eta))) = 0$  where  $f(\psi) \geq k\psi^\gamma$  and  $\gamma \neq \kappa$ .

#### Conflict of interest

There are no competing interests.

#### References

1. R. P. Agarwal, C. Zhang, T. Li, Some remarks on oscillation of second order neutral differential equations, *Appl. Math. Comput.*, **274** (2016), 178–181. doi: 10.1016/j.amc.2015.10.089.
2. M. Bohner, S. R. Grace, I. Jadlovská, Oscillation criteria for second-order neutral delay differential equations, *Electron. J. Qual. Theo. Differ. Equ.*, **60** (2017), 1–12. doi: 10.14232/ejqtde.2017.1.60.
3. A. Muhib, On oscillation of second-order noncanonical neutral differential equations, *J. Inequal. Appl.*, **2021** (2021), 1–11. doi: 10.1186/s13660-021-02595-x.
4. J. Dzurina, S. R. Grace, I. Jadlovská, T. Li, Oscillation criteria for second-order Emden–Fowler delay differential equations with a sublinear neutral term, *Math. Nachr.*, **293** (2020), 910–922. doi: 10.1002/mana.201800196.
5. G. E. Chatzarakis, O. Moaaz, T. Li, B. Qaraad, Some oscillation theorems for nonlinear second-order differential equations with an advanced argument, *Adva. Differ. Eq.*, **2020** (2020), 1–17. doi: 10.1186/s13662-020-02626-9.

6. O. Moaaz, E. M. Elabbasy, B. Qaraad, An improved approach for studying oscillation of generalized Emden–Fowler neutral differential equation, *J. Ineq. Appl.*, **2020** (2020), 1–18. doi: 10.1186/s13660-020-02332-w.
7. O. Moaaz, M. Anis, D. Baleanu, A. Muhib, More effective criteria for oscillation of second-order differential equations with neutral arguments, *Mathematics*, **8** (2020), 986. doi: 10.3390/math806098.
8. S. S. Santra, R. A. El-Nabulsi, Kh. M. Khedher, Oscillation of Second-Order Differential Equations With Multiple and Mixed Delays under a Canonical Operator, *Mathematics*, **9** (2021), 1323. doi: 10.3390/math9121323.
9. S. S. Santra, Kh. M. Khedher, O. Moaaz, A. Muhib, S-W Yao, Second-order impulsive delay differential systems: necessary and sufficient conditions for oscillatory or asymptotic behavior, *Symmetry*, **13** (2021), 722. doi: 10.3390/sym13040722.
10. S. S. Santra, A. K. Sethi, O. Moaaz, Kh. M. Khedher, S-W Yao, New oscillation theorems for second-order differential equations with canonical and non canonical operator via Riccati transformation, *Mathematics*, **9** (2021), 1111. doi: 10.3390/math9101111.
11. A. Alghamdi, C. Cesarano, B. Almarri, O. Bazighifan, Symmetry and its importance in the oscillation of solutions of differential equations, *Symmetry*, **13** (2021), 650. doi: 10.3390/sym13040650.
12. O. Moaaz, Ch. Park, A. Muhib, O. Bazighifan, Oscillation criteria for a class of even-order neutral delay differential equations, *J. Appl. Math. Comput.*, **63** (2020), 607–617. doi: 10.1007/s12190-020-01331-w.
13. O. Moaaz, C. Cesarano, A. Muhib, Some new oscillation results for fourth-order neutral differential equations, *Eur. J. Pure Appl. Math.*, **13** (2020), 185–199. doi: 10.29020/nybg.ejpam.v13i2.36.
14. O. Bazighifan, T. Abdeljawad, Q. M. Al-Mdallal, Differential equations of even-order with p-Laplacian like operators: qualitative properties of the solutions, *Adv. Differ. Equ.*, **2021** (2021), 96. doi: 10.1186/s13662-021-03254-7.
15. O. Bazighifan, F. Minhos, O. Moaaz, Sufficient conditions for oscillation of fourth-order neutral differential equations with distributed deviating arguments, *Axioms*, **9** (2020), 39. doi: 10.3390/axioms9020039.
16. O. Moaaz, S. Furuichi, A. Muhib, New comparison theorems for the nth order neutral differential equations with delay inequalities, *Mathematics*, **8** (2020), 454. doi: 10.3390/math8030454.
17. O. Moaaz, R. A. El-Nabulsi, O. Bazighifan, Oscillatory behavior of fourth-order differential equations with neutral delay, *Symmetry*, **12** (2020), 371. doi: 10.3390/sym12030371.
18. C. Park, O. Moaaz, O. Bazighifan, Oscillation results for higher order differential equations, *Axioms*, **9** (2020), 14. doi: 10.3390/axioms9010014.
19. M. Aktas, A. Tiryaki, A. Zafer, Oscillation criteria for third-order nonlinear functional differential equations, *Appl. Math. Lett.*, **23** (2010), 756–762. doi: 10.1016/j.aml.2010.03.003.

20. B. Baculikova, E. M. Elabbasy, S. H. Saker, J. Dzurina, Oscillation criteria for third-order nonlinear differential equations, *Math. Slovaca*, **58** (2008), 201–220. doi: 10.2478/s12175-008-0068-1.
21. O. Moaaz, J. Awrejcewicz, A. Muhib, Establishing new criteria for oscillation of odd-order nonlinear differential equations, *Mathematics*, **8** (2020), 607–617. doi: 10.3390/math8060937.
22. O. Moaaz, I. Dassios, W. Muhsin, A. Muhib, Oscillation Theory for Non-Linear Neutral Delay Differential Equations of Third Order, *Appl. Sci.*, **10** (2020), 4855. doi: 10.3390/app10144855.
23. O. Moaaz, E. M. Elabbasy, E. Shaaban, Oscillation criteria for a class of third order damped differential equations, *Arab J. Math. Sci.*, **24** (2018), 16–30. doi: 10.1016/j.ajmsc.2017.07.001.
24. O. Moaaz, D. Baleanu, A. Muhib, New aspects for non-existence of kneser solutions of neutral differential equations with odd-order, *Mathematics*, **8** (2020), 494. doi: 10.3390/math8040494.
25. S. H. Saker, J. Dzurina, On the oscillation of certain class of third-order nonlinear delay differential equations, *Math. Bohem.*, **135** (2010), 225–237.
26. R. P. Agarwal, M. Bohner, T. Li, C. Zhang, Oscillation of third-order nonlinear delay differential equations, *Taiwanese J. Math.*, **17** (2013), 545–558. doi: 10.11650/tjm.17.2013.2095.
27. T. Li, Yu. V. Rogovchenko, On asymptotic behavior of solutions to higher-order sublinear Emden-Fowler delay differential equations, *Appl. Math. Lett.*, **67** (2017), 53–59. doi: 10.1016/j.aml.2016.11.007.
28. C. Zhang, R. P. Agarwal, T. Li, Oscillation and asymptotic behavior of higher-order delay differential equations with  $\mathcal{L}$ -Laplacian like operators, *J. Math. Anal. Appl.*, **409** (2014), 1093–1106. doi: 10.1016/j.jmaa.2013.07.066.
29. C. Zhang, T. Li, B. Sun, E. Thandapani, On the oscillation of higher-order half-linear delay differential equations, *Appl. Math. Lett.*, **24** (2011), 1618–1621. doi: 10.1016/j.aml.2011.04.015.
30. B. Baculikova, J. Dzurina, Oscillation of third-order neutral differential equations, *Math. Comput. Model.*, **52** (2010), 215–226. doi: 10.1016/j.mcm.2010.02.011.
31. T. Li, Y. V. Rogovchenko, On the asymptotic behavior of solutions to a class of third-order nonlinear neutral differential equations, *Appl. Math. Lett.*, (2020), 106293. doi: 10.1016/j.aml.2020.106293.
32. D. Lackova, The asymptotic properties of the solutions of the n-th order functional neutral differential equations, *Comput. Appl. Math.*, **146** (2003), 385–392. doi: 10.1016/S0096-3003(02)00590-8.
33. M. Bohner, T. S. Hassan, T. Li, Fite-Hille-Wintner-type oscillation criteria for second-order half-linear dynamic equations with deviating arguments, *Indag. Math. (N.S.)*, **29** (2018), 548–560. doi: 10.1016/j.indag.2017.10.006.
34. T. Li, N. Pintus, G. Viglialoro, Properties of solutions to porous medium problems with different sources and boundary conditions, *Z. Angew. Math. Phys.*, **70** (2019), 1–18. doi: 10.1007/s00033-019-1130-2.
35. T. Li, G. Viglialoro, Boundedness for a nonlocal reaction chemotaxis model even in the attraction-dominated regime, *Differ. Integral Equ.*, **34** (2021), 315–336.

- 
36. B. Baculikova, J. Dzurina, On the oscillation of odd order advanced differential equations, *Bound. Value Probl.*, **214** (2014), 214. doi: 10.1186/s13661-014-0214-3.
37. R. P. Agarwal, S. R. Grace, D. O'Regan, Oscillation Theory for difference and functional differential equations, *Marcel Dekker*, Kluwer Academic, Dordrecht, 2000. doi: 10.1007/978-94-015-9401-1.



AIMS Press

©2022 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)