



Article

On Transformation Involving Basic Analogue to the Aleph-Function of Two Variables

Dinesh Kumar ^{1,*} , Dumitru Baleanu ^{2,3} , Frédéric Ayant ^{4,5} and Norbert Südland ⁶¹ Department of Applied Sciences, College of Agriculture-Jodhpur, Agriculture University Jodhpur, Jodhpur 342304, India² Department of Mathematics, Cankaya University, Ankara 06530, Turkey; dimitru@cankaya.edu.tr³ Institute of Space Sciences, 077125 Magurele-Bucharest, Romania⁴ Collège Jean L'herminier, Allée des Nymphéas, 83500 La Seyne-sur-Mer, France; fredericayant@gmail.com⁵ Department VAR, Avenue Joseph Raynaud Le parc Fleuri, 83140 Six-Fours-les-Plages, France⁶ Aage GmbH, Röntgenstraße 24, 73431 Aalen, Baden-Württemberg, Germany; norbert.suedland@aage-leichtbauteile.de

* Correspondence: dinesh_dino03@yahoo.com

Abstract: In our work, we derived the fractional order q -integrals and q -derivatives concerning a basic analogue to the Aleph-function of two variables (AFTV). We discussed a related application and the q -extension of the corresponding Leibniz rule. Finally, we presented two corollaries concerning the basic analogue to the I -function of two variables and the basic analogue to the Aleph-function of one variable.

Keywords: Fractional q -integral; q -derivative operators; basic analogue to the Aleph-function; basic analogue to the I -function; q -Leibniz rule



Citation: Kumar, D.; Baleanu, D.; Ayant, F.; Südland, N. On Transformation Involving Basic Analogue to the Aleph-Function of Two Variables. *Fractal Fract.* **2022**, *6*, 71. <https://doi.org/10.3390/fractalfract6020071>

Academic Editors: António M. Lopes, Alireza Alfi, Liping Chen and Sergio A. David

Received: 12 December 2021

Accepted: 19 January 2022

Published: 28 January 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Fractional calculus represents an important part of mathematical analysis. The concept of fractional calculus was born from a famous correspondence between L'Hopital and Leibniz in 1695. In the last four decades, it has gained significant recognition and found many applications in diverse research fields (see [1–6]). The fractional basic (or q -) calculus is the extension of the ordinary fractional calculus in the q -theory (see [7–10]). We recall that basic series and basic polynomials, particularly the basic (or q -) hypergeometric functions and basic (or q -) hypergeometric polynomials, are particularly applicable in several fields, e.g., Finite Vector Spaces, Lie Theory, Combinatorial Analysis, Particle Physics, Mechanical Engineering, Theory of Heat Conduction, Non-Linear Electric Circuit Theory, Cosmology, Quantum Mechanics, and Statistics. In 1952, Al-Salam introduced the q -analogue to Cauchy's formula [11] (see also [12]). Agarwal [13] studied certain fractional q -integral and q -derivative operators. In addition, various researchers reported image formulas of various q -special functions under fractional q -calculus operators, for example, Kumar et al. [14], Sahni et al. [15], Yadav and Purohit [16], Yadav et al. [17,18], and maybe more. The q -extensions of the Saigo's fractional integral operators were defined by Purohit and Yadav [19]. Several authors utilised such operators to evaluate a general class of q -polynomials, the basic analogue to Fox's H -function, basic analogue to the I -function, fractional q -calculus formulas for various special functions, etc. The readers can see more related new details in [16–18,20] on fractional q -calculus.

The purpose of the present manuscript is to discuss expansion formulas, involving the basic analogue to AFTV [21]. The q -Leibniz formula is also provided.

We recall that q -shifted factorial $(a; q)_n$ has the following form [22,23]

$$(a; q)_n = \begin{cases} 1, & (n = 0) \\ \prod_{i=0}^{n-1} (1 - aq^i), & (n \in \mathbb{N} \cup \{\infty\}) \end{cases} \quad (1)$$

such that $a, q \in \mathbb{C}$ and it is assumed that $a \neq q^{-m}$ ($m \in \mathbb{N}_0$).

The expression of the q -shifted factorial for negative subscript is written by

$$(a; q)_{-n} = \frac{1}{(1 - aq^{-1})(1 - aq^{-2}) \cdots (1 - aq^{-n})} \quad (n \in \mathbb{N}_0). \quad (2)$$

Additionally, we have

$$(a; q)_{\infty} = \prod_{i=0}^{\infty} (1 - aq^i) \quad (a, q \in \mathbb{C}; |q| < 1). \quad (3)$$

Using (1)–(3), we conclude that

$$(a; q)_n = \frac{(a; q)_{\infty}}{(aq^n; q)_{\infty}} \quad (n \in \mathbb{Z}), \quad (4)$$

its extension to $n = \alpha \in \mathbb{C}$ as:

$$(a; q)_{\alpha} = \frac{(a; q)_{\infty}}{(aq^{\alpha}; q)_{\infty}} \quad (\alpha \in \mathbb{C}; |q| < 1), \quad (5)$$

such that the principal value of q^{α} is considered.

We equivalently have a form of (1), given as

$$(a; q)_n = \frac{\Gamma_q(a+n)(1-q)^n}{\Gamma_q(a)} \quad (a \neq 0, -1, -2, \dots), \quad (6)$$

where the q -gamma function is expressed as [8]:

$$\Gamma_q(a) = \frac{(q; q)_{\infty}}{(q^a; q)_{\infty}(1-q)^{a-1}} = \frac{(q; q)_{a-1}}{(1-q)^{a-1}}, \quad (a \neq 0, -1, -2, \dots). \quad (7)$$

The expression of the q -analogue to the Riemann–Liouville fractional integral operator (RLI) of $f(x)$ has the following expression [12]:

$$I_q^{\mu} \{f(x)\} = \frac{1}{\Gamma_q(\mu)} \int_0^x (x-tq)_{\mu-1} f(t) d_q t, \quad (8)$$

here, $\Re(\mu) > 0$, $|q| < 1$ and

$$[x-y]_v = x^v \prod_{n=0}^{\infty} \left[\frac{1 - (y/x)q^n}{1 - (y/x)q^{n+v}} \right] = x^v \left(\frac{y}{x}; q \right)_v \quad (x \neq 0). \quad (9)$$

The basic integral [8] is given by

$$\int_0^x f(t) d_q t = x(1-q) \sum_{k=0}^{\infty} q^k f(xq^k). \quad (10)$$

Equation (8), in conjunction with (10); then, we have the series representation of (RLI), as follows

$$I_q^{\mu} f(x) = \frac{x^{\mu}(1-q)}{\Gamma_q(\mu)} \sum_{k=0}^{\infty} q^k [1 - q^{k+1}]_{\mu-1} f(xq^k). \quad (11)$$

We mention that for $f(x) = x^{\lambda-1}$, the following can be written [16]

$$I_q^{\mu} (x^{\lambda-1}) = \frac{\Gamma_q(\lambda)}{\Gamma_q(\lambda + \mu)} x^{\lambda + \mu - 1} \quad (\Re(\lambda + \mu) > 0). \quad (12)$$

2. Basic Analogue to Aleph-Function of Two Variables

We recall that AFTV [21] is an extension of the *I*-function possessing two variables [24]. Here, in the present article, we define a basic analogue to AFTV.

We record

$$G(q^a) = \left[\prod_{n=0}^{\infty} (1 - q^{a+n}) \right]^{-1} = \frac{1}{(q^a; q)_{\infty}}. \tag{13}$$

Next, we have

$$\begin{aligned} & \aleph(z_1, z_2; q) \\ &= \aleph_{P_i, Q_i, \tau_i; r; P_{i'}, Q_{i'}, \tau_{i'}; r'; P_{i''}, Q_{i''}, \tau_{i''}; r''}^{0, n_1; m_2, n_2; m_3, n_3} \left(\begin{array}{c} z_1 \\ z_2 \end{array} \middle| \begin{array}{l} (a_j; \alpha_j, A_j)_{1, n_1}, [\tau_i(a_{ji}; \alpha_{ji}, A_{ji})]_{n_1+1, P_i} \\ [\tau_i(b_{ji}; \beta_{ji}, B_{ji})]_{1, Q_i} \\ (c_j, \gamma_j)_{1, n_2}, [\tau_{i'}(c_{j i'}; \gamma_{j i'})]_{n_2+1, P_{i'}}; (e_j, E_j)_{1, n_3}, [\tau_{i''}(e_{j i''}; \gamma_{j i''})]_{n_3+1, P_{i''}} \\ (d_j, \delta_j)_{1, m_2}, [\tau_{i'}(d_{j i'}; \delta_{j i'})]_{m_2+1, Q_{i'}}; (f_j, F_j)_{1, m_3}, [\tau_{i''}(f_{j i''}; F_{j i''})]_{m_3+1, Q_{i''}} \end{array} \right) \\ &= \frac{1}{(2\pi\omega)^2} \int_{L_1} \int_{L_2} \pi^2 \phi(s_1, s_2; q) z_1^{s_1} z_2^{s_2} d_q s_1 d_q s_2, \end{aligned} \tag{14}$$

where $\omega = \sqrt{-1}$, and

$$\begin{aligned} \phi(s_1, s_2; q) &= \frac{\prod_{j=1}^{n_1} G(q^{1-a_j+\alpha_j s_1+A_j s_2})}{\sum_{i=1}^r \tau_i \left\{ \prod_{j=1}^{Q_i} G(q^{1-b_{ji}+\beta_{ji} s_1+B_{ji} s_2}) \prod_{j=n_1+1}^{P_i} G(q^{a_{ji}-\alpha_{ji} s_1-A_{ji} s_2}) \right\}} \\ &\times \frac{\prod_{j=1}^{m_2} G(q^{d_j-\delta_j s_1}) \prod_{j=1}^{n_2} G(q^{1-c_j+\gamma_j s_1})}{\sum_{i'=1}^{r'} \tau_{i'} \left\{ \prod_{j=m_2+1}^{Q_{i'}} G(q^{1-d_{j i'}+\delta_{j i'} s_1}) \prod_{j=n_2+1}^{P_{i'}} G(q^{c_{j i'}-\gamma_{j i'} s_1}) \right\}} G(q^{1-s_1}) \sin \pi s_1 \\ &\times \frac{\prod_{j=1}^{m_3} G(q^{f_j-F_j s_2}) \prod_{j=1}^{n_3} G(q^{1-e_j+E_j s_2})}{\sum_{i''=1}^{r''} \tau_{i''} \left\{ \prod_{j=m_3+1}^{Q_{i''}} G(q^{1-f_{j i''}+F_{j i''} s_2}) \prod_{j=n_3+1}^{P_{i''}} G(q^{e_{j i''}-E_{j i''} s_2}) \right\}} G(q^{1-s_2}) \sin \pi s_2, \end{aligned} \tag{15}$$

where $z_1, z_2 \neq 0$ and are real or complex. An empty product is elucidated as unity, and $P_i, P_{i'}, P_{i''}, Q_i, Q_{i'}, Q_{i''}, m_1, m_2, m_3, n_1, n_2, n_3$ are non-negative integers, such that $Q_i, Q_{i'}, Q_{i''} > 0; \tau_i, \tau_{i'}, \tau_{i''} > 0 (i = 1, \dots, r; i' = 1, \dots, r'; i'' = 1, \dots, r'')$. All the $A_s, \alpha_s, \gamma_s, \delta_s, E_s$, and F_s are presumed to be positive quantities for standardization intention, the a_s, b_s, c_s, d_s, e_s , and f_s are complex numbers. The definition of a basic analogue to AFTV will, however, make sense, even if some of these quantities are equal to zero. The contour L_1 is in the s_1 -plane and goes from $-\omega\infty$ to $+\omega\infty$, with loops where necessary, to make sure that the poles of $G(q^{d_j-\delta_j s_1}) (j = 1, \dots, m_2)$ are to the right-hand side and all the poles of $G(q^{1-a_j+\alpha_j s_1+A_j s_2}) (j = 1, \dots, n_1), G(q^{1-c_j+\gamma_j s_1}) (j = 1, \dots, n_2)$ lie to the left-hand side of L_1 . The contour L_2 is in the s_2 -plane and goes from $-\omega\infty$ to $+\omega\infty$, with loops where necessary, to ensure that the poles of $G(q^{f_j-F_j s_2}) (j = 1, \dots, m_3)$ are to the right-hand side and all the poles of $G(q^{1-a_j+\alpha_j s_1+A_j s_2}) (j = 1, \dots, n_1), G(q^{1-e_j+E_j s_2}) (j = 1, \dots, n_2)$ lie to the left-hand side of L_2 . For values of $|s_1|$ and $|s_2|$, the integrals converge, if $\Re(s_1 \log(z_1) - \log \sin \pi s_1) < 0$ and $\Re(s_2 \log(z_2) - \log \sin \pi s_2) < 0$.

3. Main Formulas

Here, we obtain fractional q -integral and q -derivative formulas concerning the basic analogue to AFTV. Here, we have the following notations:

$$A_1 = (a_j; \alpha_j, A_j)_{1, n_1}, [\tau_i(a_{ji}; \alpha_{ji}, A_{ji})]_{n_1+1, P_i}; B_1 = [\tau_i(b_{ji}; \beta_{ji}, B_{ji})]_{1, Q_i}. \tag{16}$$

$$C_1 = (c_j, \gamma_j)_{1, n_2}, [\tau_{i'}(c_{ji'}, \gamma_{ji'})]_{n_2+1, P_{i'}}; (e_j, E_j)_{1, n_3}, [\tau_{i''}(e_{ji''}, \gamma_{ji''})]_{n_3+1, P_{i''}}. \tag{17}$$

$$D_1 = (d_j, \delta_j)_{1, m_2}, [\tau_{i'}(d_{ji'}, \delta_{ji'})]_{m_2+1, Q_{i'}}; (f_j, F_j)_{1, m_3}, [\tau_{i''}(f_{ji''}, F_{ji''})]_{m_3+1, Q_{i''}}. \tag{18}$$

Theorem 1. Let $\Re(\mu) > 0$, $\rho_i \in \mathbb{Z}^+$ ($i = 1, 2$), and $|q| < 1$; then, the Riemann–Liouville fractional q -integral of (14) exists and is given as

$$I_q^\mu \left\{ x^{\lambda-1} \mathfrak{N}_{P_i, Q_i, \tau_i, r; P_{i'}, Q_{i'}, \tau_{i'}, r'; P_{i''}, Q_{i''}, \tau_{i''}, r''} \left(\begin{matrix} z_1 x^{\rho_1} \\ z_2 x^{\rho_2} \end{matrix} ; q \middle| \begin{matrix} A_1; C_1 \\ B_1; D_1 \end{matrix} \right) \right\} = (1-q)^\mu x^{\lambda+\mu-1} \\ \times \mathfrak{N}_{P_i+1, Q_i+1, \tau_i, r; P_{i'}, Q_{i'}, \tau_{i'}, r'; P_{i''}, Q_{i''}, \tau_{i''}, r''} \left(\begin{matrix} z_1 x^{\rho_1} \\ z_2 x^{\rho_2} \end{matrix} ; q \middle| \begin{matrix} (1-\lambda; \rho_1, \rho_2), A_1; C_1 \\ B_1, (1-\lambda-\mu; \rho_1, \rho_2); D_1 \end{matrix} \right), \tag{19}$$

where $\Re(s_i \log(z_i) - \log \sin \pi s_i) < 0$ ($i = 1, 2$).

Proof. We apply the definitions (8) and (14) in the left-hand side of (19), we have (say \mathcal{I})

$$\mathcal{I} = \frac{1}{\Gamma_q(\alpha)} \int_0^x (x-yq)_{\alpha-1} \left\{ y^{\lambda-1} \frac{1}{(2\pi\omega)^2} \int_{L_1} \int_{L_2} \pi^2 \phi(s_1, s_2; q) \prod_{i=1}^2 (z_i y^{\rho_i})^{s_i} d_q s_1 d_q s_2 \right\} d_q y. \tag{20}$$

By using standard calculations, we arrive at

$$\mathcal{I} = \frac{y^{\lambda-1}}{\Gamma_q(\alpha)} \frac{1}{(2\pi\omega)^2} \\ \times \int_{L_1} \int_{L_2} \pi^2 \phi(s_1, s_2; q) \prod_{i=1}^2 z_i^{s_i} \left\{ \int_0^x (x-yq)_{\alpha-1} y^{\lambda+\sum_{i=1}^2 \rho_i s_i - 1} d_q y \right\} d_q s_1 d_q s_2 \\ = \frac{1}{(2\pi\omega)^2} \int_{L_1} \int_{L_2} \pi^2 \phi(s_1, s_2; q) \prod_{i=1}^2 z_i^{s_i} I_q^\mu \left\{ x^{\lambda+\sum_{i=1}^2 \rho_i s_i - 1} \right\} d_q s_1 d_q s_2. \tag{21}$$

Next, we apply formula (12) to the equation above; then, we get

$$\mathcal{I} = \frac{1}{(2\pi\omega)^2} \int_{L_1} \int_{L_2} \pi^2 \phi(s_1, s_2; q) \prod_{i=1}^2 z_i^{s_i} \frac{\Gamma_q(\lambda + \sum_{i=1}^2 \rho_i s_i)}{\Gamma_q(\lambda + \mu + \sum_{i=1}^2 \rho_i s_i)} x^{\lambda+\mu+\sum_{i=1}^2 \rho_i s_i - 1} d_q s_1 d_q s_2. \tag{22}$$

Considering the above q -Mellin–Barnes double contour integrals in terms of the basic analogue to AFTV, we obtain (19). \square

If we use a fractional q -derivative operator without initial values, defined as

$$I_q^{-\mu} \{f(x)\} = D_{x,q}^\mu \{f(x)\} = \frac{1}{\Gamma_q(-\mu)} \int_0^x (x-tq)_{-\mu-1} f(t) d_q t, \tag{23}$$

where $\Re(\mu) < 0$; then, we yield the following result:

Theorem 2. For $\Re(\mu) > 0$, $\rho_i \in \mathbb{Z}^+$ ($i = 1, 2$), and $|q| < 1$, the Riemann–Liouville fractional q -derivative of (14) exists and is given by

$$D_{x,q}^\mu \left\{ x^{\lambda-1} \mathfrak{N}_{P_i, Q_i, \tau_i; r'; P_i', Q_i', \tau_i'; r''; P_i'', Q_i'', \tau_i''; r'''} \left(\begin{matrix} z_1 x^{\rho_1} \\ z_2 x^{\rho_2} \end{matrix} \middle| \begin{matrix} A_1; C_1 \\ B_1; D_1 \end{matrix} \right) \right\} = (1-q)^{-\mu} x^{\lambda-\mu-1} \\ \times \mathfrak{N}_{P_i+1, Q_i+1, \tau_i; r'; P_i', Q_i', \tau_i'; r''; P_i'', Q_i'', \tau_i''; r'''} \left(\begin{matrix} z_1 x^{\rho_1} \\ z_2 x^{\rho_2} \end{matrix} \middle| \begin{matrix} (1-\lambda; \rho_1, \rho_2), A_1; C_1 \\ B_1, (1-\lambda+\mu; \rho_1, \rho_2); D_1 \end{matrix} \right), \tag{24}$$

where $\Re(s_i \log(z_i) - \log \sin \pi s_i) < 0$ ($i = 1, 2$).

Proof. If we replace μ by $-\mu$ in (19), and follow the proof of Theorem 1, then we can easily obtain (24). \square

4. Leibniz's Formula

The q -expression of the Leibniz rule for the fractional q -derivatives [13] is written below

Lemma 1. For regular functions $U(x)$ and $V(x)$, we have

$$D_{x,q}^\alpha \{U(x)V(x)\} = \sum_{n=0}^\infty \frac{(-1)^n q^{\frac{n(n+1)}{2}} [q^{-\mu}; q]_n}{(q; q)_n} D_{x,q}^{\mu-n} \{U(xq^n)\} D_{x,q}^n \{V(x)\}. \tag{25}$$

Next, we have the following formula:

Theorem 3. For $\Re(\mu) < 0$, $\rho_i \in \mathbb{Z}^+$ ($i = 1, 2$), then the Riemann–Liouville fractional q -derivative of a product of two basic function is written as

$$\mathfrak{N}_{P_i+1, Q_i+1, \tau_i; r'; P_i', Q_i', \tau_i'; r''; P_i'', Q_i'', \tau_i''; r'''} \left(\begin{matrix} z_1 x^{\rho_1} \\ z_2 x^{\rho_2} \end{matrix} \middle| \begin{matrix} (1-\lambda; \rho_1, \rho_2), A_1; C_1 \\ B_1, (1-\lambda+\mu; \rho_1, \rho_2); D_1 \end{matrix} \right) \\ = \sum_{n=0}^\infty \frac{(-1)^n q^{n\lambda + \frac{n(n-1)}{2}} [q^{-\mu}; q]_n}{(q; q)_n (q^\lambda; q)_{n-\mu}} \\ \times \mathfrak{N}_{P_i+1, Q_i+1, \tau_i; r'; P_i', Q_i', \tau_i'; r''; P_i'', Q_i'', \tau_i''; r'''} \left(\begin{matrix} z_1 x^{\rho_1} \\ z_2 x^{\rho_2} \end{matrix} \middle| \begin{matrix} (0; \rho_1, \rho_2), A_1; C_1 \\ B_1, (n; \rho_1, \rho_2); D_1 \end{matrix} \right), \tag{26}$$

where $\Re(s_i \log(z_i) - \log \sin \pi s_i) < 0$ ($i = 1, 2$).

Proof. To apply the q -Leibniz rule, we take

$$U(x) = x^{\lambda-1} \text{ and } V(x) = \mathfrak{N}_{P_i, Q_i, \tau_i; r'; P_i', Q_i', \tau_i'; r''; P_i'', Q_i'', \tau_i''; r'''} \left(\begin{matrix} z_1 x^{\rho_1} \\ z_2 x^{\rho_2} \end{matrix} \middle| \begin{matrix} A_1; C_1 \\ B_1; D_1 \end{matrix} \right).$$

By using Lemma 1, we obtain the following relation:

$$D_{x,q}^\mu \left\{ x^{\lambda-1} \mathfrak{N}_{P_i, Q_i, \tau_i; r'; P_i', Q_i', \tau_i'; r''; P_i'', Q_i'', \tau_i''; r'''} \left(\begin{matrix} z_1 x^{\rho_1} \\ z_2 x^{\rho_2} \end{matrix} \middle| \begin{matrix} A_1; C_1 \\ B_1; D_1 \end{matrix} \right) \right\} \\ = \sum_{n=0}^\infty \frac{(-1)^n q^{\frac{n(n+1)}{2}} [q^{-\mu}; q]_n}{(q; q)_n} D_{x,q}^{\mu-n} (xq^n)^{\lambda-1} D_{x,q}^n \{ \mathfrak{N}(z_1 x^{\rho_1}, z_2 x^{\rho_2}; q) \}. \tag{27}$$

Next, by using Theorem 2 and setting $\lambda = 1$, we obtain

$$D_{x,q}^n \{ \aleph(z_1 x^{\rho_1}, z_2 x^{\rho_2}; q) \} = (1-q)^{-n} x^{-n} \aleph_{P_i+1, Q_i+1, \tau_i; r; P_i', Q_i', \tau_i'; r'; P_i'', Q_i'', \tau_i''; r''}^{0, n_1+1; m_2, n_2; m_3, n_3} \left(\begin{matrix} z_1 x^{\rho_1} \\ z_2 x^{\rho_2} \end{matrix} ; q \left| \begin{matrix} (0; \rho_1, \rho_2), A_1; C_1 \\ B_1, (n; \rho_1, \rho_2); D_1 \end{matrix} \right. \right). \tag{28}$$

By using the above equation and the following result:

$$D_{x,q}^\mu \{ x^{\lambda-1} \} = \frac{\Gamma_q(\lambda)}{\Gamma_q(\lambda-\mu)} x^{\lambda-\mu-1} \quad (\lambda \neq -1, -2, \dots), \tag{29}$$

We can easily obtain the desired result (26) after several algebraic manipulations. \square

5. Particular Cases

By setting $\tau_i, \tau_i', \tau_i'' \rightarrow 1$, the basic analogue to AFTV reduces to the basic analogue to the I-function of two variables [24].

Let

$$A'_1 = (a_j; \alpha_j, A_j)_{1, n_1}, (a_{ji}; \alpha_{ji}, A_{ji})_{n_1+1, P_i}; \quad B'_1 = (b_{ji}; \beta_{ji}, B_{ji})_{1, Q_i}. \tag{30}$$

$$C'_1 = (c_j, \gamma_j)_{1, n_2}, (c_{ji'}, \gamma_{ji'})_{n_2+1, P_i'}; (e_j, E_j)_{1, n_3}, (e_{ji}'', \gamma_{ji}'')_{n_3+1, P_i''}. \tag{31}$$

$$D'_1 = (d_j, \delta_j)_{1, m_2}, (d_{ji}', \delta_{ji}')_{m_2+1, Q_i'}; (f_j, F_j)_{1, m_3}, (f_{ji}'', F_{ji}'')_{m_3+1, Q_i''}. \tag{32}$$

We have the following result:

Corollary 1.

$$I_{P_i+1, Q_i+1, r; P_i', Q_i', r'; P_i'', Q_i'', r''}^{0, n_1+1; m_2, n_2; m_3, n_3} \left(\begin{matrix} z_1 x^{\rho_1} \\ z_2 x^{\rho_2} \end{matrix} ; q \left| \begin{matrix} (1-\lambda; \rho_1, \rho_2), A'_1; C'_1 \\ B'_1, (1-\lambda+\mu; \rho_1, \rho_2); D'_1 \end{matrix} \right. \right) = \sum_{n=0}^{\infty} \frac{(-1)^n q^{n\lambda + \frac{n(n-1)}{2}} [q^{-\mu}; q]_n}{(q; q)_n (q^\lambda; q)_{n-\mu}} \times I_{P_i+1, Q_i+1, r; P_i', Q_i', r'; P_i'', Q_i'', r''}^{0, n_1+1; m_2, n_2; m_3, n_3} \left(\begin{matrix} z_1 x^{\rho_1} \\ z_2 x^{\rho_2} \end{matrix} ; q \left| \begin{matrix} (0; \rho_1, \rho_2), A'_1; C'_1 \\ B'_1, (n; \rho_1, \rho_2); D'_1 \end{matrix} \right. \right), \tag{33}$$

where $\Re(s_i \log(z_i) - \log \sin \pi s_i) < 0 \quad (i = 1, 2)$.

Proof. By setting $\tau_i, \tau_i', \tau_i'' \rightarrow 1$ and following the proof of Theorem 3, we can easily obtain the desired result (33). \square

Remark 1. If the basic analogue to the I-function of two variables reduces to the basic analogue to the H-function of two variables [25], then we can obtain the result due to Yadav et al. [18].

The basic analogue to AFTV reduces to the basic analogue to AFTV, defined by Ahmad et al. [26].

Let

$$A = (a_j, A_j)_{1, n'} \cdots, [\tau_i(a_{ji}, A_{ji})]_{n+1, P_i}. \tag{34}$$

$$B = (b_j, B_j)_{1, m'} \cdots, [\tau_i(b_{ji}, B_{ji})]_{m+1, Q_i}. \tag{35}$$

Then, we have following relation:

Corollary 2.

$$\begin{aligned} & \mathfrak{N}_{p_i+1, q_i+1, \tau_i; r}^{m, n+1} \left(zx^\rho; q \left| \begin{array}{l} (1-\lambda; \rho), A \\ B, (1-\lambda+\mu; \rho) \end{array} \right. \right) \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n q^{n\lambda + \frac{n(n-1)}{2}} [q^{-\mu}; q]_n}{(q; q)_n (q^\lambda; q)_{n-\mu}} \mathfrak{N}_{p_i+1, q_i+1, \tau_i; r}^{m, n+1} \left(zx^\rho; q \left| \begin{array}{l} (0; \rho), A \\ B, (n; \rho) \end{array} \right. \right). \end{aligned} \quad (36)$$

If we set $\tau_i \rightarrow 1$ in (36), then the basic analogue to AFTV reduces to the basic analogue to the I -function of one variable. We have

Corollary 3.

$$\begin{aligned} & I_{p_i+1, q_i+1; r}^{m, n+1} \left(zx^\rho; q \left| \begin{array}{l} (1-\lambda; \rho), (a_j, A_j)_{1, n'}, \dots, (a_{j_i}, A_{j_i})_{n+1, p_i} \\ (b_j, B_j)_{1, m'}, \dots, (b_{j_i}, B_{j_i})_{m+1, q_i}, (1-\lambda+\mu; \rho) \end{array} \right. \right) \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n q^{n\lambda + \frac{n(n-1)}{2}} [q^{-\mu}; q]_n}{(q; q)_n} \\ &\times I_{p_i+1, q_i+1; r}^{m, n+1} \left(zx^\rho; q \left| \begin{array}{l} (0; \rho), (a_j, A_j)_{1, n'}, \dots, (a_{j_i}, A_{j_i})_{n+1, p_i} \\ (b_j, B_j)_{1, m'}, \dots, (b_{j_i}, B_{j_i})_{m+1, q_i}, (n; \rho) \end{array} \right. \right). \end{aligned} \quad (37)$$

Remark 2. If the basic analogue to AFTV reduces to the basic analogue to the H -function of one variable (see [27]), then we can report a similar expression.

Remark 3. We can generalize the q -extension of the Leibniz rule for the basic analogue to special multivariable functions; by this, we can obtain similar formulas by using similar methods.

6. Conclusions

After the famous letter between L'Hopital and Leibniz from 1695, using integral transformations, we obtained a new field in mathematics, called fractional calculus. Among other things, there are fractional derivative and fractional integrals, as well as fractional differential equations. It is also well-known that fractional calculus operators and their basic (or q -) analogues have many applications, such as signal processing, bio-medical engineering, control systems, radars, sonars, to solve dual integral and series equations in elasticity, etc. In this article, we have proposed the fractional-order q -integrals and q -derivatives involving a basic analogue to AFTV [11,12,26,28]. Some remarkable applications of these integrals and derivatives have also been discussed. By specializing the various parameters as well as the variables in the basic analogue to AFTV, we can obtain a large number of q -extensions of the Leibniz rule, involving a large set of basic functions, that is, the product of such basic functions, which are describable in terms of the basic analogue to the H -function [25,27], the basic analogue to Meijer's G -function [27], the basic analogue to MacRobert's E -function [29], and the basic analogue to the hypergeometric function [10,16–18].

Author Contributions: Conceptualization, D.K.; Data curation, D.B., F.A. and N.S.; Formal analysis, D.K., D.B., F.A. and N.S.; Funding acquisition, D.B.; Investigation, D.K. and F.A.; Methodology, D.K., D.B. and F.A.; Resources, F.A. and N.S.; Supervision, D.B. and N.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The author (D.K.) would like to thank the Agriculture University Jodhpur for supporting and encouraging this work.

Conflicts of Interest: All authors declare that they have no conflict of interest.

References

- Kumar, D.; Choi, J. Generalized fractional kinetic equations associated with Aleph function. *Proc. Jangjeon Math. Soc.* **2016**, *19*, 145–155.
- Kumar, D.; Gupta, R.K.; Shaktawat, B.S.; Choi, J. Generalized fractional calculus formulas involving the product of Aleph-function and Srivastava polynomials. *Proc. Jangjeon Math. Soc.* **2017**, *20*, 701–717.
- Kumar, D.; Ram, J.; Choi, J. Certain generalized integral formulas involving Chebyshev Hermite polynomials, generalized M -series and Aleph-function, and their application in heat conduction. *Int. J. Math. Anal.* **2015**, *9*, 1795–1803. [[CrossRef](#)]
- Ram, J.; Kumar, D. Generalized fractional integration of the \aleph -function. *J. Raj. Acad. Phy. Sci.* **2011**, *10*, 373–382.
- Samko, G.; Kilbas, A.A.; Marichev, O.I. *Fractional Integrals and Derivatives: Theory and Applications*; Gordon and Breach Science Publishers: New York, NY, USA, 1993.
- Südland, N.; Volkmann, J.; Kumar, D. Applications to give an analytical solution to the Black Scholes equation. *Integral Transform. Spec. Funct.* **2019**, *30*, 205–230. [[CrossRef](#)]
- Exton, H. q -hypergeometric functions and applications. *Ellis Horwood Series: Mathematics and its Applications*; Ellis Horwood Ltd.: Chichester, UK, 1983; 347p.
- Gasper, G.; Rahman, M. *Basic Hypergeometric Series*; Cambridge University Press: Cambridge, UK, 1990.
- Rajkovic, P.M.; Marinkovic, S.D.; Stankovic, M.S. Fractional integrals and derivatives in q -calculus. *Appl. Anal. Discrete Math.* **2007**, *1*, 311–323.
- Slater, L.J. *Generalized Hypergeometric Functions*; Cambridge University Press: Cambridge, UK, 1966.
- Al-Salam, W.A. q -analogues of Cauchy's formula. *Proc. Am. Math. Soc.* **1952–1953**, *17*, 182–184.
- Al-Salam, W.A. Some fractional q -integrals and q -derivatives. *Proc. Edinburgh Math. Soc.* **1966**, *15*, 135–140. [[CrossRef](#)]
- Agarwal, R.P. Certain fractional q -integrals and q -derivatives. *Proc. Camb. Phil. Soc.* **1969**, *66*, 365–370. [[CrossRef](#)]
- Kumar, D.; Ayant, F.Y.; Tariboon, J. On transformation involving basic analogue of multivariable H -function. *J. Funct. Spaces* **2020**, *2020*, 2616043. [[CrossRef](#)]
- Sahni, N.; Kumar, D.; Ayant, F.Y.; Singh, S. A transformation involving basic multivariable I -function of Prathima. *J. Ramanujan Soc. Math. Math. Sci.* **2021**, *8*, 95–108.
- Yadav, R.K.; Purohit, S.D. On applications of Weyl fractional q -integral operator to generalized basic hypergeometric functions. *Kyungpook Math. J.* **2006**, *46*, 235–245.
- Yadav, R.K.; Purohit, S.D.; Kalla, S.L.; Vyas, V.K. Certain fractional q -integral formulae for the generalized basic hypergeometric functions of two variables. *J. Inequal. Spec. Funct.* **2010**, *1*, 30–38.
- Yadav, R.K.; Purohit, S.D.; Vyas, V.K. On transformations involving generalized basic hypergeometric function of two variables. *Rev. Téc. Ing. Univ. Zulia* **2010**, *33*, 176–182.
- Purohit, S.D.; Yadav, R.K. On generalized fractional q -integral operators involving the q -gauss hypergeometric function. *Bull. Math. Anal. Appl.* **2010**, *2*, 35–44.
- Galué, L. Generalized Erdélyi-Kober fractional q -integral operator. *Kuwait J. Sci. Eng.* **2009**, *36*, 21–34.
- Kumar, D. Generalized fractional differintegral operators of the Aleph-function of two variables. *J. Chem. Biol. Phys. Sci. Sec. C* **2016**, *6*, 1116–1131.
- Jia, Z.; Khan, B.; Agarwal, P.; Hu, Q.; Wang, X. Two new Bailey Lattices and their applications. *Symmetry* **2021**, *13*, 958. [[CrossRef](#)]
- Jia, Z.; Khan, B.; Hu, Q.; Niu, D. Applications of generalized q -difference equations for general q -polynomials. *Symmetry* **2021**, *13*, 1222. [[CrossRef](#)]
- Sharma, C.K.; Mishra, P.L. On the I -function of two variables and its certain properties. *Acta Ciencia Indica* **1991**, *17*, 1–4.
- Saxena, R.K.; Modi, G.C.; Kalla, S.L. A basic analogue of H -function of two variable. *Rev. Tec. Ing. Univ. Zulia* **1987**, *10*, 35–39.
- Ahmad, A.; Jain, R.; Jain, D.K. q -analogue of Aleph-function and its transformation formulae with q -derivative. *J. Stat. Appl. Pro.* **2017**, *6*, 567–575. [[CrossRef](#)]
- Saxena, R.K.; Modi, G.C.; Kalla, S.L. A basic analogue of Fox's H -function. *Rev. Tec. Ing. Univ. Zulia* **1983**, *6*, 139–143.
- Ayant, F.Y.; Kumar, D. Certain finite double integrals involving the hypergeometric function and Aleph-function. *Int. J. Math. Trends Technol.* **2016**, *35*, 49–55. [[CrossRef](#)]
- Agarwal, R. A basic analogue of MacRobert's E -function. *Glasg. Math. J.* **1961**, *5*, 4–7. [[CrossRef](#)]