

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/309187612>

# A fractional finite difference inclusion

Data · October 2016

CITATIONS

0

READS

207

## 3 authors:



**Dumitru Baleanu**

Institute of Space Sciences

**1,006** PUBLICATIONS **16,275** CITATIONS

[SEE PROFILE](#)



**Shahram Rezapour**

Azerbaijan Shahid Madani University

**138** PUBLICATIONS **2,240** CITATIONS

[SEE PROFILE](#)



**Saeid Salehi**

Azarbaijan Shahid Madani University

**17** PUBLICATIONS **72** CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Approximate solution of Fractional Differential equations [View project](#)



Fractional-Order Control of Time-Delay Systems [View project](#)

Volume 20, Number 5  
ISSN:1521-1398 PRINT,1572-9206 ONLINE

May 2016



**Journal of  
Computational  
Analysis and  
Applications**

**EUDOXUS PRESS,LLC**

**Journal of Computational Analysis and Applications**

ISSNno.'s:1521-1398 PRINT,1572-9206 ONLINE

**SCOPE OF THE JOURNAL**

**An international publication of Eudoxus Press, LLC  
(fourteen times annually)**

**Editor in Chief: George Anastassiou**

**Department of Mathematical Sciences,**

**University of Memphis, Memphis, TN 38152-3240, U.S.A**

**ganastss@memphis.edu**

**<http://www.msci.memphis.edu/~ganastss/jocaaa>**

The main purpose of "J.Computational Analysis and Applications" is to publish high quality research articles from all subareas of Computational Mathematical Analysis and its many potential applications and connections to other areas of Mathematical Sciences. Any paper whose approach and proofs are computational, using methods from Mathematical Analysis in the broadest sense is suitable and welcome for consideration in our journal, except from Applied Numerical Analysis articles. Also plain word articles without formulas and proofs are excluded. The list of possibly connected mathematical areas with this publication includes, but is not restricted to: Applied Analysis, Applied Functional Analysis, Approximation Theory, Asymptotic Analysis, Difference Equations, Differential Equations, Partial Differential Equations, Fourier Analysis, Fractals, Fuzzy Sets, Harmonic Analysis, Inequalities, Integral Equations, Measure Theory, Moment Theory, Neural Networks, Numerical Functional Analysis, Potential Theory, Probability Theory, Real and Complex Analysis, Signal Analysis, Special Functions, Splines, Stochastic Analysis, Stochastic Processes, Summability, Tomography, Wavelets, any combination of the above, e.t.c.

"J.Computational Analysis and Applications" is a peer-reviewed Journal. See the instructions for preparation and submission

of articles to JoCAAA. Assistant to the Editor:

Dr.Razvan Mezei, Lenoir-Rhyne University, Hickory, NC 28601, USA.

**Journal of Computational Analysis and Applications(JoCAAA)** is published by **EUDOXUS PRESS,LLC**,1424 Beaver Trail

Drive,Cordova,TN38016,USA,[anastassioug@yahoo.com](mailto:anastassioug@yahoo.com)

<http://www.eudoxuspress.com>. **Annual Subscription Prices:**For USA and

Canada,Institutional:Print \$700, Electronic OPEN ACCESS. Individual:Print \$350. For any other part of the world add \$130 more(handling and postages) to the above prices for Print. No credit card payments.

**Copyright**©2016 by Eudoxus Press,LLC,all rights reserved.JoCAAA is printed in USA.

**JoCAAA is reviewed and abstracted by AMS Mathematical Reviews,MATHSCI,and Zentralblat MATH.**

It is strictly prohibited the reproduction and transmission of any part of JoCAAA and in any form and by any means without the written permission of the publisher.It is only allowed to educators to Xerox articles for educational purposes.The publisher assumes no responsibility for the content of published papers.

---

## Editorial Board

### Associate Editors of Journal of Computational Analysis and Applications

---

**Francesco Altomare**

Dipartimento di Matematica  
Universita' di Bari  
Via E.Orabona, 4  
70125 Bari, ITALY  
Tel+39-080-5442690 office  
+39-080-3944046 home  
+39-080-5963612 Fax  
altomare@dm.uniba.it  
Approximation Theory, Functional  
Analysis, Semigroups and Partial  
Differential Equations, Positive  
Operators.

**Ravi P. Agarwal**

Department of Mathematics  
Texas A&M University - Kingsville  
700 University Blvd.  
Kingsville, TX 78363-8202  
tel: 361-593-2600  
Agarwal@tamuk.edu  
Differential Equations, Difference  
Equations, Inequalities

**George A. Anastassiou**

Department of Mathematical Sciences  
The University of Memphis  
Memphis, TN 38152, U.S.A  
Tel. 901-678-3144  
e-mail: ganastss@memphis.edu  
Approximation Theory, Real  
Analysis,  
Wavelets, Neural Networks,  
Probability, Inequalities.

**J. Marshall Ash**

Department of Mathematics  
De Paul University  
2219 North Kenmore Ave.  
Chicago, IL 60614-3504  
773-325-4216  
e-mail: mash@math.depaul.edu  
Real and Harmonic Analysis

**Dumitru Baleanu**

Department of Mathematics and  
Computer Sciences,  
Cankaya University, Faculty of Art  
and Sciences,  
06530 Balgat, Ankara,  
Turkey, dumitru@cankaya.edu.tr

Fractional Differential Equations  
Nonlinear Analysis, Fractional  
Dynamics

**Carlo Bardaro**

Dipartimento di Matematica e  
Informatica  
Universita di Perugia  
Via Vanvitelli 1  
06123 Perugia, ITALY  
TEL+390755853822  
+390755855034  
FAX+390755855024  
E-mail carlo.bardaro@unipg.it  
Web site:  
<http://www.unipg.it/~bardaro/>  
Functional Analysis and  
Approximation Theory, Signal  
Analysis, Measure Theory, Real  
Analysis.

**Martin Bohner**

Department of Mathematics and  
Statistics, Missouri S&T  
Rolla, MO 65409-0020, USA  
bohner@mst.edu  
web.mst.edu/~bohner  
Difference equations, differential  
equations, dynamic equations on  
time scale, applications in  
economics, finance, biology.

**Jerry L. Bona**

Department of Mathematics  
The University of Illinois at  
Chicago  
851 S. Morgan St. CS 249  
Chicago, IL 60601  
e-mail: bona@math.uic.edu  
Partial Differential Equations,  
Fluid Dynamics

**Luis A. Caffarelli**

Department of Mathematics  
The University of Texas at Austin  
Austin, Texas 78712-1082  
512-471-3160  
e-mail: caffarel@math.utexas.edu  
Partial Differential Equations

**George Cybenko**

Thayer School of Engineering

Dartmouth College  
8000 Cummings Hall,  
Hanover, NH 03755-8000  
603-646-3843 (X 3546 Secr.)  
e-mail:george.cybenko@dartmouth.edu  
Approximation Theory and Neural  
Networks

**Sever S. Dragomir**

School of Computer Science and  
Mathematics, Victoria University,  
PO Box 14428,  
Melbourne City,  
MC 8001, AUSTRALIA  
Tel. +61 3 9688 4437  
Fax +61 3 9688 4050  
sever.dragomir@vu.edu.au  
Inequalities, Functional Analysis,  
Numerical Analysis, Approximations,  
Information Theory, Stochastics.

**Oktay Duman**

TOBB University of Economics and  
Technology,  
Department of Mathematics, TR-  
06530,  
Ankara, Turkey,  
oduman@etu.edu.tr  
Classical Approximation Theory,  
Summability Theory, Statistical  
Convergence and its Applications

**Saber N. Elaydi**

Department Of Mathematics  
Trinity University  
715 Stadium Dr.  
San Antonio, TX 78212-7200  
210-736-8246  
e-mail: selaydi@trinity.edu  
Ordinary Differential Equations,  
Difference Equations

**Christodoulos A. Floudas**

Department of Chemical Engineering  
Princeton University  
Princeton, NJ 08544-5263  
609-258-4595(x4619 assistant)  
e-mail: floudas@titan.princeton.edu  
Optimization Theory&Applications,  
Global Optimization

**J .A. Goldstein**

Department of Mathematical Sciences  
The University of Memphis  
Memphis, TN 38152  
901-678-3130  
jgoldste@memphis.edu

Partial Differential Equations,  
Semigroups of Operators

**H. H. Gonska**

Department of Mathematics  
University of Duisburg  
Duisburg, D-47048  
Germany  
011-49-203-379-3542  
e-mail: heiner.gonska@uni-due.de  
Approximation Theory, Computer  
Aided Geometric Design

**John R. Graef**

Department of Mathematics  
University of Tennessee at  
Chattanooga  
Chattanooga, TN 37304 USA  
John-Graef@utc.edu  
Ordinary and functional  
differential equations, difference  
equations, impulsive systems,  
differential inclusions, dynamic  
equations on time scales, control  
theory and their applications

**Weimin Han**

Department of Mathematics  
University of Iowa  
Iowa City, IA 52242-1419  
319-335-0770  
e-mail: whan@math.uiowa.edu  
Numerical analysis, Finite element  
method, Numerical PDE, Variational  
inequalities, Computational  
mechanics

**Tian-Xiao He**

Department of Mathematics and  
Computer Science  
P.O. Box 2900, Illinois Wesleyan  
University  
Bloomington, IL 61702-2900, USA  
Tel (309)556-3089  
Fax (309)556-3864  
the@iwu.edu  
Approximations, Wavelet,  
Integration Theory, Numerical  
Analysis, Analytic Combinatorics

**Margareta Heilmann**

Faculty of Mathematics and Natural  
Sciences, University of Wuppertal  
Gaußstraße 20  
D-42119 Wuppertal, Germany,  
heilmann@math.uni-wuppertal.de

Approximation Theory (Positive  
Linear Operators)

**Xing-Biao Hu**

Institute of Computational  
Mathematics  
AMSS, Chinese Academy of Sciences  
Beijing, 100190, CHINA  
hxb@lsec.cc.ac.cn  
Computational Mathematics

**Jong Kyu Kim**

Department of Mathematics  
Kyungnam University  
Masan Kyungnam, 631-701, Korea  
Tel 82-(55)-249-2211  
Fax 82-(55)-243-8609  
jongkyuk@kyungnam.ac.kr  
Nonlinear Functional Analysis,  
Variational Inequalities, Nonlinear  
Ergodic Theory, ODE, PDE,  
Functional Equations.

**Robert Kozma**

Department of Mathematical Sciences  
The University of Memphis  
Memphis, TN 38152, USA  
rkozma@memphis.edu  
Neural Networks, Reproducing Kernel  
Hilbert Spaces,  
Neural Percolation Theory

**Mustafa Kulenovic**

Department of Mathematics  
University of Rhode Island  
Kingston, RI 02881, USA  
kulenm@math.uri.edu  
Differential and Difference  
Equations

**Irena Lasiecka**

Department of Mathematical Sciences  
University of Memphis  
Memphis, TN 38152  
PDE, Control Theory, Functional  
Analysis, lasiecka@memphis.edu

**Burkhard Lenze**

Fachbereich Informatik  
Fachhochschule Dortmund  
University of Applied Sciences  
Postfach 105018  
D-44047 Dortmund, Germany  
e-mail: lenze@fh-dortmund.de  
Real Networks, Fourier Analysis,  
Approximation Theory

**Hrushikesh N. Mhaskar**

Department Of Mathematics  
California State University  
Los Angeles, CA 90032  
626-914-7002  
e-mail: hmhaska@gmail.com  
Orthogonal Polynomials,  
Approximation Theory, Splines,  
Wavelets, Neural Networks

**Ram N. Mohapatra**

Department of Mathematics  
University of Central Florida  
Orlando, FL 32816-1364  
tel.407-823-5080  
ram.mohapatra@ucf.edu  
Real and Complex Analysis,  
Approximation Th., Fourier  
Analysis, Fuzzy Sets and Systems

**Gaston M. N'Guerekata**

Department of Mathematics  
Morgan State University  
Baltimore, MD 21251, USA  
tel: 1-443-885-4373  
Fax 1-443-885-8216  
Gaston.N'Guerekata@morgan.edu  
nguerekata@aol.com  
Nonlinear Evolution Equations,  
Abstract Harmonic Analysis,  
Fractional Differential Equations,  
Almost Periodicity & Almost  
Automorphy

**M.Zuhair Nashed**

Department Of Mathematics  
University of Central Florida  
PO Box 161364  
Orlando, FL 32816-1364  
e-mail: znashed@mail.ucf.edu  
Inverse and Ill-Posed problems,  
Numerical Functional Analysis,  
Integral Equations, Optimization,  
Signal Analysis

**Mubenga N. Nkashama**

Department OF Mathematics  
University of Alabama at Birmingham  
Birmingham, AL 35294-1170  
205-934-2154  
e-mail: nkashama@math.uab.edu  
Ordinary Differential Equations,  
Partial Differential Equations

**Vassilis Papanicolaou**

Department of Mathematics

National Technical University of Athens  
Zografou campus, 157 80  
Athens, Greece  
tel:: +30(210) 772 1722  
Fax +30(210) 772 1775  
papanico@math.ntua.gr  
Partial Differential Equations,  
Probability

**Choonkil Park**

Department of Mathematics  
Hanyang University  
Seoul 133-791  
S. Korea, baak@hanyang.ac.kr  
Functional Equations

**Svetlozar (Zari) Rachev**,  
Professor of Finance, College of  
Business, and Director of  
Quantitative Finance Program,  
Department of Applied Mathematics &  
Statistics  
Stonybrook University  
312 Harriman Hall, Stony Brook, NY  
11794-3775  
tel: +1-631-632-1998,  
svetlozar.rachev@stonybrook.edu

**Alexander G. Ramm**

Mathematics Department  
Kansas State University  
Manhattan, KS 66506-2602  
e-mail: ramm@math.ksu.edu  
Inverse and Ill-posed Problems,  
Scattering Theory, Operator Theory,  
Theoretical Numerical Analysis,  
Wave Propagation, Signal Processing  
and Tomography

**Tomasz Rychlik**

Polish Academy of Sciences  
Instytut Matematyczny PAN  
00-956 Warszawa, skr. poczt. 21  
ul. Śniadeckich 8  
Poland  
trychlik@impan.pl  
Mathematical Statistics,  
Probabilistic Inequalities

**Boris Shekhtman**

Department of Mathematics  
University of South Florida  
Tampa, FL 33620, USA  
Tel 813-974-9710  
shekhtma@usf.edu

Approximation Theory, Banach  
spaces, Classical Analysis

**T. E. Simos**

Department of Computer  
Science and Technology  
Faculty of Sciences and Technology  
University of Peloponnese  
GR-221 00 Tripolis, Greece  
Postal Address:  
26 Menelaou St.  
Anfithea - Paleon Faliron  
GR-175 64 Athens, Greece  
tsimos@mail.ariadne-t.gr  
Numerical Analysis

**H. M. Srivastava**

Department of Mathematics and  
Statistics  
University of Victoria  
Victoria, British Columbia V8W 3R4  
Canada  
tel.250-472-5313; office,250-477-  
6960 home, fax 250-721-8962  
harimsri@math.uvic.ca  
Real and Complex Analysis,  
Fractional Calculus and Appl.,  
Integral Equations and Transforms,  
Higher Transcendental Functions and  
Appl., q-Series and q-Polynomials,  
Analytic Number Th.

**I. P. Stavroulakis**

Department of Mathematics  
University of Ioannina  
451-10 Ioannina, Greece  
ipstav@cc.uoi.gr  
Differential Equations  
Phone +3-065-109-8283

**Manfred Tasche**

Department of Mathematics  
University of Rostock  
D-18051 Rostock, Germany  
manfred.tasche@mathematik.uni-  
rostock.de  
Numerical Fourier Analysis, Fourier  
Analysis, Harmonic Analysis, Signal  
Analysis, Spectral Methods,  
Wavelets, Splines, Approximation  
Theory

**Roberto Triggiani**

Department of Mathematical Sciences  
University of Memphis  
Memphis, TN 38152  
PDE, Control Theory, Functional

Analysis, rtrggani@memphis.edu

**Juan J. Trujillo**

University of La Laguna  
Departamento de Analisis Matematico  
C/Astr.Fco.Sanchez s/n  
38271. LaLaguna. Tenerife.  
SPAIN  
Tel/Fax 34-922-318209  
Juan.Trujillo@ull.es  
Fractional: Differential Equations-  
Operators-Fourier Transforms,  
Special functions, Approximations,  
and Applications

**Ram Verma**

International Publications  
1200 Dallas Drive #824 Denton,  
TX 76205, USA  
[Verma99@msn.com](mailto:Verma99@msn.com)  
Applied Nonlinear Analysis,  
Numerical Analysis, Variational  
Inequalities, Optimization Theory,  
Computational Mathematics, Operator  
Theory

**Xiang Ming Yu**

Department of Mathematical Sciences  
Southwest Missouri State University  
Springfield, MO 65804-0094  
417-836-5931  
xmy944f@missouristate.edu  
Classical Approximation Theory,  
Wavelets

**Lotfi A. Zadeh**

Professor in the Graduate School  
and Director, Computer Initiative,  
Soft Computing (BISC)  
Computer Science Division  
University of California at  
Berkeley  
Berkeley, CA 94720  
Office: 510-642-4959  
Sec: 510-642-8271  
Home: 510-526-2569  
FAX: 510-642-1712  
zadeh@cs.berkeley.edu  
Fuzzyness, Artificial Intelligence,  
Natural language processing, Fuzzy  
logic

**Richard A. Zalik**

Department of Mathematics  
Auburn University  
Auburn University, AL 36849-5310

USA.

Tel 334-844-6557 office  
678-642-8703 home  
Fax 334-844-6555  
zalik@auburn.edu  
Approximation Theory, Chebychev  
Systems, Wavelet Theory

**Ahmed I. Zayed**

Department of Mathematical Sciences  
DePaul University  
2320 N. Kenmore Ave.  
Chicago, IL 60614-3250  
773-325-7808  
e-mail: azayed@condor.depaul.edu  
Shannon sampling theory, Harmonic  
analysis and wavelets, Special  
functions and orthogonal  
polynomials, Integral transforms

**Ding-Xuan Zhou**

Department Of Mathematics  
City University of Hong Kong  
83 Tat Chee Avenue  
Kowloon, Hong Kong  
852-2788 9708, Fax: 852-2788 8561  
e-mail: mazhou@cityu.edu.hk  
Approximation Theory, Spline  
functions, Wavelets

**Xin-long Zhou**

Fachbereich Mathematik, Fachgebiet  
Informatik  
Gerhard-Mercator-Universitat  
Duisburg  
Lotharstr.65, D-47048 Duisburg,  
Germany  
e-mail: Xzhou@informatik.uni-  
duisburg.de  
Fourier Analysis, Computer-Aided  
Geometric Design, Computational  
Complexity, Multivariate  
Approximation Theory, Approximation  
and Interpolation Theory



---

**Instructions to Contributors**  
**Journal of Computational Analysis and Applications**  
An international publication of Eudoxus Press, LLC, of TN.

**Editor in Chief: George Anastassiou**  
Department of Mathematical Sciences  
University of Memphis  
Memphis, TN 38152-3240, U.S.A.

**1. Manuscripts files in Latex and PDF and in English, should be submitted via email to the Editor-in-Chief:**

**Prof. George A. Anastassiou**  
Department of Mathematical Sciences  
The University of Memphis  
Memphis, TN 38152, USA.  
Tel. 901.678.3144  
e-mail: [ganastss@memphis.edu](mailto:ganastss@memphis.edu)

**Authors may want to recommend an associate editor the most related to the submission to possibly handle it.**

**Also authors may want to submit a list of six possible referees, to be used in case we cannot find related referees by ourselves.**

**2. Manuscripts should be typed using any of TEX, LaTeX, AMS-TEX, or AMS-LaTeX and according to EUDOXUS PRESS, LLC. LATEX STYLE FILE. (Click [HERE](#) to save a copy of the style file.) They should be carefully prepared in all respects. Submitted articles should be brightly typed (not dot-matrix), double spaced, in ten point type size and in 8(1/2)x11 inch area per page. Manuscripts should have generous margins on all sides and should not exceed 24 pages.**

**3. Submission is a representation that the manuscript has not been published previously in this or any other similar form and is not currently under consideration for publication elsewhere. A statement transferring from the authors (or their employers, if they hold the copyright) to Eudoxus Press, LLC, will be required before the manuscript can be accepted for publication. The Editor-in-Chief will supply the necessary forms for this transfer. Such a written transfer of copyright, which previously was assumed to be implicit in the act of submitting a manuscript, is necessary under the U.S. Copyright Law in order for the publisher to carry through the dissemination of research results and reviews as widely and effectively as possible.**

**4. The paper starts with the title of the article, author's name(s) (no titles or degrees), author's affiliation(s) and e-mail addresses. The affiliation should comprise the department, institution (usually university or company), city, state (and/or nation) and mail code.**

**The following items, 5 and 6, should be on page no. 1 of the paper.**

**5. An abstract is to be provided, preferably no longer than 150 words.**

**6. A list of 5 key words is to be provided directly below the abstract. Key words should express the precise content of the manuscript, as they are used for indexing purposes.**

**The main body of the paper should begin on page no. 1, if possible.**

**7. All sections should be numbered with Arabic numerals (such as: 1. INTRODUCTION) .**

**Subsections should be identified with section and subsection numbers (such as 6.1. Second-Value Subheading).**

**If applicable, an independent single-number system (one for each category) should be used to label all theorems, lemmas, propositions, corollaries, definitions, remarks, examples, etc. The label (such as Lemma 7) should be typed with paragraph indentation, followed by a period and the lemma itself.**

**8. Mathematical notation must be typeset. Equations should be numbered consecutively with Arabic numerals in parentheses placed flush right, and should be thusly referred to in the text [such as Eqs.(2) and (5)]. The running title must be placed at the top of even numbered pages and the first author's name, et al., must be placed at the top of the odd numbered pages.**

**9. Illustrations (photographs, drawings, diagrams, and charts) are to be numbered in one consecutive series of Arabic numerals. The captions for illustrations should be typed double space. All illustrations, charts, tables, etc., must be embedded in the body of the manuscript in proper, final, print position. In particular, manuscript, source, and PDF file version must be at camera ready stage for publication or they cannot be considered.**

**Tables are to be numbered (with Roman numerals) and referred to by number in the text. Center the title above the table, and type explanatory footnotes (indicated by superscript lowercase letters) below the table.**

**10. List references alphabetically at the end of the paper and number them consecutively. Each must be cited in the text by the appropriate Arabic numeral in square brackets on the baseline.**

**References should include (in the following order):  
initials of first and middle name, last name of author(s)  
title of article,**

name of publication, volume number, inclusive pages, and year of publication.

Authors should follow these examples:

**Journal Article**

1. H.H.Gonska, Degree of simultaneous approximation of bivariate functions by Gordon operators, (journal name in italics) *J. Approx. Theory*, 62,170-191(1990).

**Book**

2. G.G.Lorentz, (title of book in italics) *Bernstein Polynomials* (2nd ed.), Chelsea, New York, 1986.

**Contribution to a Book**

3. M.K.Khan, Approximation properties of beta operators, in (title of book in italics) *Progress in Approximation Theory* (P.Nevai and A.Pinkus, eds.), Academic Press, New York, 1991, pp.483-495.

**11. All acknowledgements (including those for a grant and financial support) should occur in one paragraph that directly precedes the References section.**

**12. Footnotes should be avoided. When their use is absolutely necessary, footnotes should be numbered consecutively using Arabic numerals and should be typed at the bottom of the page to which they refer. Place a line above the footnote, so that it is set off from the text. Use the appropriate superscript numeral for citation in the text.**

**13. After each revision is made please again submit via email Latex and PDF files of the revised manuscript, including the final one.**

**14. Effective 1 Nov. 2009 for current journal page charges, contact the Editor in Chief. Upon acceptance of the paper an invoice will be sent to the contact author. The fee payment will be due one month from the invoice date. The article will proceed to publication only after the fee is paid. The charges are to be sent, by money order or certified check, in US dollars, payable to Eudoxus Press, LLC, to the address shown on the Eudoxus [homepage](#).**

**No galleys will be sent and the contact author will receive one (1) electronic copy of the journal issue in which the article appears.**

**15. This journal will consider for publication only papers that contain proofs for their listed results.**

# Barnes-type Peters of the first kind and poly-Cauchy of the first kind mixed-type polynomials

Dae San Kim

Department of Mathematics, Sogang University  
Seoul 121-742, Republic of Korea  
dskim@sogang.ac.kr

Taekyun Kim

Department of Mathematics, Kwangwoon University  
Seoul 139-701, Republic of Korea  
tkkim@kw.ac.kr

Takao Komatsu

Graduate School of Science and Technology, Hirosaki University  
Hirosaki 036-8561, Japan  
komatsu@cc.hirosaki-u.ac.jp

Dmitry V. Dolgy

Institute of Natural Sciences, Far Eastern Federal University  
690950 Vladivostok, Russia  
d\_dol@mail.ru

MR Subject Classifications: 05A15, 05A40, 11B68, 11B75, 65Q05

## Abstract

In this paper, by considering Barnes-type Peters polynomials of the first kind as well as poly-Cauchy polynomials of the first kind, we define and investigate the mixed-type polynomials of these polynomials. From the properties of Sheffer sequences of these polynomials arising from umbral calculus, we derive new and interesting identities.

## 1 Introduction

In this paper, we consider the polynomials

$$s_n^{(k)}(x) = s_n^{(k)}(x|\lambda; \mu) = s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r)$$

called the Barnes-type Peters of the first kind and poly-Cauchy of the first kind mixed-type polynomials, whose generating function is given by

$$\prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t))(1+t)^x = \sum_{n=0}^{\infty} s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \frac{t^n}{n!}, \quad (1)$$

where  $\lambda_1, \dots, \lambda_r, \mu_1, \dots, \mu_r \in \mathbb{C}$  with  $\lambda_1, \dots, \lambda_r \neq 0$ . Here,  $\text{Lif}_k(x)$  ( $k \in \mathbb{Z}$ ) is the polyfactorial function ([6]) defined by

$$\text{Lif}_k(x) = \sum_{m=0}^{\infty} \frac{x^m}{m!(m+1)^k}.$$

When  $x = 0$ ,  $s_n^{(k)} = s_n^{(k)}(0) = s_n^{(k)}(0|\lambda; \mu) = s_n^{(k)}(0; \lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r)$  is called the the Barnes-type Peters of the first kind and poly-Cauchy of the first kind mixed-type number.

Recall that the Barnes-type Peters polynomials of the first kind, denoted by  $s_n(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r)$ , are given by the generating function as

$$\prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} (1+t)^x = \sum_{n=0}^{\infty} s_n(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \frac{t^n}{n!}.$$

If  $r = 1$ , then  $s_n(x|\lambda; \mu)$  are the Peters polynomials of the first kind. Peters polynomials were mentioned in [9, p.128] and have been investigated in e.g. [5].

The poly-Cauchy polynomials of the first kind, denoted by  $c_n^{(k)}(x)$  ([3, 7]), are given by the generating function as

$$\text{Lif}_k(\ln(1+t))(1+t)^x = \sum_{n=0}^{\infty} c_n^{(k)}(-x) \frac{t^n}{n!}.$$

The generalized Barnes-type Euler polynomials  $E_n(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r)$  are defined by the generating function

$$\prod_{j=1}^r \left( \frac{2}{1 + e^{\lambda_j t}} \right)^{\mu_j} e^{xt} = \sum_{n=0}^{\infty} E_n(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \frac{t^n}{n!}.$$

If  $\mu_1 = \dots = \mu_r = 1$ , then  $E_n(x|\lambda_1, \dots, \lambda_r) = E_n(x|\lambda_1, \dots, \lambda_r; 1, \dots, 1)$  are called the Barnes-type Euler polynomials. If further  $\lambda_1 = \dots = \lambda_r = 1$ , then  $E_n^{(r)}(x) = E_n(x|1, \dots, 1; 1, \dots, 1)$  are called the Euler polynomials of order  $r$ .

In this paper, by considering Barnes-type Peters polynomials of the first kind as well as poly-Cauchy polynomials of the first kind, we define and investigate the mixed-type polynomials of these polynomials. From the properties of Sheffer sequences of these polynomials arising from umbral calculus, we derive new and interesting identities.

## 2 Umbral calculus

Let  $\mathbb{C}$  be the complex number field and let  $\mathcal{F}$  be the set of all formal power series in the variable  $t$ :

$$\mathcal{F} = \left\{ f(t) = \sum_{k=0}^{\infty} \frac{a_k}{k!} t^k \mid a_k \in \mathbb{C} \right\}. \tag{2}$$

Let  $\mathbb{P} = \mathbb{C}[x]$  and let  $\mathbb{P}^*$  be the vector space of all linear functionals on  $\mathbb{P}$ .  $\langle L|p(x) \rangle$  is the action of the linear functional  $L$  on the polynomial  $p(x)$ , and we recall that the vector space operations on  $\mathbb{P}^*$  are defined by  $\langle L + M|p(x) \rangle = \langle L|p(x) \rangle + \langle M|p(x) \rangle$ ,  $\langle cL|p(x) \rangle = c \langle L|p(x) \rangle$ , where  $c$  is a complex constant in  $\mathbb{C}$ . For  $f(t) \in \mathcal{F}$ , let us define the linear functional on  $\mathbb{P}$  by setting

$$\langle f(t)|x^n \rangle = a_n, \quad (n \geq 0). \tag{3}$$

In particular,

$$\langle t^k|x^n \rangle = n! \delta_{n,k} \quad (n, k \geq 0), \tag{4}$$

where  $\delta_{n,k}$  is the Kronecker's symbol.

For  $f_L(t) = \sum_{k=0}^{\infty} \frac{\langle L|x^k \rangle}{k!} t^k$ , we have  $\langle f_L(t)|x^n \rangle = \langle L|x^n \rangle$ . That is,  $L = f_L(t)$ . The map  $L \mapsto f_L(t)$  is a vector space isomorphism from  $\mathbb{P}^*$  onto  $\mathcal{F}$ . Henceforth,  $\mathcal{F}$  denotes both the algebra of formal power series in  $t$  and the vector space of all linear functionals on  $\mathbb{P}$ , and so an element  $f(t)$  of  $\mathcal{F}$  will be thought of as both a formal power series and a linear functional. We call  $\mathcal{F}$  the *umbral algebra* and the *umbral calculus* is the study of umbral algebra. The order  $O(f(t))$  of a power series  $f(t) (\neq 0)$  is the smallest integer  $k$  for which the coefficient of  $t^k$  does not vanish. If  $O(f(t)) = 1$ , then  $f(t)$  is called a *delta series*; if  $O(f(t)) = 0$ , then  $f(t)$  is called an *invertible series*. For  $f(t), g(t) \in \mathcal{F}$  with  $O(f(t)) = 1$  and  $O(g(t)) = 0$ , there exists a unique sequence  $s_n(x)$  ( $\deg s_n(x) = n$ ) such that  $\langle g(t)f(t)^k|s_n(x) \rangle = n! \delta_{n,k}$ , for  $n, k \geq 0$ . Such a sequence  $s_n(x)$  is called the *Sheffer sequence* for  $(g(t), f(t))$  which is denoted by  $s_n(x) \sim (g(t), f(t))$ .

For  $f(t), g(t) \in \mathcal{F}$  and  $p(x) \in \mathbb{P}$ , we have

$$\langle f(t)g(t)|p(x) \rangle = \langle f(t)|g(t)p(x) \rangle = \langle g(t)|f(t)p(x) \rangle \tag{5}$$

and

$$f(t) = \sum_{k=0}^{\infty} \langle f(t)|x^k \rangle \frac{t^k}{k!}, \quad p(x) = \sum_{k=0}^{\infty} \langle t^k|p(x) \rangle \frac{x^k}{k!} \tag{6}$$

([9, Theorem 2.2.5]). Thus, by (6), we get

$$t^k p(x) = p^{(k)}(x) = \frac{d^k p(x)}{dx^k} \quad \text{and} \quad e^{yt} p(x) = p(x + y). \tag{7}$$

Sheffer sequences are characterized in the generating function ([9, Theorem 2.3.4]).

**Lemma 1** *The sequence  $s_n(x)$  is Sheffer for  $(g(t), f(t))$  if and only if*

$$\frac{1}{g(\bar{f}(t))} e^{y\bar{f}(t)} = \sum_{k=0}^{\infty} \frac{s_k(y)}{k!} t^k \quad (y \in \mathbb{C}),$$

where  $\bar{f}(t)$  is the compositional inverse of  $f(t)$ .

For  $s_n(x) \sim (g(t), f(t))$ , we have the following equations ([9, Theorem 2.3.7, Theorem 2.3.5, Theorem 2.3.9]):

$$f(t)s_n(x) = ns_{n-1}(x) \quad (n \geq 0), \tag{8}$$

$$s_n(x) = \sum_{j=0}^n \frac{1}{j!} \left\langle g(\bar{f}(t))^{-1} \bar{f}(t)^j | x^n \right\rangle x^j, \tag{9}$$

$$s_n(x+y) = \sum_{j=0}^n \binom{n}{j} s_j(x) p_{n-j}(y), \tag{10}$$

where  $p_n(x) = g(t)s_n(x)$ .

Assume that  $p_n(x) \sim (1, f(t))$  and  $q_n(x) \sim (1, g(t))$ . Then the transfer formula ([9, Corollary 3.8.2]) is given by

$$q_n(x) = x \left( \frac{f(t)}{g(t)} \right)^n x^{-1} p_n(x) \quad (n \geq 1).$$

For  $s_n(x) \sim (g(t), f(t))$  and  $r_n(x) \sim (h(t), l(t))$ , assume that

$$s_n(x) = \sum_{m=0}^n C_{n,m} r_m(x) \quad (n \geq 0).$$

Then we have ([9, p.132])

$$C_{n,m} = \frac{1}{m!} \left\langle \frac{h(\bar{f}(t))}{g(\bar{f}(t))} l(\bar{f}(t))^m | x^n \right\rangle. \tag{11}$$

### 3 Main results

From the definition (1),  $s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r)$  is the Sheffer sequence for the pair

$$g(t) = \prod_{j=1}^r (1 + e^{\lambda_j t})^{\mu_j} \frac{1}{\text{Lif}_k(t)} \quad \text{and} \quad f(t) = e^t - 1.$$

So,

$$s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \sim \left( \prod_{j=1}^r (1 + e^{\lambda_j t})^{\mu_j} \frac{1}{\text{Lif}_k(t)}, e^t - 1 \right). \tag{12}$$

### 3.1 Explicit expressions

Let  $(n)_j = n(n-1)\cdots(n-j+1)$  ( $j \geq 1$ ) with  $(n)_0 = 1$ . The (signed) Stirling numbers of the first kind  $S_1(n, m)$  are defined by

$$(x)_n = \sum_{m=0}^n S_1(n, m)x^m.$$

#### Theorem 1

$$\begin{aligned} & s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\ &= 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n S_1(n, m) \sum_{l=0}^m \frac{\binom{m}{l}}{(l+1)^k} E_{m-l}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \end{aligned} \quad (13)$$

$$= \sum_{j=0}^n \sum_{l=0}^{n-j} \binom{n}{l} S_1(n-l, j) s_l^{(k)} x^j \quad (14)$$

$$= \sum_{j=0}^n \sum_{l=0}^{n-j} \sum_{i=0}^l \binom{n}{l} \binom{l}{i} S_1(n-l, j) c_i^{(k)} s_{l-i}(\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) x^j \quad (15)$$

$$= \sum_{l=0}^n \binom{n}{l} s_{n-l}(\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) c_l^{(k)}(-x), \quad (16)$$

$$= \sum_{l=0}^n \binom{n}{l} s_l(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) c_{n-l}^{(k)}. \quad (17)$$

*Proof.* Since

$$\prod_{j=1}^r (1 + e^{\lambda_j t})^{\mu_j} \frac{1}{\text{Lif}_k(t)} s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \sim (1, e^t - 1) \quad (18)$$

and

$$(x)_n \sim (1, e^t - 1), \quad (19)$$



we have

$$\begin{aligned}
 s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) &= s_n^{(k)}(x) \\
 &= \prod_{j=1}^r (1 + e^{\lambda_j t})^{-\mu_j} \text{Lif}_k(t)(x)_n \\
 &= \sum_{m=0}^n S_1(n, m) \prod_{j=1}^r (1 + e^{\lambda_j t})^{-\mu_j} \text{Lif}_k(t)x^m \\
 &= \sum_{m=0}^n S_1(n, m) \prod_{j=1}^r (1 + e^{\lambda_j t})^{-\mu_j} \sum_{l=0}^m \frac{t^l}{l!(l+1)^k} x^m \\
 &= \sum_{m=0}^n S_1(n, m) \prod_{j=1}^r (1 + e^{\lambda_j t})^{-\mu_j} \sum_{l=0}^m \frac{\binom{m}{l}}{l!(l+1)^k} x^{m-l} \\
 &= \sum_{m=0}^n S_1(n, m) \sum_{l=0}^m \frac{\binom{m}{l}}{(l+1)^k} \prod_{j=1}^r (1 + e^{\lambda_j t})^{-\mu_j} x^{m-l} \\
 &= 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n S_1(n, m) \sum_{l=0}^m \frac{\binom{m}{l}}{(l+1)^k} \prod_{j=1}^r \left(\frac{2}{1 + e^{\lambda_j t}}\right)^{\mu_j} x^{m-l} \\
 &= 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n S_1(n, m) \sum_{l=0}^m \frac{\binom{m}{l}}{(l+1)^k} E_{m-l}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r).
 \end{aligned}$$

So, we get (13).

By (9) with (12), we get

$$\begin{aligned}
 &\left\langle g(\bar{f}(t))^{-1} \bar{f}(t)^j | x^n \right\rangle \\
 &= \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) (\ln(1+t))^j | x^n \right\rangle \\
 &= \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) \left| j! \sum_{l=j}^{\infty} S_1(l, j) \frac{t^l}{l!} x^n \right. \right\rangle \\
 &= j! \sum_{l=j}^n \binom{n}{l} S_1(l, j) \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) | x^{n-l} \right\rangle \\
 &= j! \sum_{l=j}^n \binom{n}{l} S_1(l, j) \left\langle \sum_{i=0}^{\infty} s_i^{(k)} \frac{t^i}{i!} | x^{n-l} \right\rangle \\
 &= j! \sum_{l=j}^n \binom{n}{l} S_1(l, j) s_{n-l}^{(k)}.
 \end{aligned}$$

On the other hand,

$$\begin{aligned}
 & \left\langle g(\bar{f}(t))^{-1} \bar{f}(t)^j |x^n \right\rangle \\
 &= j! \sum_{l=j}^n \binom{n}{l} S_1(l, j) \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \Big| \text{Lif}_k(\ln(1+t)) x^{n-l} \right\rangle \\
 &= j! \sum_{l=j}^n \binom{n}{l} S_1(l, j) \sum_{i=0}^{n-l} \binom{n-l}{i} c_i^{(k)} \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \Big| x^{n-l-i} \right\rangle \\
 &= j! \sum_{l=j}^n \binom{n}{l} S_1(l, j) \sum_{i=0}^{n-l} \binom{n-l}{i} c_i^{(k)} \left\langle \sum_{m=0}^{\infty} s_m(\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \frac{t^m}{m!} \Big| x^{n-l-i} \right\rangle \\
 &= j! \sum_{l=j}^n \sum_{i=0}^{n-l} \binom{n}{l} \binom{n-l}{i} S_1(l, j) c_i^{(k)} s_{n-l-i}(\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r).
 \end{aligned}$$

Thus, we obtain

$$\begin{aligned}
 s_n^{(k)}(x) &= \sum_{j=0}^n \sum_{l=0}^{n-j} \binom{n}{l} S_1(n-l, j) s_l^{(k)} x^j \\
 &= \sum_{j=0}^n \sum_{l=0}^{n-j} \sum_{i=0}^l \binom{n}{l} \binom{l}{i} S_1(n-l, j) c_i^{(k)} s_{l-i}(\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) x^j,
 \end{aligned}$$

which are the identities (14) and (15).

Next,

$$\begin{aligned}
 s_n^{(k)}(y|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) &= \left\langle \sum_{i=0}^{\infty} s_i^{(k)}(y|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \frac{t^i}{i!} \middle| x^n \right\rangle \\
 &= \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t))(1+t)^y \middle| x^n \right\rangle \\
 &= \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \middle| \text{Lif}_k(\ln(1+t))(1+t)^y x^n \right\rangle \\
 &= \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \middle| \sum_{l=0}^{\infty} c_l^{(k)}(-y) \frac{t^l}{l!} x^n \right\rangle \\
 &= \sum_{l=0}^n \binom{n}{l} c_l^{(k)}(-y) \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \middle| x^{n-l} \right\rangle \\
 &= \sum_{l=0}^n \binom{n}{l} c_l^{(k)}(-y) \left\langle \sum_{i=0}^{\infty} s_i(\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \frac{t^i}{i!} \middle| x^{n-l} \right\rangle \\
 &= \sum_{l=0}^n \binom{n}{l} c_l^{(k)}(-y) s_{n-l}(\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r).
 \end{aligned}$$

Thus, we obtain (16).

Finally, we obtain that

$$\begin{aligned}
 s_n^{(k)}(y|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) &= \left\langle \sum_{i=0}^{\infty} s_i^{(k)}(y|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \frac{t^i}{i!} \middle| x^n \right\rangle \\
 &= \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t))(1+t)^y \middle| x^n \right\rangle \\
 &= \left\langle \text{Lif}_k(\ln(1+t)) \middle| \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} (1+t)^y x^n \right\rangle \\
 &= \left\langle \text{Lif}_k(\ln(1+t)) \middle| \sum_{l=0}^{\infty} s_l(y|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \frac{t^l}{l!} x^n \right\rangle \\
 &= \sum_{l=0}^n s_l(y|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \binom{n}{l} \left\langle \text{Lif}_k(\ln(1+t)) \middle| x^{n-l} \right\rangle \\
 &= \sum_{l=0}^n s_l(y|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \binom{n}{l} \left\langle \sum_{i=0}^{\infty} c_i^{(k)} \frac{t^i}{i!} \middle| x^{n-l} \right\rangle \\
 &= \sum_{l=0}^n \binom{n}{l} s_l(y|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) c_{n-l}^{(k)}.
 \end{aligned}$$

Thus, we get the identity (17). ■

### 3.2 Sheffer identity

#### Theorem 2

$$s_n^{(k)}(x + y | \lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) = \sum_{j=0}^n \binom{n}{j} s_j^{(k)}(x | \lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) (y)_{n-j}. \quad (20)$$

*Proof.* By (12) with

$$\begin{aligned} p_n(x) &= \prod_{j=1}^r (1 + e^{\lambda_j t})^{\mu_j} \frac{1}{\text{Lif}_k(t)} s_n(x | \lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\ &= (x)_n \sim (1, e^t - 1), \end{aligned}$$

using (10), we have (20). ■

### 3.3 Difference relations

#### Theorem 3

$$\begin{aligned} s_n^{(k)}(x + 1 | \lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) - s_n^{(k)}(x | \lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\ = n s_{n-1}^{(k)}(x | \lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r). \quad (21) \end{aligned}$$

*Proof.* By (8) with (12), we get

$$(e^t - 1) s_n^{(k)}(x | \lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) = n s_{n-1}^{(k)}(x | \lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r).$$

By (7), we have (21). ■

### 3.4 Recurrence

#### Theorem 4

$$\begin{aligned}
 & s_{n+1}^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\
 &= x s_n^{(k)}(x-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\
 &\quad - 2^{-1-\sum_{j=1}^r \mu_j} \sum_{m=0}^n \sum_{l=0}^m \sum_{i=1}^r S_1(n, m) \frac{\binom{m}{l}}{(l+1)^k} \mu_i \lambda_i E_{m-l}(x+\lambda_i-1|\lambda; \mu+e_i) \\
 &\quad + 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n \sum_{l=0}^m \frac{\binom{m}{l}}{(m+2-l)^k} S_1(n, m) E_l(x-1|\lambda; \mu) \tag{22}
 \end{aligned}$$

$$\begin{aligned}
 &= x s_n^{(k)}(x-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\
 &\quad - \frac{1}{2} \sum_{j=0}^n \sum_{l=0}^{n-j} \sum_{i=1}^r \binom{n}{l} \mu_i \lambda_i^{j+1} S_1(n-l, j) s_l^{(k)} E_j\left(\frac{x+\lambda_i-1}{\lambda_i}\right) \\
 &\quad + 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n \sum_{l=0}^m \frac{\binom{m}{l}}{(m+2-l)^k} S_1(n, m) E_l(x-1|\lambda; \mu). \tag{23}
 \end{aligned}$$

*Proof.* By applying

$$s_{n+1}(x) = \left(x - \frac{g'(t)}{g(t)}\right) \frac{1}{f'(t)} s_n(x) \tag{24}$$

([9, Corollary 3.7.2]) with (12), we get

$$\begin{aligned}
 & s_{n+1}^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\
 &= x s_n^{(k)}(x-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) - e^{-t} \frac{g'(t)}{g(t)} s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r).
 \end{aligned}$$

Since

$$\begin{aligned}
 \frac{g'(t)}{g(t)} &= (\ln g(t))' \\
 &= \left(\sum_{i=1}^r \mu_i \ln(1+e^{\lambda_i t}) - \ln \text{Lif}_k(t)\right)' \\
 &= \sum_{i=1}^r \frac{\mu_i \lambda_i e^{\lambda_i t}}{1+e^{\lambda_i t}} - \frac{\text{Lif}'_k(t)}{\text{Lif}_k(t)},
 \end{aligned}$$

by (13), we have

$$\begin{aligned}
 & \frac{g'(t)}{g(t)} s_n^{(k)}(x) \\
 &= \left( \sum_{i=1}^r \frac{\mu_i \lambda_i e^{\lambda_i t}}{1 + e^{\lambda_i t}} - \frac{\text{Lif}'_k(t)}{\text{Lif}_k(t)} \right) s_n^{(k)}(x) \\
 &= 2^{-1 - \sum_{j=1}^r \mu_j} \sum_{m=0}^n \sum_{l=0}^m S_1(n, m) \frac{\binom{m}{l}}{(l+1)^k} \sum_{i=1}^r \mu_i \lambda_i e^{\lambda_i t} \frac{2}{1 + e^{\lambda_i t}} \prod_{j=1}^r \left( \frac{2}{1 + e^{\lambda_j t}} \right)^{\mu_j} x^{m-l} \\
 &\quad - \sum_{m=0}^n S_1(n, m) \prod_{j=1}^r (1 + e^{\lambda_j t})^{-\mu_j} \text{Lif}'_k(t) x^m. \tag{25}
 \end{aligned}$$

The first term in (25) is

$$2^{-1 - \sum_{j=1}^r \mu_j} \sum_{m=0}^n \sum_{l=0}^m \sum_{i=1}^r S_1(n, m) \frac{\binom{m}{l}}{(l+1)^k} \mu_i \lambda_i E_{m-l}(x + \lambda_i | \lambda; \mu + e_i),$$

where  $\lambda = (\lambda_1, \dots, \lambda_r)$ ,  $\mu = (\mu_1, \dots, \mu_r)$  and  $e_i = (\underbrace{0, \dots, 0}_{i-1}, 1, \underbrace{0, \dots, 0}_{r-i})$  ( $i = 1, 2, \dots, r$ ).

Since

$$\text{Lif}_{k-1}(t) - \text{Lif}_k(t) = \left( \frac{1}{2^{k-1}} - \frac{1}{2^k} \right) t + \dots,$$

the second term in (25) is

$$\begin{aligned}
 & 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n S_1(n, m) \text{Lif}'_k(t) \prod_{j=1}^r \left( \frac{2}{1 + e^{\lambda_j t}} \right)^{\mu_j} x^m \\
 &= 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n S_1(n, m) \frac{\text{Lif}_{k-1}(t) - \text{Lif}_k(t)}{t} E_m(x | \lambda; \mu) \\
 &= 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n S_1(n, m) (\text{Lif}_{k-1}(t) - \text{Lif}_k(t)) \frac{E_{m+1}(x | \lambda; \mu)}{m+1} \\
 &= 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n \frac{S_1(n, m)}{m+1} \left( \sum_{l=0}^{m+1} \frac{t^l}{l!(l+1)^{k-1}} E_{m+1-l}(x | \lambda; \mu) - \sum_{l=0}^{m+1} \frac{t^l}{l!(l+1)^k} E_{m+1-l}(x | \lambda; \mu) \right) \\
 &= 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n \frac{S_1(n, m)}{m+1} \left( \sum_{l=0}^{m+1} \frac{\binom{m+1}{l}}{(l+1)^{k-1}} E_{m+1-l}(x | \lambda; \mu) - \sum_{l=0}^{m+1} \frac{\binom{m+1}{l}}{(l+1)^k} E_{m+1-l}(x | \lambda; \mu) \right) \\
 &= 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n \frac{S_1(n, m)}{m+1} \sum_{l=1}^{m+1} \frac{\binom{m+1}{l} l}{(l+1)^k} E_{m+1-l}(x | \lambda; \mu) \\
 &= 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n S_1(n, m) \sum_{l=0}^m \frac{\binom{m}{l}}{(m+2-l)^k} E_l(x | \lambda; \mu).
 \end{aligned}$$

Thus, we obtain

$$\begin{aligned}
 s_{n+1}^{(k)}(x) &= x s_n^{(k)}(x-1) \\
 &\quad - 2^{-1-\sum_{j=1}^r \mu_j} \sum_{m=0}^n \sum_{l=0}^m \sum_{i=1}^r S_1(n, m) \frac{\binom{m}{l}}{(l+1)^k} \mu_i \lambda_i E_{m-l}(x + \lambda_i - 1 | \lambda; \mu + e_i) \\
 &\quad + 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n \sum_{l=0}^m \frac{\binom{m}{l}}{(m+2-l)^k} S_1(n, m) E_l(x-1 | \lambda; \mu),
 \end{aligned}$$

which is (22).

On the other hand, by (14) with (22), we have

$$\begin{aligned}
 &\frac{g'(t)}{g(t)} s_n^{(k)}(x) \\
 &= \left( \sum_{i=1}^r \frac{\mu_i \lambda_i e^{\lambda_i t}}{1 + e^{\lambda_i t}} - \frac{\text{Lif}'_k(t)}{\text{Lif}_k(t)} \right) s_n^{(k)}(x) \\
 &= \frac{1}{2} \sum_{i=1}^r \mu_i \lambda_i e^{\lambda_i t} \frac{2}{1 + e^{\lambda_i t}} \sum_{j=0}^n \sum_{l=0}^{n-j} \binom{n}{l} S_1(n-l, j) s_l^{(k)} x^j \\
 &\quad - 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n \sum_{l=0}^m \frac{\binom{m}{l}}{(m+2-l)^k} S_1(n, m) E_l(x | \lambda; \mu). \tag{26}
 \end{aligned}$$

The first term in (26) is

$$\begin{aligned}
 &\frac{1}{2} \sum_{j=0}^n \sum_{l=0}^{n-j} \binom{n}{l} S_1(n-l, j) s_l^{(k)} \sum_{i=1}^r \mu_i \lambda_i e^{\lambda_i t} \frac{2}{1 + e^{\lambda_i t}} x^j \\
 &= \frac{1}{2} \sum_{j=0}^n \sum_{l=0}^{n-j} \binom{n}{l} S_1(n-l, j) s_l^{(k)} \sum_{i=1}^r \mu_i \lambda_i e^{\lambda_i t} \lambda_i^j E_j \left( \frac{x}{\lambda_i} \right) \\
 &= \frac{1}{2} \sum_{j=0}^n \sum_{l=0}^{n-j} \binom{n}{l} S_1(n-l, j) s_l^{(k)} \sum_{i=1}^r \mu_i \lambda_i^{j+1} E_j \left( \frac{x + \lambda_i}{\lambda_i} \right).
 \end{aligned}$$

Thus, we obtain

$$\begin{aligned}
 s_{n+1}^{(k)}(x) &= x s_n^{(k)}(x-1) \\
 &\quad - \frac{1}{2} \sum_{j=0}^n \sum_{l=0}^{n-j} \sum_{i=1}^r \binom{n}{l} \mu_i \lambda_i^{j+1} S_1(n-l, j) s_l^{(k)} E_j \left( \frac{x + \lambda_i - 1}{\lambda_i} \right) \\
 &\quad + 2^{-\sum_{j=1}^r \mu_j} \sum_{m=0}^n \sum_{l=0}^m \frac{\binom{m}{l}}{(m+2-l)^k} S_1(n, m) E_l(x-1 | \lambda; \mu).
 \end{aligned}$$

which is (23). ■

### 3.5 Differentiation

#### Theorem 5

$$\frac{d}{dx} s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) = n! \sum_{l=0}^{n-1} \frac{(-1)^{n-l-1}}{l!(n-l)} s_l^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r). \quad (27)$$

*Proof.* We shall use

$$\frac{d}{dx} s_n(x) = \sum_{l=0}^{n-1} \binom{n}{l} \langle \bar{f}(t) | x^{n-l} \rangle s_l(x)$$

(Cf. [9, Theorem 2.3.12]). Since

$$\begin{aligned} \langle \bar{f}(t) | x^{n-l} \rangle &= \langle \ln(1+t) | x^{n-l} \rangle \\ &= \left\langle \sum_{m=1}^{\infty} \frac{(-1)^{m-1} t^m}{m} \middle| x^{n-l} \right\rangle \\ &= \sum_{m=1}^{n-l} \frac{(-1)^{m-1}}{m} \langle t^m | x^{n-l} \rangle \\ &= \sum_{m=1}^{n-l} \frac{(-1)^{m-1}}{m} (n-l)! \delta_{m,n-l} \\ &= (-1)^{n-l-1} (n-l-1)!, \end{aligned}$$

with (12), we have

$$\begin{aligned} &\frac{d}{dx} s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\ &= \sum_{l=0}^{n-1} \binom{n}{l} (-1)^{n-l-1} (n-l-1)! s_l^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\ &= n! \sum_{l=0}^{n-1} \frac{(-1)^{n-l-1}}{l!(n-l)} s_l^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r), \end{aligned}$$

which is the identity (27). ■

### 3.6 A more relation

The classical Cauchy numbers  $c_n$  are defined by

$$\frac{t}{\ln(1+t)} = \sum_{n=0}^{\infty} c_n \frac{t^n}{n!}$$

(see e.g. [2, 6]).



**Theorem 6** For  $n \geq 1$ , we have

$$\begin{aligned}
 & s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\
 &= x s_{n-1}^{(k)}(x-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) + \frac{1}{n} \sum_{l=1}^n \binom{n}{l} c_{n-l} (s_l^{(k-1)}(x-1) - s_l^{(k)}(x-1)) \\
 &\quad - \sum_{i=1}^r \mu_i \lambda_i s_{n-1}^{(k)}(x+\lambda_i-1|\lambda; \mu+e_i). \tag{28}
 \end{aligned}$$

*Proof.* For  $n \geq 1$ , we have

$$\begin{aligned}
 & s_n^{(k)}(y|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\
 &= \left\langle \sum_{l=0}^{\infty} s_l^{(k)}(y|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \frac{t^l}{l!} \middle| x^n \right\rangle \\
 &= \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t))(1+t)^y \middle| x^n \right\rangle \\
 &= \left\langle \partial_t \left( \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t))(1+t)^y \right) \middle| x^{n-1} \right\rangle \\
 &= \left\langle \left( \partial_t \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \right) \text{Lif}_k(\ln(1+t))(1+t)^y \middle| x^{n-1} \right\rangle \\
 &\quad + \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \left( \partial_t \text{Lif}_k(\ln(1+t)) \right) (1+t)^y \middle| x^{n-1} \right\rangle \\
 &\quad + \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) (\partial_t (1+t)^y) \middle| x^{n-1} \right\rangle.
 \end{aligned}$$

The third term is

$$\begin{aligned}
 & y \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t))(1+t)^{y-1} \middle| x^{n-1} \right\rangle \\
 &= y s_{n-1}^{(k)}(y-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r).
 \end{aligned}$$

Since

$$\text{Lif}_{k-1}(\ln(1+t)) - \text{Lif}_k(\ln(1+t)) = \left( \frac{1}{2^{k-1}} - \frac{1}{2^k} \right) t + \dots,$$

the second term is

$$\begin{aligned}
 & \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \frac{\text{Lif}_{k-1}(\ln(1+t)) - \text{Lif}_k(\ln(1+t))}{(1+t)\ln(1+t)} (1+t)^y \middle| x^{n-1} \right\rangle \\
 &= \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \frac{\text{Lif}_{k-1}(\ln(1+t)) - \text{Lif}_k(\ln(1+t))}{t} (1+t)^{y-1} \middle| \frac{t}{\ln(1+t)} x^{n-1} \right\rangle \\
 &= \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \frac{\text{Lif}_{k-1}(\ln(1+t)) - \text{Lif}_k(\ln(1+t))}{t} (1+t)^{y-1} \middle| \sum_{l=0}^{\infty} c_l \frac{t^l}{l!} x^{n-1} \right\rangle \\
 &= \sum_{l=0}^{n-1} \binom{n-1}{l} c_l \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} (1+t)^{y-1} \middle| \frac{\text{Lif}_{k-1}(\ln(1+t)) - \text{Lif}_k(\ln(1+t))}{t} x^{n-1-l} \right\rangle \\
 &= \sum_{l=0}^{n-1} \frac{1}{n-l} \binom{n-1}{l} c_l \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} (1+t)^{y-1} \middle| (\text{Lif}_{k-1}(\ln(1+t)) - \text{Lif}_k(\ln(1+t))) x^{n-l} \right\rangle \\
 &= \sum_{l=0}^{n-1} \frac{1}{n-l} \binom{n}{l} c_l \left( \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_{k-1}(\ln(1+t)) (1+t)^{y-1} \middle| x^{n-l} \right\rangle \right. \\
 &\quad \left. - \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) (1+t)^{y-1} \middle| x^{n-l} \right\rangle \right) \\
 &= \frac{1}{n} \sum_{l=0}^{n-1} \binom{n}{l} c_l (s_{n-l}^{(k-1)}(y-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) - s_{n-l}^{(k)}(y-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r)) \\
 &= \frac{1}{n} \sum_{l=1}^n \binom{n}{l} c_{n-l} (s_l^{(k-1)}(y-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) - s_l^{(k)}(y-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r)).
 \end{aligned}$$

Since

$$\begin{aligned} & \partial_t \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \\ &= - \sum_{i=1}^r \mu_i \lambda_i (1+t)^{\lambda_i-1} (1 + (1+t)^{\lambda_i})^{-1} \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j}, \end{aligned}$$

the first term is

$$\begin{aligned} & - \sum_{i=1}^r \mu_i \lambda_i \left\langle (1 + (1+t)^{\lambda_i})^{-1} \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t))(1+t)^{y+\lambda_i-1} | x^{n-1} \right\rangle \\ &= - \sum_{i=1}^r \mu_i \lambda_i s_{n-1}^{(k)}(y + \lambda_i - 1 | \lambda; \mu + e_i). \end{aligned}$$

Therefore, we obtain

$$\begin{aligned} & s_n^{(k)}(x | \lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\ &= x s_{n-1}^{(k)}(x-1 | \lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) + \frac{1}{n} \sum_{l=1}^n \binom{n}{l} c_{n-l} (s_l^{(k-1)}(x-1) - s_l^{(k)}(x-1)) \\ & \quad - \sum_{i=1}^r \mu_i \lambda_i s_{n-1}^{(k)}(x + \lambda_i - 1 | \lambda; \mu + e_i), \end{aligned}$$

which is the identity (28). ■

### 3.7 A relation including the Stirling numbers of the first kind

**Theorem 7** For  $n - 1 \geq m \geq 1$ , we have

$$\begin{aligned} & m \sum_{l=0}^{n-m} \binom{n}{l} S_1(n-l, m) s_l^{(k)}(\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\ &= (m-1) \sum_{l=0}^{n-m} \binom{n-1}{l} S_1(n-l-1, m-1) s_l^{(k)}(-1 | \lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\ & \quad + \sum_{l=0}^{n-m} \binom{n-1}{l} S_1(n-l-1, m-1) s_l^{(k-1)}(-1 | \lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\ & \quad - m \sum_{l=0}^{n-m-1} \sum_{i=1}^r \binom{n-1}{l} S_1(n-l-1, m) \mu_i \lambda_i s_l^{(k)}(\lambda_i - 1 | \lambda; \mu + e_i). \end{aligned} \tag{29}$$

*Proof.* We shall compute

$$\left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) (\ln(1+t))^m \Big| x^n \right\rangle$$

in two different ways. On the one hand, it is equal to

$$\begin{aligned} & \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) \Big| (\ln(1+t))^m x^n \right\rangle \\ &= \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) \Big| m! \sum_{l=m}^{\infty} S_1(l, m) \frac{t^l}{l!} x^n \right\rangle \\ &= m! \sum_{l=m}^n \binom{n}{l} S_1(l, m) \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) \Big| x^{n-l} \right\rangle \\ &= m! \sum_{l=m}^n \binom{n}{l} S_1(l, m) \left\langle \sum_{i=0}^{\infty} s_i^{(k)}(\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \frac{t^i}{i!} \Big| x^{n-l} \right\rangle \\ &= m! \sum_{l=m}^n \binom{n}{l} S_1(l, m) s_{n-l}^{(k)}(\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\ &= m! \sum_{l=0}^{n-m} \binom{n}{l} S_1(n-l, m) s_l^{(k)}(\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r). \end{aligned}$$

On the other hand, it is equal to

$$\begin{aligned} & \left\langle \partial_t \left( \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) (\ln(1+t))^m \right) \Big| x^{n-1} \right\rangle \\ &= \left\langle \left( \partial_t \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \right) \text{Lif}_k(\ln(1+t)) (\ln(1+t))^m \Big| x^{n-1} \right\rangle \\ & \quad + \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} (\partial_t \text{Lif}_k(\ln(1+t))) (\ln(1+t))^m \Big| x^{n-1} \right\rangle \\ & \quad + \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) (\partial_t (\ln(1+t))^m) \Big| x^{n-1} \right\rangle. \end{aligned} \tag{30}$$

The third term of (30) is equal to

$$\begin{aligned}
 & m \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t))(1+t)^{-1} \middle| (\ln(1+t))^{m-1} x^{n-1} \right\rangle \\
 &= m \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t))(1+t)^{-1} \middle| \right. \\
 &\quad \left. (m-1)! \sum_{l=m-1}^{\infty} S_1(l, m-1) \frac{t^l}{l!} x^{n-1} \right\rangle \\
 &= m! \sum_{l=m-1}^{n-1} \binom{n-1}{l} S_1(l, m-1) \\
 &\quad \times \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t))(1+t)^{-1} \middle| x^{n-1-l} \right\rangle \\
 &= m! \sum_{l=m-1}^{n-1} \binom{n-1}{l} S_1(l, m-1) s_{n-1-l}^{(k)}(-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\
 &= m! \sum_{l=0}^{n-m} \binom{n-1}{l} S_1(n-l-1, m-1) s_l^{(k)}(-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r).
 \end{aligned}$$

The second term of (30) is equal to

$$\begin{aligned}
 & \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \left( \frac{\text{Lif}_{k-1}(\ln(1+t)) - \text{Lif}_k(\ln(1+t))}{(1+t)\ln(1+t)} \right) (\ln(1+t))^m \middle| x^{n-1} \right\rangle \\
 &= \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_{k-1}(\ln(1+t))(1+t)^{-1} \middle| (\ln(1+t))^{m-1} x^{n-1} \right\rangle \\
 &\quad - \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t))(1+t)^{-1} \middle| (\ln(1+t))^{m-1} x^{n-1} \right\rangle \\
 &= (m-1)! \sum_{l=0}^{n-m} \binom{n-1}{l} S_1(n-l-1, m-1) s_l^{(k-1)}(-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\
 &\quad - (m-1)! \sum_{l=0}^{n-m} \binom{n-1}{l} S_1(n-l-1, m-1) s_l^{(k)}(-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r).
 \end{aligned}$$

The first term of (30) is equal to

$$\begin{aligned}
 & \left\langle \left( \partial_t \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \right) \text{Lif}_k(\ln(1+t)) (\ln(1+t))^m \Big| x^{n-1} \right\rangle \\
 &= - \sum_{i=1}^r \mu_i \lambda_i \left\langle (1 + (1+t)^{\lambda_i})^{-1} \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \right. \\
 & \quad \left. \text{Lif}_k(\ln(1+t)) (1+t)^{\lambda_i-1} \Big| (\ln(1+t))^m x^{n-1} \right\rangle \\
 &= - \sum_{i=1}^r \mu_i \lambda_i \left\langle (1 + (1+t)^{\lambda_i})^{-1} \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \right. \\
 & \quad \left. \text{Lif}_k(\ln(1+t)) (1+t)^{\lambda_i-1} \Big| m! \sum_{l=m}^{\infty} S_1(l, m) \frac{t^l}{l!} x^{n-1} \right\rangle \\
 &= -m! \sum_{i=1}^r \mu_i \lambda_i \sum_{l=m}^{n-1} \binom{n-1}{l} S_1(l, m) \\
 & \quad \times \left\langle (1 + (1+t)^{\lambda_i})^{-1} \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) (1+t)^{\lambda_i-1} \Big| x^{n-1-l} \right\rangle \\
 &= -m! \sum_{i=1}^r \mu_i \lambda_i \sum_{l=m}^{n-1} \binom{n-1}{l} S_1(l, m) s_{n-1-l}^{(k)}(\lambda_i - 1 | \lambda; \mu + e_i) \\
 &= -m! \sum_{l=0}^{n-m-1} \sum_{i=1}^r \binom{n-1}{l} S_1(n-1-l, m) \mu_i \lambda_i s_l^{(k)}(\lambda_i - 1 | \lambda; \mu + e_i).
 \end{aligned}$$

Therefore, we get, for  $n - 1 \geq m \geq 1$ ,

$$\begin{aligned}
 & m! \sum_{l=0}^{n-m} \binom{n}{l} S_1(n-l, m) s_l^{(k)}(\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\
 &= m! \sum_{l=0}^{n-m} \binom{n-1}{l} S_1(n-l-1, m-1) s_l^{(k)}(-1) \\
 & \quad + (m-1)! \sum_{l=0}^{n-m} \binom{n-1}{l} S_1(n-l-1, m-1) s_l^{(k-1)}(-1) \\
 & \quad - (m-1)! \sum_{l=0}^{n-m} \binom{n-1}{l} S_1(n-l-1, m-1) s_l^{(k)}(-1) \\
 & \quad - m! \sum_{l=0}^{n-m-1} \sum_{i=1}^r \binom{n-1}{l} S_1(n-l-1, m) \mu_i \lambda_i s_l^{(k)}(\lambda_i - 1 | \lambda; \mu + e_i).
 \end{aligned}$$

Dividing both sides by  $(m - 1)!$ , we obtain for  $n - 1 \geq m \geq 1$

$$\begin{aligned} & m \sum_{l=0}^{n-m} \binom{n}{l} S_1(n-l, m) s_l^{(k)}(\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\ &= (m-1) \sum_{l=0}^{n-m} \binom{n-1}{l} S_1(n-l-1, m-1) s_l^{(k)}(-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\ & \quad + \sum_{l=0}^{n-m} \binom{n-1}{l} S_1(n-l-1, m-1) s_l^{(k-1)}(-1|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\ & \quad - m \sum_{l=0}^{n-m-1} \sum_{i=1}^r \binom{n-1}{l} S_1(n-l-1, m) \mu_i \lambda_i s_l^{(k)}(\lambda_i - 1|\lambda; \mu + e_i). \end{aligned}$$

Thus, we get (29). ■

### 3.8 A relation with the falling factorials

#### Theorem 8

$$s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) = \sum_{m=0}^n \binom{n}{m} s_{n-m}^{(k)}(x)_m. \tag{31}$$

*Proof.* For (12) and (19), assume that  $s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) = \sum_{m=0}^n C_{n,m}(x)_m$ . By (11), we have

$$\begin{aligned} C_{n,m} &= \frac{1}{m!} \left\langle \frac{1}{\prod_{j=1}^r (1 + e^{\lambda_j \ln(1+t)})^{\mu_j}} \text{Lif}_k(\ln(1+t)) t^m \middle| x^n \right\rangle \\ &= \frac{1}{m!} \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) \middle| t^m x^n \right\rangle \\ &= \binom{n}{m} \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) \middle| x^{n-m} \right\rangle \\ &= \binom{n}{m} s_{n-m}^{(k)}. \end{aligned}$$

Thus, we get the identity (31). ■

### 3.9 A relation with higher-order Frobenius-Euler polynomials

For  $\alpha \in \mathbb{C}$  with  $\alpha \neq 1$ , the Frobenius-Euler polynomials of order  $r$ ,  $H_n^{(r)}(x|\alpha)$  are defined by the generating function

$$\left( \frac{1 - \alpha}{e^t - \alpha} \right)^r e^{xt} = \sum_{n=0}^{\infty} H_n^{(r)}(x|\alpha) \frac{t^n}{n!}$$

(see e.g. [4]).

**Theorem 9**

$$s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) = \sum_{m=0}^n \left( \sum_{j=0}^{n-m} \sum_{l=0}^{n-m-j} \binom{s}{j} \binom{n-j}{l} (n)_j \right. \\ \left. \times (1-\alpha)^{-j} S_1(n-j-l, m) s_l^{(k)} \right) H_m^{(s)}(x|\alpha). \tag{32}$$

*Proof.* For (12) and

$$H_n^{(s)}(x|\alpha) \sim \left( \left( \frac{e^t - \alpha}{1 - \alpha} \right)^s, t \right), \tag{33}$$

assume that  $s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) = \sum_{m=0}^n C_{n,m} H_m^{(s)}(x|\alpha)$ . By (11), similarly to the proof of (29), we have

$$C_{n,m} = \frac{1}{m!} \left\langle \frac{\left( \frac{e^{\ln(1+t)} - \alpha}{1 - \alpha} \right)^s}{\prod_{j=1}^r (1 + e^{\lambda_j \ln(1+t)})^{\mu_j}} \text{Lif}_k(\ln(1+t)) (\ln(1+t))^m \Big| x^n \right\rangle \\ = \frac{1}{m!(1-\alpha)^s} \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) (\ln(1+t))^m (1-\alpha+t)^s \Big| x^n \right\rangle \\ = \frac{1}{m!(1-\alpha)^s} \\ \times \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) (\ln(1+t))^m \Big| \sum_{i=0}^{\min\{s,n\}} \binom{s}{i} (1-\alpha)^{s-i} t^i x^n \right\rangle \\ = \frac{1}{m!(1-\alpha)^s} \sum_{i=0}^{n-m} \binom{s}{i} (1-\alpha)^{s-i} (n)_i \\ \times \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) (\ln(1+t))^m \Big| x^{n-i} \right\rangle \\ = \frac{1}{m!(1-\alpha)^s} \sum_{i=0}^{n-m} \binom{s}{i} (1-\alpha)^{s-i} (n)_i \sum_{l=0}^{n-m-i} m! \binom{n-i}{l} S_1(n-i-l, m) s_l^{(k)} \\ = \sum_{i=0}^{n-m} \sum_{l=0}^{n-m-i} \binom{s}{i} \binom{n-i}{l} (n)_i (1-\alpha)^{-i} S_1(n-i-l, m) s_l^{(k)}.$$

Thus, we get the identity (32). ■



### 3.10 A relation with higher-order Bernoulli polynomials

Bernoulli polynomials  $\mathfrak{B}_n^{(r)}(x)$  of order  $r$  are defined by

$$\left(\frac{t}{e^t - 1}\right)^r e^{xt} = \sum_{n=0}^{\infty} \frac{\mathfrak{B}_n^{(r)}(x)}{n!} t^n$$

(see e.g. [9, Section 2.2]). In addition, Cauchy numbers of the first kind  $\mathfrak{C}_n^{(r)}$  of order  $r$  are defined by

$$\left(\frac{t}{\ln(1+t)}\right)^r = \sum_{n=0}^{\infty} \frac{\mathfrak{C}_n^{(r)}}{n!} t^n$$

(see e.g. [1, (2.1)], [8, (6)]).

**Theorem 10**

$$\begin{aligned} & s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) \\ &= \sum_{m=0}^n \left( \sum_{i=0}^{n-m} \sum_{l=0}^{n-m-i} \binom{n}{i} \binom{n-i}{l} \mathfrak{C}_i^{(s)} S_1(n-i-l, m) s_l^{(k)} \right) \mathfrak{B}_m^{(s)}(x). \end{aligned} \tag{34}$$

*Proof.* For (12) and

$$\mathfrak{B}_n^{(s)}(x) \sim \left( \left( \frac{e^t - 1}{t} \right)^s, t \right), \tag{35}$$

assume that  $s_n^{(k)}(x|\lambda_1, \dots, \lambda_r; \mu_1, \dots, \mu_r) = \sum_{m=0}^n C_{n,m} \mathfrak{B}_m^{(s)}(x)$ . By (11), similarly to the proof of (29), we have

$$\begin{aligned} C_{n,m} &= \frac{1}{m!} \left\langle \frac{\left(\frac{e^{\ln(1+t)} - 1}{\ln(1+t)}\right)^s}{\prod_{j=1}^r (1 + e^{\lambda_j \ln(1+t)})^{\mu_j}} \text{Lif}_k(\ln(1+t)) (\ln(1+t))^m \Big| x^n \right\rangle \\ &= \frac{1}{m!} \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) (\ln(1+t))^m \Big| \left(\frac{t}{\ln(1+t)}\right)^s x^n \right\rangle \\ &= \frac{1}{m!} \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) (\ln(1+t))^m \Big| \sum_{i=0}^{\infty} \mathfrak{C}_i^{(s)} \frac{t^i}{i!} x^n \right\rangle \\ &= \frac{1}{m!} \sum_{i=0}^{n-m} \mathfrak{C}_i^{(s)} \binom{n}{i} \left\langle \prod_{j=1}^r (1 + (1+t)^{\lambda_j})^{-\mu_j} \text{Lif}_k(\ln(1+t)) (\ln(1+t))^m \Big| x^{n-i} \right\rangle \\ &= \frac{1}{m!} \sum_{i=0}^{n-m} \mathfrak{C}_i^{(s)} \binom{n}{i} \sum_{l=0}^{n-m-i} m! \binom{n-i}{l} S_1(n-i-l, m) s_l^{(k)} \\ &= \sum_{i=0}^{n-m} \sum_{l=0}^{n-m-i} \binom{n}{i} \binom{n-i}{l} \mathfrak{C}_i^{(s)} S_1(n-i-l, m) s_l^{(k)}. \end{aligned}$$

Thus, we get the identity (34). ■

ACKNOWLEDGEMENTS. This paper is supported by grant NO 14-11-00022 of Russian Scientific fund.

## References

- [1] L. Carlitz, *A note on Bernoulli and Euler polynomials of the second kind*, Scripta Math. **25** (1961), 323–330.
- [2] L. Comtet, *Advanced Combinatorics*, Reidel, Dordrecht, 1974.
- [3] D. S. Kim, T. Kim, S.-H. Lee, S.-H. Rim, *Umbral calculus and Euler polynomials*, Ars Combin. **112** (2013), 293–306.
- [4] D. S. Kim and T. Kim, *Some identities of Frobenius-Euler polynomials arising from umbral calculus*, Adv. Difference Equ. **2012** (2012), #196.
- [5] D. S. Kim and T. Kim, *Poly-Cauchy and Peters mixed-type polynomials*, Adv. Difference Equ. **2014**, (2014), #4.
- [6] D.S. Kim and T. Kim, *Higher-order Cauchy of the first kind and poly-Cauchy of the first kind mixed type polynomials*, Adv. Stud. Contemp. Math. **23** (2013), no. 4, 621–636.
- [7] D. S. Kim, T. Kim and S.-H. Lee, *Poly-Cauchy numbers and polynomials with umbral calculus viewpoint*, Int. J. Math. Anal. (Ruse) **7** (2013), no. 45-48, 2235–2253.
- [8] T. Kim, *Identities involving Laguerre polynomials derived from umbral calculus*, Russ. J. Math. Phys. **21** (2014), no. 1, 36–45.
- [9] S. Roman, *The umbral Calculus*, Dover, New York, 2005.

# On the hyperstability of a functional equation in commutative groups

Muaadh Almahalebi<sup>1</sup> and Choonkil Park<sup>2\*</sup>

<sup>1</sup>*Department of Mathematics, Ibn Tofail University, Kenitra, Morocco: muaadh1979@hotmail.fr*

<sup>2</sup>*Research Institute for Natural Sciences, Hanyang University, Seoul 133-791, Korea: baak@hanyang.ac.kr*

**Abstract.** Using the fixed point method, we prove the hyperstability of the functional equation

$$f(ax + by) = \frac{(a + b)}{2}f(x + y) + \frac{(a - b)}{2}f(x - y),$$

where  $a, b$  are different integers greater than 1, in the class of functions from a commutative group into a commutative complete metric group.

## 1. Introduction and preliminaries

The stability problem of functional equations originated from a question of Ulam [33] concerning the stability of group homomorphisms. Hyers [19] gave a first affirmative partial answer to the question of Ulam for Banach spaces. Hyers' Theorem was generalized by Aoki [4] for additive mappings and by Rassias [29] for linear mappings by considering an unbounded Cauchy difference. A generalization of the Th.M. Rassias theorem was obtained by Găvruta [17] by replacing the unbounded Cauchy difference by a general control function in the spirit of Th.M. Rassias' approach. The stability problems of several functional equations have been extensively investigated by a number of authors and there are many interesting results concerning this problem (see [6, 12, 13, 14, 15, 16, 20, 21, 25, 27, 28, 30, 31, 34]).

We say a functional equation  $\mathfrak{D}$  is *hyperstable* if any function  $f$  satisfying the equation  $\mathfrak{D}$  approximately is a true solution of  $\mathfrak{D}$ . It seems that the first hyperstability result was published in [6] and concerned the ring homomorphisms. However, The term *hyperstability* has been used for the first time in [24]. Quite often the hyperstability is confused with superstability, which admits also bounded functions. Numerous papers on this subject have been published and we refer to [1, 2, 3, 5, 7, 8, 9, 11, 18, 24, 26, 32].

Throughout this paper, we will denote the set of natural numbers by  $\mathbb{N}$ , the set of integers by  $\mathbb{Z}$  and the set of real numbers by  $\mathbb{R}$ . Let  $\mathbb{N}_+$  be the set of positive integers. By  $\mathbb{N}_m$ ,  $m \in \mathbb{N}_+$ , we will denote the set of all integers greater than or equal to  $m$ . Let  $\mathbb{R}_0 = [0, \infty)$  the set of nonnegative real numbers and  $\mathbb{R}_+ = (0, \infty)$  the set of positive real numbers. We write  $B^A$  to mean “ the family of all functions mapping from a nonempty set  $A$  into a nonempty set  $B$  ”.

**Definition 1.1.** Let  $X$  be a nonempty set,  $(Y, d)$  be a metric space,  $\varepsilon \in \mathbb{R}_0^{X^n}$  and  $\mathcal{F}_1, \mathcal{F}_2$  be operators mapping from a nonempty set  $\mathcal{D} \subset Y^X$  into  $Y^X$ . We say that the operator equation

$$\mathcal{F}_1\varphi(x_1, \dots, x_n) = \mathcal{F}_2\varphi(x_1, \dots, x_n), \quad (x_1, \dots, x_n \in X) \tag{1.1}$$

---

<sup>0</sup>2010 Mathematics Subject Classification: 39B52, 54E50, 39B82.

<sup>0</sup>Keywords: Hyers-Ulam stability; hyperstability; fixed point method; complete metric space.

\*Corresponding author: Choonkil Park (email: baak@hanyang.ac.kr).

is  $\varepsilon$ -hyperstable provided that every  $\varphi_0 \in \mathcal{D}$  which satisfies

$$d(\mathcal{F}_1\varphi_0(x_1, \dots, x_n), \mathcal{F}_2\varphi_0(x_1, \dots, x_n)) \leq \varepsilon(x_1, \dots, x_n), \quad (x_1, \dots, x_n \in X)$$

fulfills the equation (1.1).

In 2011, Kenary [22] introduced and proved the Hyers-Ulam stability for the following functional equation

$$f(ax + by) = \frac{(a + b)}{2}f(x + y) + \frac{(a - b)}{2}f(x - y) \tag{1.2}$$

in non-Archimedean normed spaces and in random normed spaces, where  $m, n$  are different integers greater than 1. In 2011, Kenary, Jang and Park [23] proved the Hyers-Ulam stability of (1.2) in various normed spaces by using the fixed point method.

In this paper, using the fixed point method derived from [10, Theorem 1], we prove the hyperstability of (1.2) in the class of functions from a commutative group into a commutative complete metric group.

Before proceeding to the main results, we state the following theorem which is useful for our purpose.

**Theorem 1.2.** ([10, Theorem 1]) *Let  $X$  be a nonempty set,  $(Y, d)$  a complete metric space,  $f_1, \dots, f_s: X \rightarrow X$  and  $L_1, \dots, L_s: X \rightarrow \mathbb{R}_0$  be given mappings. Let  $\Lambda: \mathbb{R}_0^X \rightarrow \mathbb{R}_0^X$  be a linear operator defined by*

$$\Lambda\delta(x) := \sum_{i=1}^s L_i(x)\delta(f_i(x)), \tag{1.3}$$

for  $\delta \in \mathbb{R}_0^X$  and  $x \in X$ . If  $\mathcal{T}: Y^X \rightarrow Y^X$  is an operator satisfying the inequality

$$d(\mathcal{T}\xi(x), \mathcal{T}\mu(x)) \leq \sum_{i=1}^s L_i(x)d(\xi(f_i(x)), \mu(f_i(x))), \quad \xi, \mu \in Y^X, x \in X,$$

and a function  $\varepsilon: X \rightarrow \mathbb{R}_0$  and a mapping  $\varphi: X \rightarrow Y$  satisfy

$$d(\mathcal{T}\varphi(x), \varphi(x)) \leq \varepsilon(x), \quad (x \in X),$$

$$\varepsilon^*(x) := \sum_{k=0}^{\infty} \Lambda^k \varepsilon(x) < \infty, \quad (x \in X),$$

then for every  $x \in X$ , the limit

$$\psi(x) := \lim_{n \rightarrow \infty} \mathcal{T}^n \varphi(x),$$

exists and the function  $\psi \in Y^X$  so defined is a unique fixed point of  $\mathcal{T}$  with

$$d(\varphi(x), \psi(x)) \leq \varepsilon^*(x), \quad (x \in X).$$

## 2. Main results

Suppose  $(G, +)$  and  $(H, +)$  are abelian groups, and  $d$  is a metric on  $H$  such that

(i):  $d$  is invariant with respect to  $+$ , that is,

$$d(u + w, v + w) = d(u, v), \quad (u, v, w \in H);$$

(ii):  $(H, d)$  is a complete metric space.

We will denote by  $Aut(G)$  the family of all automorphisms of  $G$ . Moreover, for each  $u: G \rightarrow G$  we write  $ux := u(x)$  for  $x \in G$  and we define  $u'$  by  $u'x := x - ux$  for  $x \in G$ .

Let

$$l(G) := \{u \in Aut(G) : (u' - u), (au' + bu) \in Aut(G),$$

$$\alpha_u := \frac{2}{a+b}\lambda(au' + bu) + \frac{|a-b|}{a+b}\lambda(u' - u) < 1\} \neq \emptyset, \tag{2.1}$$

where

$$\lambda(u) := \inf \{t \in \mathbb{R}_0 : \varepsilon(ux, uy) \leq t\varepsilon(x, y), \quad \forall x, y \in G\}$$

for  $u \in Aut(G)$  and  $\varepsilon: G^2 \rightarrow \mathbb{R}_0$ . The following theorem is a result concerning the hyperstability of the functional equation (1.2).

**Theorem 2.1.** *Let  $f: G \rightarrow H$  be a mapping satisfying the inequality*

$$d\left(f(x+y), \frac{2}{a+b}f(ax+by) - \frac{a-b}{a+b}f(x-y)\right) \leq \varepsilon(x, y) \tag{2.2}$$

for all  $x, y \in G$ , where  $\varepsilon: G^2 \rightarrow \mathbb{R}_0$  is an arbitrary function. Assume that there exists a nonempty subset  $\mathcal{U} \subset l(G)$  such that

$$u \circ v = v \circ u, \quad \forall u, v \in \mathcal{U},$$

and

$$\inf \{\varepsilon(u'x, ux) : u \in \mathcal{U}\} = 0, \quad \forall x \in G,$$

$$\sup \{\alpha_u : u \in \mathcal{U}\} < 1, \tag{2.3}$$

then  $f$  is a solution of (1.2) on  $G$ .

*Proof.* Let us fix  $u \in \mathcal{U}$ . Replacing  $x$  with  $u'x$  and  $y$  with  $ux$  in (2.2), we get

$$d\left(f(x), \frac{2}{a+b}f((au' + bu)x) - \frac{a-b}{a+b}f((u' - u)x)\right) \leq \varepsilon(u'x, ux) := \varepsilon_u(x) \tag{2.4}$$

for all  $x \in G$ . We define the operators  $\mathcal{T}_u: H^G \rightarrow H^G$  and  $\Lambda_u: \mathbb{R}_0^G \rightarrow \mathbb{R}_0^G$  by

$$\mathcal{T}_u\xi(x) := \frac{2}{a+b}\xi((au' + bu)x) - \frac{a-b}{a+b}\xi((u' - u)x), \tag{2.5}$$

$$\Lambda_u\delta(x) := \frac{2}{a+b}\delta((au' + bu)x) + \frac{|a-b|}{a+b}\delta((u' - u)x)$$

for all  $x \in G$ ,  $\xi \in H^G$  and  $\delta \in \mathbb{R}_0^G$ . Then (2.4) becomes

$$d(f(x), \mathcal{T}_u f(x)) \leq \varepsilon_u(x)$$

for all  $x \in G$ .

The operator  $\Lambda_u: \mathbb{R}_0^G \rightarrow \mathbb{R}_0^G$  has the form given by (1.3) with  $s = 2$  and  $f_1(x) = (au' + bu)x$ ,  $f_2(x) = (u' - u)x$ ,  $L_1(x) = \frac{2}{a+b}$ ,  $L_2(x) = \frac{|a-b|}{a+b}$  for all  $x \in G$ .

Further,

$$\begin{aligned}
 d(\mathcal{T}_u\xi(x), \mathcal{T}_u\mu(x)) &= d\left(\frac{2}{a+b}\xi((au' + bu)x) - \frac{a-b}{a+b}\xi((u' - u)x), \right. \\
 &\quad \left. \frac{2}{a+b}\mu((au' + bu)x) - \frac{a-b}{a+b}\mu((u' - u)x)\right) \\
 &\leq \frac{2}{a+b}d\left(\xi((au' + bu)x), \mu((au' + bu)x)\right) + \frac{|a-b|}{a+b}d\left(\xi((u' - u)x), \mu((u' - u)x)\right)
 \end{aligned}$$

for all  $x \in G$  and  $\xi, \mu \in H^G$ .

Note that, in view of the definition of  $\lambda(u)$ ,

$$\varepsilon(ux, uy) \leq \lambda(u)\varepsilon(x, y), \quad x, y \in G,$$

So it is easy to show by induction on  $k$  that

$$\Lambda_u^k \varepsilon_u(x) \leq \alpha_u^k \varepsilon(u'x, ux),$$

for all  $x \in G$ , where

$$\alpha_u = \left( \frac{2}{a+b}\lambda(au' + bu) + \frac{|a-b|}{a+b}\lambda(u' - u) \right).$$

Hence

$$\varepsilon^*(x) := \sum_{k=0}^{\infty} \Lambda_u^k \varepsilon_u(x) \leq \varepsilon(u'x, ux) \sum_{k=0}^{\infty} \alpha_u^k = \frac{\varepsilon(u'x, ux)}{1 - \alpha_u} < \infty$$

for all  $x \in G$ . By Theorem 1.2, there exists a unique solution  $F_u : G \rightarrow H$  of the equation

$$F_u(x) = \frac{2}{a+b}F_u((au' + bu)x) - \frac{a-b}{a+b}F_u((u' - u)x)$$

for all  $x \in G$ , which is a fixed point of  $\mathcal{T}_u$  such that

$$d(F_u(x), f(x)) \leq \frac{\varepsilon(u'x, ux)}{1 - \alpha_u}$$

for all  $x \in G$ . Moreover,

$$F_u(x) = \lim_{k \rightarrow \infty} \mathcal{T}_u^k f(x)$$

for all  $x \in G$ .

To prove that  $F_u$  satisfies the functional equation (1.2) on  $G$ , just prove the following inequality

$$d\left(\mathcal{T}_u^n f(x + y), \frac{2}{a+b}\mathcal{T}_u^n f(ax + by) - \frac{a-b}{a+b}\mathcal{T}_u^n f(x - y)\right) \leq \alpha_u^n \varepsilon(x, y) \tag{2.6}$$

for all  $x, y \in G$ , and  $n \in \mathbb{N}$ .

Indeed, if  $n = 0$  then (2.6) is simply (2.2). So, take  $n \in \mathbb{N}_+$  and suppose that (2.6) holds for  $n$

and  $x, y \in G$ . Then, by using (2.5) and the triangle inequality, we have

$$\begin{aligned} & d\left(\mathcal{T}_u^{n+1}f(x+y), \frac{2}{a+b}\mathcal{T}_u^{n+1}f(ax+by) - \frac{a-b}{a+b}\mathcal{T}_u^{n+1}f(x-y)\right) \\ &= d\left(\frac{2}{a+b}\mathcal{T}_u^n f((au'+bu)(x+y)) - \frac{a-b}{a+b}\mathcal{T}_u^n f((u'-u)(x+y)),\right. \\ &\quad \left. \left(\frac{2}{a+b}\right)^2\mathcal{T}_u^n f((au'+bu)(ax+by)) - \left(\frac{2}{a+b}\right)\left(\frac{a-b}{a+b}\right)\mathcal{T}_u^n f((u'-u)(ax+by))\right. \\ &\quad \left. - \left(\frac{2}{a+b}\right)\left(\frac{a-b}{a+b}\right)\mathcal{T}_u^n f((au'+bu)(x-y)) + \left(\frac{a-b}{a+b}\right)^2\mathcal{T}_u^n f((u'-u)(x-y))\right) \\ &\leq \frac{2}{a+b}d\left(\mathcal{T}_u^n f((au'+bu)(x+y)),\right. \\ &\quad \left. \frac{2}{a+b}\mathcal{T}_u^n f((au'+bu)(ax+by)) - \frac{a-b}{a+b}\mathcal{T}_u^n f((au'+bu)(x-y))\right) \\ &\quad + \frac{|a-b|}{a+b}d\left(\mathcal{T}_u^n f((u'-u)(x+y)),\right. \\ &\quad \left. \frac{2}{a+b}\mathcal{T}_u^n f((u'-u)(ax+by)) - \frac{a-b}{a+b}\mathcal{T}_u^n f((u'-u)(x-y))\right) \\ &\leq \alpha_u^n \left(\frac{2}{a+b}\varepsilon((au'+bu)x, (au'+bu)y) + \frac{|a-b|}{a+b}\varepsilon((u'-u)x, (u'-u)y)\right) \\ &\leq \varepsilon(x, y)\alpha_u^n \left(\frac{2}{a+b}\lambda(au'+bu) + \frac{|a-b|}{a+b}\lambda(u'-u)\right) \\ &= \alpha_u^{n+1}\varepsilon(x, y). \end{aligned}$$

By induction, we have shown that (2.6) holds for all  $x, y \in G$ . Letting  $n \rightarrow \infty$  in (2.6), we get

$$F_u(ax+by) = \frac{a+b}{2}F_u(x+y) + \frac{a-b}{2}F_u(x-y)$$

for all  $x, y \in G$ . Thus, we have proved that for every  $u \in \mathcal{U}$  there exists a function  $F_u : G \rightarrow H$  which is a solution of the functional equation (1.2) on  $G$  and satisfies

$$d(f(x), F_u(x)) \leq \frac{\varepsilon(u'x, ux)}{1 - \alpha_u}$$

for all  $x \in G$ . By (2.3), we get

$$\begin{aligned} d(f(x), F_u(x)) &\leq \frac{\inf_{u \in \mathcal{U}} \varepsilon(u'x, ux)}{1 - \sup_{u \in \mathcal{U}} \alpha_u} \\ &= 0 \end{aligned}$$

for all  $x \in G$ . This means that  $F_u(x) = f(x)$  for all  $x \in G$  and  $u \in \mathcal{U}$ , and hence

$$f(ax+by) = \frac{a+b}{2}f(x+y) + \frac{a-b}{2}f(x-y)$$

for all  $x, y \in G$ , which implies that  $f$  satisfies the functional equation (1.2) on  $G$ . □

In the next theorem, we will study the hyperstability of the functional equation (1.2) on  $G$  without 0 the neutral element, because of the reason that one can easily deduce some applications.

**Theorem 2.2.** Let  $f : G \rightarrow H$  be a mapping satisfying the inequality

$$d\left(f(x+y), \frac{2}{a+b}f(ax+by) - \frac{a-b}{a+b}f(x-y)\right) \leq \varepsilon(x,y)$$

for all  $x, y \in G \setminus \{0\}$ , where  $0$  is the neutral element of the group  $(G, +)$  and  $\varepsilon : (G \setminus \{0\})^2 \rightarrow \mathbb{R}_0$  is an arbitrary function. Assume that there exists a nonempty subset  $\mathcal{U} \subset l(G)$  such that

$$u \circ v = v \circ u \quad (u, v \in \mathcal{U}),$$

and

$$\begin{aligned} \inf \{ \varepsilon(u'x, ux) : u \in \mathcal{U} \} &= 0, \quad \forall x \in G \setminus \{0\}, \\ \sup \{ \alpha_u : u \in \mathcal{U} \} &< 1, \end{aligned} \tag{2.7}$$

then  $f$  is a solution of the functional equation (1.2) on  $G \setminus \{0\}$ .

*Proof.* The proof is the same as in the proof of Theorem 2.1. □

### 3. Some consequences

From Theorem 2.2, we can obtain the following corollaries as natural results.

**Corollary 3.1.** Let  $E$  and  $F$  be a normed space and a Banach space, respectively. Assume that  $X$  is a subgroup of the group  $(E, +)$ ,  $p < 0$ ,  $q < 0$  and  $c \geq 0$ . If  $f : X \rightarrow F$  satisfies

$$\left\| f(ax+by) - \frac{a+b}{2}f(x+y) - \frac{a-b}{2}f(x-y) \right\| \leq c(\|x\|^p + \|y\|^q) \tag{3.1}$$

for all  $x, y \in X \setminus \{0\}$ , then  $f$  satisfies the functional equation (1.2) on  $X \setminus \{0\}$ .

*Proof.* The proof follows from Theorem 2.2 by taking

$$\varepsilon(x, y) = c(\|x\|^p + \|y\|^q), \quad x, y \in X \setminus \{0\},$$

with some real numbers  $c \geq 0$ ,  $p < 0$ ,  $q < 0$  and  $d(x, y) = \|x - y\|$ . For each  $m \in \mathbb{N}_+$  define  $u_m : X \setminus \{0\} \rightarrow X \setminus \{0\}$  by  $u_mx := -mx$  and  $u'_m : X \setminus \{0\} \rightarrow X \setminus \{0\}$  by  $u'_mx := (1+m)x$ . Then

$$\begin{aligned} \varepsilon(u_mx, u_ky) &= \varepsilon(-mx, -ky) = c(\|-mx\|^p + \|-ky\|^q) \\ &= cm^p \|x\|^p + ck^q \|y\|^q \leq (m^p + k^q) c(\|x\|^p + \|y\|^q) \\ &= (m^p + k^q) \varepsilon(x, y) \end{aligned}$$

for all  $x \in X \setminus \{0\}$ ,  $k, m \in \mathbb{N}_+$ . Hence

$$\lim_{m \rightarrow \infty} \varepsilon(u'_m x, u_m y) \leq \lim_{m \rightarrow \infty} ((1+m)^p + m^q) \varepsilon(x, y) = 0$$

for all  $x, y \in X \setminus \{0\}$ . Then (2.7) is valid with  $\lambda(u_m) = m^p + m^q$  for  $m \in \mathbb{N}_+$ , and there exists  $n_0 \in \mathbb{N}_+$  such that

$$\frac{2}{a+b} \left( \left| a + (a-b)m \right|^p + \left| a + (a-b)m \right|^q \right) + \frac{|a-b|}{a+b} \left( (2m+1)^p + (2m+1)^q \right) < 1 \quad (m \geq n_0).$$

So it easily seen that (2.1) is fulfilled with

$$\mathcal{U} := \{u_m \in \text{Aut } X : m \in \mathbb{N}_{n_0}\}.$$

Therefore, by Theorem 2.2, every  $f : X \rightarrow F$  satisfying (3.1) is a solution of the functional equation (1.2) on  $X \setminus \{0\}$ . □



**Corollary 3.2.** *Let  $E$  and  $F$  be a normed space and a Banach space, respectively. Assume that  $X$  is a subgroup of the group  $(E, +)$ ,  $p, q \in \mathbb{R}$ ,  $p + q < 0$  and  $c \geq 0$ . If  $f : X \rightarrow F$  satisfies*

$$\left\| f(ax + by) - \frac{a+b}{2}f(x+y) - \frac{a-b}{2}f(x-y) \right\| \leq c \|x\|^p \|y\|^q$$

for all  $x, y \in X \setminus \{0\}$ , then  $f$  satisfies the functional equation (1.2) on  $X \setminus \{0\}$ .

*Proof.* It is easily seen that the function  $\varepsilon$  given by

$$\varepsilon(x, y) = c \|x\|^p \|y\|^q \quad x, y \in X \setminus \{0\},$$

satisfies (2.7), since

$$\varepsilon(mx, ky) = c \|mx\|^p \|ky\|^q = c |m|^p |k|^q \|x\|^p \|y\|^q = |m|^p |k|^q \varepsilon(x, y)$$

for all  $x, y \in X \setminus \{0\}$ ,  $k, m \in \mathbb{Z}$ , and  $km \neq 0$ .

The remainder of the proof is similar to the proof of Corollary 3.1. □

By an analogous conclusion, the function  $\varepsilon$  given by

$$\varepsilon(x, y) = c (\|x\|^p + \|y\|^q + \|x\|^p \|y\|^q), \quad x, y \in X \setminus \{0\},$$

satisfies (2.7), since

$$\begin{aligned} \varepsilon(mx, ky) &= c (\|mx\|^p + \|ky\|^q + \|mx\|^p \|ky\|^q) \\ &= c (|m|^p \|x\|^p + |k|^q \|y\|^q + |m|^p |k|^q \|x\|^p \|y\|^q) \\ &\leq (|m|^p + |k|^q + |m|^p |k|^q) \varepsilon(x, y) \end{aligned}$$

for all  $x, y \in X \setminus \{0\}$ ,  $k, m \in \mathbb{Z}$ , and  $km \neq 0$ . So we have the following corollary.

**Corollary 3.3.** *Let  $E$  and  $F$  be a normed space and a Banach space, respectively. Assume that  $X$  is a subgroup of the group  $(E, +)$ , and  $p < 0$ ,  $q < 0$ ,  $p + q < 0$  and  $c \geq 0$ . If  $f : X \rightarrow F$  satisfies*

$$\left\| f(ax + by) - \frac{a+b}{2}f(x+y) - \frac{a-b}{2}f(x-y) \right\| \leq c (\|x\|^p + \|y\|^q + \|x\|^p \|y\|^q)$$

for all  $x, y \in X \setminus \{0\}$ , then  $f$  satisfies the functional equation (1.2) on  $X \setminus \{0\}$ .

#### REFERENCES

- [1] M. Almahalebi, A. Charifi and S. Kabbaj, Hyperstability of a monomial functional equation, (preprint).
- [2] M. Almahalebi, A. Charifi and S. Kabbaj, Hyperstability of a Cauchy functional equation, (preprint).
- [3] M. Almahalebi and S. Kabbaj, Hyperstability of Cauchy-Jensen type functional equation. (preprint).
- [4] T. Aoki, On the stability of the linear transformation in Banach spaces, *J. Math. Soc. Japan* **2** (1950), 64-66.
- [5] A. Bahyrycz and M. Piszczek, Hyperstability of the Jensen functional equation, *Acta Math. Hungar.* **142** (2014), 353-365.
- [6] D. G. Bourgin, Approximately isometric and multiplicative transformations on continuous function rings, *Duke Math. J.* **16** (1949), 385-397.
- [7] J. Brzdęk, Hyperstability of the Cauchy equation on restricted domains. *Acta Math. Hungar.* **141** (2013), 58-67.
- [8] J. Brzdęk, Remarks on hyperstability of the Cauchy functional equation, *Aequationes Math.* **86** (2013), 255-267.
- [9] J. Brzdęk, A hyperstability result for the Cauchy equation, *Bull. Aust. Math. Soc.* **89** (2014), 33-40.
- [10] J. Brzdęk, J. Chudziak and Zs. Páles, A fixed point approach to stability of functional equations, *Nonlinear Anal.* **74** (2011), 6728-6732.
- [11] J. Brzdęk and K. Cieplinski, Hyperstability and superstability, *Abs. Appl. Anal.* **2013**, Article ID 401756, 13 pp. (2013).

- [12] L. Cădariu, L. Găvruta and P. Găvruta, On the stability of an affine functional equation, *J. Nonlinear Sci. Appl.* **6** (2013), 60-67.
- [13] L. Cădariu and V. Radu, Fixed points and the stability of Jensen's functional equation, *J. Inequal. Pure Appl. Math.* **4** (2003), no.1, Article 4.
- [14] A. Chahbi and N. Bounader, On the generalized stability of d'Alembert functional equation, *J. Nonlinear Sci. Appl.* **6** (2013), 198-204.
- [15] M. Eshaghi Gordji, H. Khodaei and M. Kamyar, Stability of Cauchy-Jensen type functional equation in generalized fuzzy normed spaces, *Comput. Math. Appl.* **62** (2011), 2950-2960.
- [16] G. Z. Eskandani and P. Găvruta, Hyers-Ulam-Rassias stability of pexiderized Cauchy functional equation in 2-Banach spaces, *J. Nonlinear Sci. Appl.* **5** (2012), 459-465.
- [17] P. Găvruta, A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings. *J. Math. Anal. Appl.* **184** (1994), 431-436.
- [18] E. Gselmann, Hyperstability of a functional equation, *Acta Math. Hungar.* **124** (2009), 179-188.
- [19] D. H. Hyers, On the stability of the linear functional equation, *Proc. Nat. Acad. Sci. U. S. A.* **27** (1941), 222-224.
- [20] K. Jun, H. Kim and J.M. Rassias, Extended Hyers-Ulam stability for Cauchy-Jensen mappings, *J. Difference Equ. Appl.* **13** (2007), 1139-1153.
- [21] S. Jung, Stability of the quadratic equation of Pexider type, *Abh. Math. Sem. Univ. Hamburg*, **70** (2000), 175-190.
- [22] H. A. Kenary, On the Hyers-Ulam-Rassias stability of a functional equation in non-Archimedean and random normed spaces, *Acta Univ. Apulensis, Math. Inform.* **27** (2011), 173-186.
- [23] H. A. Kenary, S. Y. Jang and C. Park, A fixed point approach to the Hyers-Ulam stability of a functional equation in various normed spaces, *Fixed Point Theory Appl.* **2011**, 2011:67, 14 pp. (2011).
- [24] Gy. Maksa and Zs. Páles, Hyperstability of a class of linear functional equations, *Acta Math. Acad. Paedag. Nyíregyháziensis*, **17** (2001), 107-112.
- [25] C. Park, Orthogonal stability of a cubic-quartic functional equation, *J. Nonlinear Sci. Appl.* **5** (2012), 28-36.
- [26] M. Piszczek, Remark on hyperstability of the general linear equation, *Aequationes Math.* (to appear).
- [27] J. M. Rassias, On approximation of approximately linear mappings by linear mappings, *J. Funct. Anal.* **46** (1982), 126-130.
- [28] J. M. Rassias, On the Ulam stability of Jensen and Jensen type mappings on restricted domains, *J. Math. Anal. Appl.* **281** (2003), 516-524.
- [29] Th. M. Rassias, On the stability of linear mapping in Banach spaces, *Proc. Amer. Math. Soc.* **72** (1978), 297-300.
- [30] Th. M. Rassias, On a modified Hyers-Ulam sequence, *J. Math. Anal. Appl.* **158** (1991), 106-113.
- [31] K. Ravi, E. Thandapani and B. V. Senthil Kumar, Solution and stability of a reciprocal type functional equation in several variables, *J. Nonlinear Sci. Appl.* **7** (2014), 18-27.
- [32] M. Sirouni, M. Almahalebi and S. Kabbaj, On the hyperstability of the Jensen's functional equation in complete metric spaces. (preprint).
- [33] S. M. Ulam, *Problems in Modern Mathematics*, Wiley, New York, 1960.
- [34] C. Zaharia, On the probabilistic stability of the monomial functional equation, *J. Nonlinear Sci. Appl.* **6** (2013), 51-59.

## A fractional finite difference inclusion

Dumitru Baleanu<sup>1,2</sup>, Shahram Rezapour<sup>3</sup>, Saeid Salehi<sup>3</sup>

<sup>1</sup>Department of Mathematics, Cankaya University,

Ogretmenler Cad. 14 06530, Balgat, Ankara, Turkey, email:dumitru@cankaya.edu.tr

<sup>2</sup>Institute of Space Sciences, Magurele, Bucharest, Romania

<sup>3</sup>Department of Mathematics, Azarbaijan Shahid Madani University,Iran

**Abstract.** In this manuscript we investigated the fractional finite difference inclusion  $\Delta_{\mu-2}^\mu x(t) \in F(t, x(t), \Delta x(t))$  via the boundary conditions  $\Delta x(b + \mu) = A$  and  $x(\mu - 2) = B$ , where  $1 < \mu \leq 2$ ,  $A, B \in \mathbb{R}$  and  $F : \mathbb{N}_{\mu-2}^{b+\mu+2} \times \mathbb{R} \times \mathbb{R} \rightarrow 2^{\mathbb{R}}$  is a compact valued multifunction.

**Keywords:** Fixed point of multifunction, Fractional finite difference inclusion, Hausdorff metric.

### 1 Introduction

We recall that there are several published works on the existence of solutions for some fractional finite difference equations. In [1, 3, 7] the boundary value problems for discrete fractional equations were explain in detail. In [4] the discrete nabla fractional Taylor formulae were described together with some well known inequalities. In [8] the authors investigated a k-dimensional system of fractional finite difference equations. The papers [11, 12, 13, 14] reported several results on discrete boundary value problem and existence results for fractional difference equations. Further results can be seen in [17, 19, 20, 21] and the references therein. The readers can find more details about elementary notions and definitions of fractional finite difference equations in [5, 6, 10, 15] and [16].

In [2] it was proved the existence of solutions for nonlinear fractional q-difference inclusions involving convex as well as non-convex valued maps with nonlocal Robin (separated) conditions. However, in the of our knowledge there is no research on fractional finite difference inclusions so far. Here, we give a motivation about the importance of studying the fractional finite difference inclusions. For example, consider the fractional finite difference equation  $\Delta^\mu y(t) = h(t + \mu - 2, y(t + \mu - 2))$  via the boundary conditions  $y(\mu - 3) = 0$ ,  $\Delta^\alpha y(\mu - 1 - \alpha) = 0$  and  $\Delta^\beta y(\mu + b + 1 - \beta) = 0$ , where  $t \in \mathbb{N}_0^{b+3}$ ,  $b \in \mathbb{N}_0$ ,  $2 < \mu \leq 3$ ,  $1 < \alpha \leq 2$ ,  $0 < \beta \leq 1$  and  $h : \mathbb{N}_{\mu-2}^{b+\mu+1} \times \mathbb{R} \rightarrow \mathbb{R}$  is a map. Define the compact valued multifunction  $T_h : \mathbb{N}_{\mu-2}^{b+\mu+1} \times \mathbb{R} \rightarrow 2^{\mathbb{R}}$  by  $T_h(t) = \{h(t + \mu - 2, y(t + \mu - 2))\}$ . It is easy to check that each solution of the fractional finite difference equation  $\Delta^\mu y(t) = h(t + \mu - 2, y(t + \mu - 2))$  is a solution for the fractional finite difference inclusion  $\Delta^\mu y(t) \in T_h(t)$ . Thus, studying the fractional finite difference inclusions needs more potential mathematical abilities. In this paper,

we provide some preliminaries to investigate the existence of solution for a fractional finite difference inclusion.

We recall some basic definitions utilized in the rest of the manuscript. The Gamma function is defined by  $\Gamma(z) = \int_0^\infty e^{-t}t^{z-1}dt$  for the complex numbers  $z$  in which the real part of  $z$  is positive (see [18]). Now, we define  $t^\mu := \frac{\Gamma(t+1)}{\Gamma(t+1-\mu)}$  for all  $t, \mu \in \mathbb{R}$  whenever the right-hand side is defined (see [7] and [16]). In this paper, we use the notations  $\mathbb{N}_p = \{p, p + 1, p + 2, \dots\}$  for all  $p \in \mathbb{R}$  and  $\mathbb{N}_p^q = \{p, p + 1, p + 2, \dots, q\}$  for all real numbers  $p$  and  $q$  whenever  $q - p$  is a natural number. Let  $\mu > 0$  with  $m - 1 < \mu < m$  for some natural number  $m$ . The  $\mu$ -th fractional sum of  $f$  based at  $a$  is defined by  $\Delta_a^{-\mu}f(t) = \frac{1}{\Gamma(\mu)} \sum_{k=a}^{t-\mu} (t - \sigma(k))^{\mu-1}f(k)$  for all  $t \in \mathbb{N}_{a+\mu}$ , where  $\sigma(k) = k + 1$  is the forward jump operator (see [3, 8]). Similarly, we define  $\Delta_a^\mu f(t) = \frac{1}{\Gamma(-\mu)} \sum_{k=a}^{t+\mu} (t - \sigma(k))^{-\mu-1}f(k)$  for all  $t \in \mathbb{N}_{a+m-\mu}$  (see [7, 16]). Note that, the domain of  $\Delta_a^r f$  is  $\mathbb{N}_{a+m-r}$  for  $r > 0$  and  $\mathbb{N}_{a-r}$  for  $r < 0$ . Also, for the natural number  $\mu = m$ , we have the known formula  $\Delta_a^\mu f(t) = \Delta^m f(t) = \sum_{i=0}^m (-1)^i \binom{m}{i} f(t + m - i)$  (see [7, 16]). Finally, set the trivial sum  $\Delta_a^0 f(t) = f(t)$  for all  $t \in \mathbb{N}_a$ .

**Lemma 1.1.** [7] *Let  $m \geq 1$ ,  $m - 1 < \mu \leq m$  and  $h : \mathbb{N}_{\mu-2}^{b+\mu+2} \rightarrow \mathbb{R}$  be a map. The general solution of the equation  $\Delta_{\mu-2}^\mu y(t) = h(t)$  is given by*

$$y(t) = \sum_{i=1}^m c_i t^{\mu-i} + \frac{1}{\Gamma(\mu)} \sum_{s=0}^{t-\mu} (t - \sigma(s))^{\mu-1} h(s)$$

for all  $t \in \mathbb{N}_{\mu-2}$ , where  $c_1, \dots, c_m$  are arbitrary constants.

Let  $(\mathcal{X}, d)$  be a metric space. The Hausdorff metric  $H_d : 2^{\mathcal{X}} \times 2^{\mathcal{X}} \rightarrow [0, \infty)$  is defined by

$$H_d(C, E) = \max\{\sup_{c \in C} d(c, E), \sup_{e \in E} d(C, e)\},$$

where  $d(C, e) = \inf_{c \in C} d(c, e)$  ([18]). We denote the set of compact subsets of  $\mathcal{X}$  by  $P_{cp}(\mathcal{X})$  and the set of closed subsets of  $\mathcal{X}$  by  $C(\mathcal{X})$ . Let  $T : \mathcal{X} \rightarrow 2^{\mathcal{X}}$  be a multifunction. An element  $x \in \mathcal{X}$  is called a fixed point of  $T$  whenever  $x \in Tx$  ([18]). A multifunction  $T : \mathcal{X} \rightarrow C(\mathcal{X})$  is called a contraction whenever there exists  $\lambda \in (0, 1)$  such that  $H_d(T(x), T(y)) \leq \lambda d(x, y)$  for all  $x, y \in \mathcal{X}$  ([18]). In 1970, Covitz and Nadler proved next theorem ([9]).

**Theorem 1.2.** *Each closed valued contraction multifunction on a complete metric space has a fixed point.*

## 2 Main result

Now, we are ready to investigate the existence of solutions for fractional finite difference inclusion

$$\Delta_{\mu-2}^\mu x(t) \in F(t, x(t), \Delta x(t)) \tag{1}$$

via the boundary conditions  $\Delta x(b + \mu) = A$  and  $x(\mu - 2) = B$ , where  $x : \mathbb{N}_{\mu-2}^{b+\mu+2} \rightarrow \mathbb{R}$  is a map,  $1 < \mu \leq 2$  and  $F : \mathbb{N}_{\mu-2}^{b+\mu+2} \times \mathbb{R} \times \mathbb{R} \rightarrow 2^{\mathbb{R}}$  is a compact valued multifunction.

**Lemma 2.1.** *Let  $1 < \mu \leq 2$  and  $x : \mathbb{N}_{\mu-2}^{b+\mu+2} \rightarrow \mathbb{R}$  and  $y : \mathbb{N}_0^{b+2} \rightarrow \mathbb{R}$  be two maps. Then  $x_0$  is a solution of the fractional finite difference equation  $\Delta_{\mu-2}^{\mu} x(t) = y(t)$  via the boundary conditions  $\Delta x(b + \mu) = A$  and  $x(\mu - 2) = B$  if and only if  $x_0$  is a solution of the fractional sum equation  $x(t) = \varphi(t) + \sum_{s=0}^{b+1} G(t, s)y(s)$ , where  $\varphi(t) = \left[ \frac{A}{(\mu-1)(\mu+b)^{\underline{\mu-2}}} - \frac{B(\mu-2)}{(b+3)\Gamma(\mu)} \right] t^{\underline{\mu-1}} + \frac{B}{\Gamma(\mu-1)} t^{\underline{\mu-2}}$ ,  $G(t, s) = -\frac{t^{\underline{\mu-1}}(b + \mu - \sigma(s))^{\underline{\mu-2}}}{(\mu + b)^{\underline{\mu-2}}} + (t - \sigma(s))^{\underline{\mu-1}}$  whenever  $0 \leq s \leq t - \mu \leq b + 1$  and  $G(t, s) = -\frac{t^{\underline{\mu-1}}(b + \mu - \sigma(s))^{\underline{\mu-2}}}{(\mu + b)^{\underline{\mu-2}}}$  whenever  $0 \leq t - \mu < s \leq b + 1$ .*

*Proof.* Let  $x_0$  be a solution of the equation  $\Delta_{\mu-2}^{\mu} x(t) = y(t)$  via the boundary conditions  $\Delta x(b + \mu) = A$  and  $x(\mu - 2) = B$ . By using Lemma 1.1, we get

$$x_0(t) = c_1 t^{\underline{\mu-1}} + c_2 t^{\underline{\mu-2}} + \frac{1}{\Gamma(\mu)} \sum_{s=0}^{t-\mu} (t - \sigma(s))^{\underline{\mu-1}} y(s)$$

where  $c_1, c_2 \in \mathbb{R}$ . By using the boundary condition  $x_0(\mu - 2) = B$ , we obtain  $c_2 = \frac{B}{\Gamma(\mu-1)}$ . Since  $\Delta x_0(t) = c_1(\mu - 1)t^{\underline{\mu-2}} + c_2(\mu - 2)t^{\underline{\mu-3}} + \frac{1}{\Gamma(\mu-1)} \sum_{s=0}^{t-\mu+1} (t - \sigma(s))^{\underline{\mu-2}} y(s)$ , by using the boundary condition  $\Delta x_0(b + \mu) = A$  we get

$$c_1 = \frac{A}{(\mu - 1)(\mu + b)^{\underline{\mu-2}}} - \frac{B(\mu - 2)}{(b + 3)\Gamma(\mu)} - \frac{1}{(\mu + b)^{\underline{\mu-2}}\Gamma(\mu)} \sum_{s=0}^{b+1} (b + \mu - \sigma(s))^{\underline{\mu-2}} y(s).$$

Hence,

$$\begin{aligned} x_0(t) &= \left[ \frac{A}{(\mu - 1)(\mu + b)^{\underline{\mu-2}}} - \frac{B(\mu - 2)}{(b + 3)\Gamma(\mu)} \right] t^{\underline{\mu-1}} + \frac{B}{\Gamma(\mu - 1)} t^{\underline{\mu-2}} \\ &- \frac{t^{\underline{\mu-1}}}{(\mu + b)^{\underline{\mu-2}}\Gamma(\mu)} \sum_{s=0}^{b+1} (b + \mu - \sigma(s))^{\underline{\mu-2}} y(s) + \frac{1}{\Gamma(\mu)} \sum_{s=0}^{t-\mu} (t - \sigma(s))^{\underline{\mu-1}} y(s) = \varphi(t) + \sum_{s=0}^{b+1} G(t, s)y(s). \end{aligned}$$

Now let  $x_0$  be a solution of the equation  $x(t) = \varphi(t) + \sum_{s=0}^{b+1} G(s, t)y(s)$ . Then, we have

$$\begin{aligned} x_0(t) &= \left[ \frac{A}{(\mu - 1)(\mu + b)^{\underline{\mu-2}}} - \frac{B(\mu - 2)}{(b + 3)\Gamma(\mu)} \right] t^{\underline{\mu-1}} + \frac{B}{\Gamma(\mu - 1)} t^{\underline{\mu-2}} \\ &- \frac{t^{\underline{\mu-1}}}{(\mu + b)^{\underline{\mu-2}}\Gamma(\mu)} \sum_{s=0}^{b+1} (b + \mu - \sigma(s))^{\underline{\mu-2}} y(s) + \frac{1}{\Gamma(\mu)} \sum_{s=0}^{t-\mu} (t - \sigma(s))^{\underline{\mu-1}} y(s). \end{aligned}$$

Since  $(\mu - 2)^{\underline{\mu-1}} = 0$ ,  $(\mu - 2)^{\underline{\mu-2}} = \Gamma(\mu - 1)$  and  $\sum_{s=0}^{-2} (\mu - 2 - \sigma(s))^{\underline{\mu-1}} y(s) = 0$ , we get  $x_0(\mu - 2) = B$ . A simple calculation shows that  $\Delta x_0(\mu + b) = A$ . On the other hand, it is easy to check that  $x_0(t) = c_1 t^{\underline{\mu-1}} + c_2 t^{\underline{\mu-2}} + \frac{1}{\Gamma(\mu)} \sum_{s=0}^{t-\mu} (t - \sigma(s))^{\underline{\mu-1}} y(s)$  is a solution for the equation  $\Delta_{\mu-2}^{\mu} x(t) = y(t)$  and so  $\Delta^{\mu} x_0(t) = y(t)$ . This completes the proof.  $\square$

A function  $x : \mathbb{N}_{\mu-2}^{b+\mu+2} \rightarrow \mathbb{R}$  is a solution for the problem (1) whenever it satisfies the boundary conditions and there exists a function  $y : \mathbb{N}_0^{b+2} \rightarrow \mathbb{R}$  such that  $y(t) \in F(t, x(t), \Delta x(t))$  for all  $t \in \mathbb{N}_0^{b+2}$  and

$$x(t) = \left[ \frac{A}{(\mu-1)(\mu+b)^{\underline{\mu-2}}} - \frac{B(\mu-2)}{(b+3)\Gamma(\mu)} \right] t^{\underline{\mu-1}} + \frac{B}{\Gamma(\mu-1)} t^{\underline{\mu-2}} - \frac{t^{\underline{\mu-1}}}{(\mu+b)^{\underline{\mu-2}}\Gamma(\mu)} \sum_{s=0}^{b+1} (b+\mu-\sigma(s))^{\underline{\mu-2}} y(s) + \frac{1}{\Gamma(\mu)} \sum_{s=0}^{t-\mu} (t-\sigma(s))^{\underline{\mu-1}} y(s).$$

Let  $\mathcal{X}$  be the set of all functions  $x : \mathbb{N}_{\mu-2}^{b+\mu+2} \rightarrow \mathbb{R}$  endowed with the norm

$$\|x\| = \max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} |x(t)| + \max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} |\Delta x(t)|.$$

Suppose that  $\{x_n\}$  is a Cauchy sequence in  $\mathcal{X}$ . For each  $\epsilon > 0$ , choose a natural number  $N$  such that  $\|x_n - x_m\| < \epsilon$  for all  $m, n > N$ . Hence,  $\max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} |x_n(t) - x_m(t)| < \epsilon$  and  $\max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} |\Delta x_n(t) - \Delta x_m(t)| < \epsilon$ . Choose  $x(t), z(t) \in \mathbb{R}$  such that  $x_n(t) \rightarrow x(t)$  and  $\Delta x_n(t) \rightarrow z(t)$  for all  $t \in \mathbb{N}_{\mu-2}^{b+\mu+2}$ . Note that,  $\Delta x_n(t) = x_n(t+1) - x_n(t)$  for all  $n$  and so  $\Delta x(t) = x(t+1) - x(t) = z(t)$ . This implies that there exists a natural number  $M$  such that  $|x_n(t) - x(t)| < \frac{\epsilon}{2}$  and  $|\Delta x_n(t) - \Delta x(t)| < \frac{\epsilon}{2}$  for all  $t \in \mathbb{N}_{\mu-2}^{b+\mu+2}$  and  $n > M$ . Thus,

$$\|x_n - x\| = \max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} |x_n(t) - x(t)| + \max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} |\Delta x_n(t) - \Delta x(t)| < \epsilon.$$

This shows that  $(\mathcal{X}, \|\cdot\|)$  is a Banach space. For each  $x \in \mathcal{X}$ , put

$$S_{F,x} = \{y : \mathbb{N}_0^{b+2} \rightarrow \mathbb{R} : y(t) \in F(t, x(t), \Delta x(t)) \text{ for all } t \in \mathbb{N}_0^{b+2}\}.$$

Note that, the selection principle implies that  $S_{F,x}$  is nonempty because  $F(t, x(t), \Delta x(t)) \neq \emptyset$ . Put

$$G_1 = \max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} \frac{1}{(\mu+b)^{\underline{\mu-2}}\Gamma(\mu)} \sum_{s=0}^{b+2} \left[ (b+\mu-\sigma(s))^{\underline{\mu-2}} t^{\underline{\mu-1}} + (\mu+b)^{\underline{\mu-2}} (t-\sigma(s))^{\underline{\mu-1}} \right]$$

and

$$G_2 = \max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} \frac{1}{(\mu+b)^{\underline{\mu-2}}\Gamma(\mu-1)} \sum_{s=0}^{b+2} \left[ (b+\mu-\sigma(s))^{\underline{\mu-2}} t^{\underline{\mu-2}} + (\mu+b)^{\underline{\mu-2}} (t-\sigma(s))^{\underline{\mu-2}} \right].$$

Since every Green function is bounded and the summations are finite,  $G_1$  and  $G_2$  are real numbers.

**Theorem 2.2.** Let  $g : \mathbb{N}_{\mu-2}^{b+\mu+2} \rightarrow \mathbb{R}$  be a map such that  $0 < \max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} |g(t)|(G_1 + G_2) < 1$ . Suppose that  $F : \mathbb{N}_{\mu-2}^{b+\mu+2} \times \mathbb{R} \times \mathbb{R} \rightarrow P_{cp}(\mathbb{R})$  is a multifunction such that

$$H_d(F(t, x_1, x_2), F(t, z_1, z_2)) \leq g(t)(|x_1 - z_1| + |x_2 - z_2|) \tag{2}$$

for all  $t \in \mathbb{N}_{\mu-2}^{b+\mu+2}$  and  $x_1, x_2, z_1, z_2 \in \mathbb{R}$ . Then the inclusion problem (1) has a solution.

*Proof.* Choose  $y \in S_{F,x}$ . Define  $h \in \mathcal{X}$  by

$$h(t) = \left[ \frac{A}{(\mu-1)(\mu+b)^{\mu-2}} - \frac{B(\mu-2)}{(b+3)\Gamma(\mu)} \right] t^{\mu-1} + \frac{B}{\Gamma(\mu-1)} t^{\mu-2} - \frac{t^{\mu-1}}{(\mu+b)^{\mu-2}\Gamma(\mu)} \sum_{s=0}^{b+1} (b+\mu-\sigma(s))^{\mu-2} y(s) + \frac{1}{\Gamma(\mu)} \sum_{s=0}^{t-\mu} (t-\sigma(s))^{\mu-1} y(s)$$

for all  $t \in \mathbb{N}_{\mu-2}^{\mu+b+2}$ . This shows that the set

$$\left\{ h \in \mathcal{X} : \text{there exists } y \in S_{F,x} \text{ such that } h(t) = w(t) \text{ for all } t \in \mathbb{N}_{\mu-2}^{\mu+b+2} \right\}$$

is nonempty, where

$$w(t) = \left[ \frac{A}{(\mu-1)(\mu+b)^{\mu-2}} - \frac{B(\mu-2)}{(b+3)\Gamma(\mu)} \right] t^{\mu-1} + \frac{B}{\Gamma(\mu-1)} t^{\mu-2} - \frac{t^{\mu-1}}{(\mu+b)^{\mu-2}\Gamma(\mu)} \sum_{s=0}^{b+1} (b+\mu-\sigma(s))^{\mu-2} y(s) + \frac{1}{\Gamma(\mu)} \sum_{s=0}^{t-\mu} (t-\sigma(s))^{\mu-1} y(s).$$

Now, define the multifunction  $T : \mathcal{X} \rightarrow 2^{\mathcal{X}}$  by

$$T(x) = \left\{ h \in \mathcal{X} : \text{there exists } y \in S_{F,x} \text{ such that } h(t) = w(t) \text{ for all } t \in \mathbb{N}_{\mu-2}^{\mu+b+2} \right\}.$$

First, we show that  $T(x)$  is a closed subset of  $\mathcal{X}$  for all  $x \in \mathcal{X}$ . Let  $x \in \mathcal{X}$  and  $\{u_n\}_{n \geq 1}$  be a sequence in  $T(x)$  with  $u_n \rightarrow u$ . Choose  $y_n \in S_{F,x}$  such that

$$u_n(t) = \left[ \frac{A}{(\mu-1)(\mu+b)^{\mu-2}} - \frac{B(\mu-2)}{(b+3)\Gamma(\mu)} \right] t^{\mu-1} + \frac{B}{\Gamma(\mu-1)} t^{\mu-2} - \frac{t^{\mu-1}}{(\mu+b)^{\mu-2}\Gamma(\mu)} \sum_{s=0}^{b+1} (b+\mu-\sigma(s))^{\mu-2} y_n(s) + \frac{1}{\Gamma(\mu)} \sum_{s=0}^{t-\mu} (t-\sigma(s))^{\mu-1} y_n(s)$$

for all  $t \in \mathbb{N}_{\mu-2}^{b+\mu+2}$  and  $n \geq 1$ . Since  $F$  has compact values,  $\{y_n\}_{n \geq 1}$  has a subsequence which converges to some  $y : \mathbb{N}_0^{b+2} \rightarrow \mathbb{R}$ . We denote this subsequence again by  $\{y_n\}_{n \geq 1}$ . It is easy to check that  $y \in S_{F,x}$  and

$$u_n(t) \rightarrow u(t) = \left[ \frac{A}{(\mu-1)(\mu+b)^{\mu-2}} - \frac{B(\mu-2)}{(b+3)\Gamma(\mu)} \right] t^{\mu-1} + \frac{B}{\Gamma(\mu-1)} t^{\mu-2}$$

$$-\frac{t^{\mu-1}}{(\mu+b)^{\mu-2}\Gamma(\mu)} \sum_{s=0}^{b+1} (b+\mu-\sigma(s))^{\mu-2} y(s) + \frac{1}{\Gamma(\mu)} \sum_{s=0}^{t-\mu} (t-\sigma(s))^{\mu-1} y(s)$$

for all  $t \in \mathbb{N}_{\mu-2}^{b+\mu+2}$ . This implies that  $u \in T(x)$  and so the multifunction  $T$  has closed values. Now, we show that  $T$  is a contraction multifunction with the constant

$$\lambda = \max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} |g(t)|(G_1 + G_2) < 1.$$

Let  $x, z \in \mathcal{X}$ ,  $h_1 \in T(x)$  and  $h_2 \in T(z)$ . Choose  $y_1 \in S_{F,x}$  and  $y_2 \in S_{F,z}$  such that

$$h_1(t) = \left[ \frac{A}{(\mu-1)(\mu+b)^{\mu-2}} - \frac{B(\mu-2)}{(b+3)\Gamma(\mu)} \right] t^{\mu-1} + \frac{B}{\Gamma(\mu-1)} t^{\mu-2} - \frac{t^{\mu-1}}{(\mu+b)^{\mu-2}\Gamma(\mu)} \sum_{s=0}^{b+1} (b+\mu-\sigma(s))^{\mu-2} y_1(s) + \frac{1}{\Gamma(\mu)} \sum_{s=0}^{t-\mu} (t-\sigma(s))^{\mu-1} y_1(s)$$

and

$$h_2(t) = \left[ \frac{A}{(\mu-1)(\mu+b)^{\mu-2}} - \frac{B(\mu-2)}{(b+3)\Gamma(\mu)} \right] t^{\mu-1} + \frac{B}{\Gamma(\mu-1)} t^{\mu-2} - \frac{t^{\mu-1}}{(\mu+b)^{\mu-2}\Gamma(\mu)} \sum_{s=0}^{b+1} (b+\mu-\sigma(s))^{\mu-2} y_2(s) + \frac{1}{\Gamma(\mu)} \sum_{s=0}^{t-\mu} (t-\sigma(s))^{\mu-1} y_2(s)$$

for all  $t \in \mathbb{N}_{\mu-2}^{b+\mu+2}$ . Since

$$H_d(F(t, x(t), \Delta x(t)), F(t, z(t), \Delta z(t))) \leq g(t)(|x(t) - z(t)| + |\Delta x(t) - \Delta z(t)|),$$

we get  $|y_1(t) - y_2(t)| \leq g(t)(|x(t) - z(t)| + |\Delta x(t) - \Delta z(t)|)$  for all  $t \in \mathbb{N}_0^{b+2}$ . Hence,

$$|h_1(t) - h_2(t)| \leq \sum_{s=0}^{b+1} \frac{(b+\mu-\sigma(s))^{\mu-2} t^{\mu-1}}{\Gamma(\mu)(\mu+b)^{\mu-2}} |y_1(s) - y_2(s)| + \sum_{s=0}^{t-\mu} \frac{(t-\sigma(s))^{\mu-1}}{\Gamma(\mu)} |y_1(s) - y_2(s)|.$$

Since  $\sum_{s=t-\mu+1}^{b+2} \frac{(t-\sigma(s))^{\mu-1}}{\Gamma(\mu)} = 0$  and  $\frac{(b+\mu-\sigma(b+2))^{\mu-2}}{\Gamma(\mu)(\mu+b)^{\mu-2}} = 0$ , we obtain

$$|h_1(t) - h_2(t)| \leq \sum_{s=0}^{b+2} \frac{(b+\mu-\sigma(s))^{\mu-2} t^{\mu-1} + (\mu+b)^{\mu-2} (t-\sigma(s))^{\mu-1}}{(\mu+b)^{\mu-2}\Gamma(\mu)} |y_1(s) - y_2(s)|$$

and so

$$\begin{aligned} \max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} |h_1(t) - h_2(t)| &\leq G_1 \max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} |g(t)| \left( \max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} |x(t) - z(t)| + \max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} |\Delta x(t) - \Delta z(t)| \right) \\ &= G_1 \max_{t \in \mathbb{N}_{\mu-2}^{b+\mu+2}} |g(t)| \|x - z\|. \end{aligned}$$



Since

$$\Delta h_1(t) = \left[ \frac{A}{(\mu + b)^{\mu-2}} - \frac{B(\mu - 2)}{(b + 3)\Gamma(\mu - 1)} \right] t^{\mu-2} + \frac{B}{\Gamma(\mu - 2)} t^{\mu-3} - \frac{t^{\mu-2}}{(\mu + b)^{\mu-2}\Gamma(\mu - 1)} \sum_{s=0}^{b+1} (b + \mu - \sigma(s))^{\mu-2} y_1(s) + \frac{1}{\Gamma(\mu - 1)} \sum_{s=0}^{t-\mu+1} (t - \sigma(s))^{\mu-2} y_1(s),$$

we get

$$|\Delta h_1(t) - \Delta h_2(t)| \leq \sum_{s=0}^{b+2} \frac{(b + \mu - \sigma(s))^{\mu-2} t^{\mu-2} + (\mu + b)^{\mu-2} (t - \sigma(s))^{\mu-2}}{(\mu + b)^{\mu-2}\Gamma(\mu - 1)} |y_1(s) - y_2(s)|$$

and so

$$\begin{aligned} \max_{t \in \mathbb{N}_{\mu-2}^{\mu+b+2}} |\Delta h_1(t) - \Delta h_2(t)| &\leq G_2 \max_{t \in \mathbb{N}_{\mu-2}^{\mu+b+2}} |g(t)| \left( \max_{t \in \mathbb{N}_{\mu-2}^{\mu+b+2}} |x(t) - z(t)| + \max_{t \in \mathbb{N}_{\mu-2}^{\mu+b+2}} |\Delta x(t) - \Delta z(t)| \right) \\ &= G_2 \max_{t \in \mathbb{N}_{\mu-2}^{\mu+b+2}} |g(t)| \|x - z\|. \end{aligned}$$

Hence,  $\|h_1 - h_2\| \leq \max_{t \in \mathbb{N}_{\mu-2}^{\mu+b+2}} |g(t)| (G_1 + G_2) \|x - z\| = \lambda \|x - z\|$  for all  $x, z \in \mathcal{X}$ ,  $h_1 \in T(x)$  and  $h_2 \in T(z)$ . This implies that  $H_d(T(x), T(z)) \leq \lambda \|x - z\|$  for all  $x, z \in \mathcal{X}$  and so the multifunction  $T$  is a contraction with closed values. Now by using Lemma 1.2,  $T$  has a fixed point which is a solution for the inclusion problem (1).  $\square$

Now, we present an example to illustrate the problem.

**Example 2.1.** Consider the fractional finite difference inclusion

$$\Delta_{-0.5}^{1.5} x(t) \in \left[ 0, e^t + \frac{\sin(x(t))}{e^7} + 8t^2 - \frac{|\Delta x(t)|}{e^7 + |\Delta x(t)|} \right] \tag{3}$$

via the boundary conditions  $\Delta x(6.5) = 5$  and  $x(-0.5) = -15$ . Put  $\mu = 1.5$ ,  $b = 5$ ,  $A = 5$ ,  $B = -15$  and  $F(t, x_1, x_2) = \left[ 0, e^t + \frac{\sin x_1}{e^7} + 8t^2 - \frac{|x_2|}{e^7 + |x_2|} \right]$  for all  $t \in \mathbb{N}_{-0.5}^{8.5}$  and  $x_1, x_2 \in \mathbb{R}$ . Note that,  $e^t + \frac{\sin x_1}{e^7} + 8t^2 - \frac{|x_2|}{e^7 + |x_2|} > 0$  for all  $t \in \mathbb{N}_{-0.5}^{8.5}$  and  $x_1, x_2 \in \mathbb{R}$  and so  $F$  is a compact valued multifunction on  $\mathbb{N}_{-0.5}^{8.5} \times \mathbb{R} \times \mathbb{R}$ . Define  $g(t) = \frac{1}{165|t|}$ . Note that  $\max_{t \in \mathbb{N}_{-0.5}^{8.5}} |g(t)| = \frac{2}{165}$ ,

$$G_1 = \max_{t \in \mathbb{N}_{-0.5}^{8.5}} \frac{1}{(6.5)^{-0.5}\Gamma(1.5)} \sum_{s=0}^7 (5.5 - s)^{-0.5} t^{0.5} + (6.5)^{-0.5} (t - \sigma(s))^{0.5} \simeq 66.8457$$

and

$$G_2 = \max_{t \in \mathbb{N}_{-0.5}^{8.5}} \frac{1}{(6.5)^{-0.5}\Gamma(0.5)} \sum_{s=0}^7 (5.5 - s)^{-0.5} t^{-0.5} + (6.5)^{-0.5} (t - \sigma(s))^{-0.5} \simeq 15.3947.$$

Also,  $\lambda = \frac{2}{165}(66.8457 + 15.3947) < 1$ . On the other hand, we have

$$\sup_{b \in F(t, y_1, y_2)} d(F(t, x_1, x_2), b) = \frac{\sin y_1}{e^7} - \frac{|y_2|}{e^7 + |y_2|} - \frac{\sin x_1}{e^7} + \frac{|x_2|}{e^7 + |x_2|}$$

whenever  $\frac{\sin y_1}{e^7} - \frac{|y_2|}{e^7 + |y_2|} > \frac{\sin x_1}{e^7} - \frac{|x_2|}{e^7 + |x_2|}$  and  $\sup_{b \in F(t, y_1, y_2)} d(F(t, x_1, x_2), b) = 0$  otherwise. This implies that

$$\begin{aligned} H_d(F(t, x_1, x_2), F(t, y_1, y_2)) &\leq \left| \frac{\sin y_1}{e^7} - \frac{|y_2|}{e^7 + |y_2|} - \frac{\sin x_1}{e^7} + \frac{|x_2|}{e^7 + |x_2|} \right| \\ &\leq \frac{1}{e^7} |\sin y_1 - \sin x_1| + \frac{1}{e^7} |y_2 - x_2| \leq g(t)(|x_1 - y_1| + |x_2 - y_2|) \end{aligned}$$

for all  $t \in \mathbb{N}_{-0.5}^{8.5}$  and  $x_1, x_2, y_1, y_2 \in \mathbb{R}$ . Now by using Theorem 2.2, the problem (3) has at least one solution.

**Remark 2.1.** The values  $A$  and  $B$  are arbitrary constants in the problem. In particular, the problem (3) has a solution via the boundary conditions  $\Delta x(6.5) = A$  and  $x(-0.5) = B$  for all  $A, B \in \mathbb{R}$ .

## Acknowledgments

Research of the authors Shahram Rezapour and Saeid Salehi was supported by Azarbaijan Shahid Madani University.

## References

- [1] N. Acar, F. M. Atici, *Exponential functions of discrete fractional calculus*, Appl. Anal. Discrete Math. 7 (2013) 343–353.
- [2] B. Ahmad, S.K. Ntouyas, *Existence of solutions for nonlinear fractional  $q$ -difference inclusions with nonlocal Robin (separated) conditions*, Mediteranean J. Math. 10 (2013) 1333-1351.
- [3] Kh. Alyousef, *Boundary value problems for discrete fractional equations*, Ph.D. Thesis, University of Nebraska-Lincoln (2012).
- [4] G. A. Anastassiou, *Nabla discrete fractional calculus and nabla inequalities*, Math. Computer Modelling 51 (2010) 562–571.
- [5] F. M. Atici, S. Sengul, *Modeling with fractional difference equations*, J. Math. Anal. Appl. 369 (2010) 1-9.
- [6] F. M. Atici, P. W. Eloe, *Initial value problems in discrete fractional calculus*, Ame. Math. Soc. 137 (2009) 981–989.

- [7] P. Awasthi, *Boundary value problems for discrete fractional equations*, Ph.D. Thesis, Ann Arbor, MI, University of Nebraska-Lincoln (2013).
- [8] D. Baleanu, Sh. Rezapour, S. Salehi, *A  $k$ -dimensional system of fractional finite difference equations*, Abst. Appl. Anal. (2014) Article ID 312578, 8 pages.
- [9] H. Covitz, S. Nadler, *Multivalued contraction mappings in generalized metric spaces*, Israel J. Math. 8 (1970) 5–11.
- [10] S. N. Elaydi, *An introduction to difference equations*, Springer-Verlag (1996).
- [11] C. S. Goodrich, *On a fractional boundary value problem with fractional boundary conditions*, Appl. Math. Let. (2012) 1101–1105.
- [12] C. S. Goodrich, *On discrete sequential fractional boundary value problems*, J. Math. Anal. Appl. 385 (2012) 111–124.
- [13] C. S. Goodrich, *Solutions to a discrete right-focal fractional boundary value problem*, Int. J. Diff. Eq. 5 (2010) 195–216.
- [14] C. S. Goodrich, *Some new existence results for fractional difference equations*, Int. J. Dynamical Syst. Diff. Eq. 3 (2011) 145–162.
- [15] M. Holm, *Sum and differences compositions in discrete fractional calculus*, CUBO. A Math. J. 13 (2011) 153–184.
- [16] M. Holm, *The theory of discrete fractional calculus: development and applications*, Ph.D. Thesis, Ann Arbor, MI, University of Nebraska-Lincoln (2011).
- [17] F. Jarad, K. Tas, *On Sumudu Transform Method in Discrete Fractional Calculus*, Abs. Appl. Anal. (2012) Article ID 270106, 16 pages.
- [18] M. Kisielewicz, *Differential Inclusions and Optimal Control*, Kluwer Academic, Dordrecht (1991).
- [19] Y. Lu, R. Ma, *Global structure of positive solutions for second-order difference equation with nonlinear boundary value condition*, Adv. Diff. Eq. (2014) 2014:188.
- [20] W. Lv, J. Feng, *Nonlinear discrete fractional mixed type sum-difference equation boundary value problems in Banach spaces*, Adv. Diff. Eq. (2014) 2014:184.
- [21] Y. Pan, Z. Han, S. Sun, Y. Zhao, *The existence of solutions to a system of discrete fractional boundary value problems*, Abs. Appl. Anal. (2012) 1-15.

# Some implicit properties of the second kind Bernoulli polynomials of order $\alpha$

C. S. Ryoo, J. Y. Kang

Department of Mathematics,  
Hannam University, Daejeon 306-791, Korea

**Abstract** In this paper we define the second kind Bernoulli polynomials of order  $\alpha$  in the complex plane and find some interesting properties, symmetric identities of this polynomials. We also derive some relations between the second kind Bernoulli polynomials of order  $\alpha$ , Euler polynomials of the second kind of order  $\alpha$ , the stirling numbers and other polynomials.

**2000 Mathematics Subject Classification** - 11B68, 11B73, 11S05

**Key words**- the second kind Bernoulli polynomials of order  $\alpha$ , Euler polynomials of the second kind of order  $\alpha$ , the stirling numbers, the stirling polynomials, central factorial numbers

## 1. Introduction

The Bernoulli, Euler and Genocchi polynomials have been a subject of investigations and have been used in various branches of mathematics such as theory of numbers, calculus of finite differences, combinatorial analysis,  $p$ -adic analytic number theory etc. These polynomials which are divided the first kind or the second kind have been researched by many mathematicians. Those polynomials have been extended and generalized in various direction, in particular, Bernoulli polynomials have been discussed by C. S. Ryoo, Q.-M. Luo, H. M. Srivastava, and T. Kim, etc(see [1-23]).

In this paper we use the following notations.  $\mathbb{N}$  denotes the set of natural numbers,  $\mathbb{Z}$  denotes the ring of integers,  $\mathbb{Q}$  denotes the field of rational numbers,  $\mathbb{C}$  denotes the set of complex numbers and  $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$ .

The second kind Bernoulli numbers  $\tilde{B}_n$  and polynomials  $\tilde{B}_n(x)$  are by means of the

following generating function (see [16-18]):

$$\begin{aligned} \sum_{n=0}^{\infty} \tilde{B}_n \frac{t^n}{n!} &= \frac{te^t}{e^{2t}-1} \quad (|t| < \pi), \\ \sum_{n=0}^{\infty} \tilde{B}_n(x) \frac{t^n}{n!} &= \left( \frac{te^t}{e^{2t}-1} \right) e^{tx} \quad (|t| < \pi), \end{aligned} \tag{1.1}$$

respectively. From the above generating function, we can easily see that

$$\tilde{B}_n := \tilde{B}_n(0).$$

Also, the Euler polynomials  $\tilde{E}_n(x)$  of second kinds are given by

$$\sum_{n=0}^{\infty} \tilde{E}_n(x) \frac{t^n}{n!} = \left( \frac{2e^t}{e^{2t}+1} \right) e^{tx} \quad (|t| < \frac{\pi}{2}). \tag{1.2}$$

In [7], they expanded the Euler polynomials using a real or complex parameter  $\alpha$  and introduced the Euler numbers and polynomials of the second kind of order  $\alpha$  as follows:

$$\begin{aligned} \sum_{n=0}^{\infty} \tilde{E}_n^{(\alpha)} \frac{t^n}{n!} &= \left( \frac{2e^t}{e^{2t}+1} \right)^\alpha \quad (|t| < \frac{\pi}{2}), \\ \sum_{n=0}^{\infty} \tilde{E}_n^{(\alpha)}(x) \frac{t^n}{n!} &= \left( \frac{2e^t}{e^{2t}+1} \right)^\alpha e^{tx} \quad (|t| < \frac{\pi}{2}). \end{aligned} \tag{1.3}$$

We usually define the central factorial numbers by the following expansion formula.

$$\sum_{k=0}^n T(n, k) x(x-1^2)(x-2^2) \cdots (x-(k-1)^2) = x^n. \tag{1.4}$$

or by the generating function

$$(2k)! \sum_{n=k}^{\infty} T(n, k) \frac{x^{2n}}{(2n)!} = (e^x + e^{-x} - 2)^k. \tag{1.5}$$

By using (1.4) and (1.5), we can find some properties of the central factorial numbers  $T(n, k)$  (see [8,23]).

The Stirling numbers of the second kind are the number of ways to partition a set of  $n$  objects into  $k$  non-empty subsets and are denoted by  $S_2(n, k)$ . The Stirling numbers of each kind according to the parameters  $n, k$  can form mutually inverse triangular matrices. The Stirling numbers of the second kind  $S_2(n, k)$  can be defined by means of

$$x^n = \sum_{k=0}^n S_2(n, k) x(x-1)(x-2) \cdots (x-n+1), \tag{1.6}$$

or by the generating function

$$(e^x - 1)^k = k! \sum_{n=k}^{\infty} S_2(n, k) \frac{x^n}{n!}. \tag{1.7}$$

It follows from (1.6) or (1.7) that

$$S_2(n, k) = S_2(n - 1, k - 1) + kS_2(n - 1, k),$$

with  $S_2(n, 0) = 0(n > 0)$ ,  $S_2(n, n) = 1$ ,  $S_2(n, 1) = 1(n > 0)$ ,  $S_2(n, k) = 0(k > n \text{ or } k < 0)$  (see [4,10,22-23]).

Associated Stirling numbers of the second kind  $b(n, k)$  are defined by

$$(e^x - 1 - x)^k = k! \sum_{n=2k}^{\infty} b(n, k) \frac{x^n}{n!}. \tag{1.8}$$

It follows from (1.8) that

$$b(n, k) = (n - 1)b(n - 2, k - 1) + kb(n - 1, k),$$

with  $b(n, 0) = 0(n > 0)$ ,  $b(0, 0) = 1$ ,  $b(n, 1) = 1(n > 1)$ ,  $b(n, k) = 0(2k > n \text{ or } k < 0)$ (see [3-4, 10-11]).

$S_n(x)$  is the stirling polynomials, defined by

$$\sum_{n=0}^{\infty} \frac{S_n(x)}{n!} t^n = \left( \frac{t}{1 - e^{-t}} \right)^{x+1} \tag{1.9}$$

The stirling polynomials are related to the stirling numbers of the first kind  $S_1(n, m)$  by

$$S_n(m) = \frac{(-1)^n}{\binom{m}{n}} S_1(m + 1, m - n + 1),$$

where  $\binom{m}{n}$  is a binomial coefficient and  $m$  is an integer with  $m \geq n$ (see [19]).

This paper is organized as follows. In section 2, we construct the second kind Bernoulli polynomials of order  $\alpha$  in the complex plane. We also study some interesting and basic properties of their polynomials. In section 3, we find some relations between the second kind Bernoulli polynomials of order  $\alpha$  and Euler polynomials of the second kind of order  $\alpha$ , the stirling numbers and other polynomials. We also define the power sum polynomials in order to derive some symmetric identities about the second kind Bernoulli polynomials of order  $\alpha$  in the complex plane.

2. Some properties of the second kind Bernoulli polynomials of order  $\alpha$

In this section, we construct the second kind Bernoulli polynomials of order  $\alpha$  in the complex plane. We get some basic properties of their polynomials.

**Definition 2.1.** For a real or complex parameter  $\alpha$ , the second kind Bernoulli numbers  $\tilde{B}_n^{(\alpha)}$  and the second kind Bernoulli polynomials  $\tilde{B}_n^{(\alpha)}(x)$ , each of degree  $n$  in  $x$  as well as in  $\alpha$ , are defined by means of the following generating functions:

$$\sum_{n=0}^{\infty} \tilde{B}_n^{(\alpha)} \frac{t^n}{n!} = \left( \frac{te^t}{e^{2t}-1} \right)^\alpha,$$

and

$$\sum_{n=0}^{\infty} \tilde{B}_n^{(\alpha)}(x) \frac{t^n}{n!} = \left( \frac{te^t}{e^{2t}-1} \right)^\alpha e^{tx}, \quad (|t| < \pi; 1^\alpha := 1),$$

respectively. Note that

$$\tilde{B}_n^{(\alpha)}(0) := \tilde{B}_n^{(\alpha)}, \quad \tilde{B}_n^{(1)} = \tilde{B}_n(x), \quad \tilde{B}_n^{(0)}(x) = x^n, \quad (n \in \mathbb{N}_0).$$

From Definition 2.1, we can find the above polynomials by using computer.

$n$	$\tilde{B}_n$	$\tilde{B}_n^{(\alpha)}$	$\tilde{B}_n^{(\alpha)}(x)$
$n=0$	$\frac{1}{2}$	$\frac{1}{2^\alpha}$	$\frac{1}{2^\alpha}$
$n=1$	0	0	$\frac{1}{2^\alpha}x$
$n=2$	$-\frac{1}{6}$	$\frac{1}{2^\alpha} \left( -\frac{1}{3}\alpha \right)$	$\frac{1}{2^\alpha} \left( -\frac{1}{3}\alpha + x^2 \right)$
$n=3$	0	0	$\frac{1}{2^\alpha} (-\alpha x + x^3)$
$n=4$	$\frac{7}{30}$	$\frac{1}{2^\alpha} \left( \frac{1}{270}\alpha + \frac{1}{27}\alpha^2 \right)$	$\frac{1}{2^\alpha} \left( \frac{2}{15}\alpha + \frac{1}{3}\alpha^2 - 2\alpha x^2 + x^4 \right)$
$n=5$	0	0	$\frac{1}{2^\alpha} \left( \left( \frac{2}{3}\alpha + \frac{5}{3}\alpha^2 \right) x - \frac{10}{3}\alpha x^3 + x^5 \right)$
$n=6$	$-\frac{31}{42}$	$\frac{1}{2^\alpha} \left( -\frac{16}{63}\alpha - \frac{6}{9}\alpha^2 - \frac{5}{9}\alpha^3 \right)$	$\frac{1}{2^\alpha} \left( -\frac{16}{63}\alpha + \frac{10}{9}\alpha^2 - \frac{16}{9}\alpha^2 - \frac{5}{9}\alpha^3 + (2\alpha + 5\alpha^2)x^2 - 5\alpha x^4 + x^6 \right)$
...	...	...	...

Table 1: The comparison of  $\tilde{B}_n$ ,  $\tilde{B}_n^{(\alpha)}$  and  $\tilde{B}_n^{(\alpha)}(x)$

From now on, we will find the basic properties of the second kind Bernoulli polynomials of order  $\alpha$  from Definition 2.1.

**Theorem 2.2.** Let  $\alpha, \beta \in \mathbb{C}$ . Then we find that

$$\tilde{B}_n^{(\alpha+\beta)}(x+y) = \sum_{k=0}^n \binom{n}{k} \tilde{B}_k^{(\alpha)}(x) \tilde{B}_{n-k}^{(\beta)}(y).$$

*Proof.* From Definition 2.1, we obtain as the following equation by the binomial theorem.

$$\begin{aligned} \sum_{n=0}^{\infty} \tilde{B}_n^{(\alpha+\beta)}(x+y) \frac{t^n}{n!} &= \left( \frac{te^t}{e^{2t}-1} \right)^{\alpha+\beta} e^{t(x+y)} \\ &= \sum_{n=0}^{\infty} \tilde{B}_n^{(\alpha)}(x) \frac{t^n}{n!} \sum_{n=0}^{\infty} \tilde{B}_n^{(\beta)}(y) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^n \binom{n}{k} \tilde{B}_k^{(\alpha)}(x) \tilde{B}_{n-k}^{(\beta)}(y) \frac{t^n}{n!}. \end{aligned}$$

Comparing the both sides of  $\frac{t^n}{n!}$ , we complete the proof.

□

**Corollary 2.3.** Let  $\alpha, \beta \in \mathbb{C}$ . From Theorem 2.2, we can see that

(i) If we suppose  $\beta = 0$ , then

$$\tilde{B}_n^{(\alpha)}(x+y) = \sum_{k=0}^n \binom{n}{k} \tilde{B}_k^{(\alpha)}(x) y^{n-k} = \sum_{k=0}^n \binom{n}{k} \tilde{B}_k^{(\alpha)}(y) x^{n-k}.$$

(ii) If we suppose  $\beta = 0$  and  $y = 0$ , then

$$\tilde{B}_n^{(\alpha)}(x) = \sum_{k=0}^n \binom{n}{k} \tilde{B}_k^{(\alpha)} x^{n-k} = \sum_{k=0}^n \binom{n}{k} \tilde{B}_k^{(\alpha)} x^k.$$

**Theorem 2.4.** Let  $\alpha, x \in \mathbb{C}$  and  $n \in \mathbb{N}_0$ . Then we get that

$$\tilde{B}_n^{(\alpha)}(-x) = (-1)^n \tilde{B}_n^{(\alpha)}(x).$$

*Proof.* Substituting  $-x$  instead of  $x$  we get the following equation.

$$\begin{aligned} \sum_{n=0}^{\infty} \tilde{B}_n^{(\alpha)}(-x) \frac{t^n}{n!} &= \left( \frac{te^t}{e^{2t}-1} \right)^{\alpha} e^{-tx} \\ &= \left( \frac{-te^{-t}}{e^{-2t}-1} \right)^{\alpha} e^{-tx} \\ &= \sum_{n=0}^{\infty} (-1)^n \tilde{B}_n^{(\alpha)}(x) \frac{t^n}{n!}. \end{aligned}$$

Hence the proof of Theorem 2.4. is clear.



□

**Theorem 2.5.** Let  $\alpha \in \mathbb{C}$  and  $n \in \mathbb{N}_0$ . Then we have

$$\tilde{B}_n^{(\alpha)}(x+2) - \tilde{B}_n^{(\alpha)}(x) = n\tilde{B}_n^{(\alpha-1)}(x+1) = \sum_{k=0}^n \binom{n}{k} k\tilde{B}_{n-k}^{(\alpha-1)}(x).$$

*Proof.* By using Definition 2.1, we get that

$$\begin{aligned} \tilde{B}_n^{(\alpha)}(x+2) - \tilde{B}_n^{(\alpha)}(x) \frac{t^n}{n!} &= \left(\frac{te^t}{e^{2t}-1}\right)^\alpha e^{t(x+2)} - \left(\frac{te^t}{e^{2t}-1}\right)^\alpha e^{tx} \\ &= \left(\frac{te^t}{e^{2t}-1}\right)^{\alpha-1} te^{t(x+1)}. \end{aligned}$$

We can find out the following equation.

$$\left(\frac{te^t}{e^{2t}-1}\right)^{\alpha-1} te^{t(x+1)} = \sum_{n=0}^{\infty} \tilde{B}_{n-1}^{(\alpha-1)}(x) \frac{t^n}{n!} \sum_{n=0}^{\infty} n \frac{t^n}{n!} = \sum_{n=0}^{\infty} \sum_{k=0}^n \binom{n}{k} k\tilde{B}_{n-k}^{(\alpha-1)}(x) \frac{t^n}{n!}.$$

We also find the other equation as the follows:

$$\left(\frac{te^t}{e^{2t}-1}\right)^{\alpha-1} te^{t(x+1)} = \sum_{n=0}^{\infty} \tilde{B}_n^{(\alpha-1)}(x+1) \frac{t^{n+1}}{n!} = \sum_{n=0}^{\infty} n\tilde{B}_{n-1}^{(\alpha-1)}(x+1) \frac{t^n}{n!}.$$

Hence we terminate the proof of Theorem 2.5.

□

**Theorem 2.6.** Let  $\alpha \in \mathbb{C}$  and  $n \in \mathbb{N}_0$ . Then we derive that

- (i)  $\sum_{k=0}^{n-1} \binom{n-1}{k} \tilde{B}_k^{(\alpha-1)}(x) = \frac{1}{n} [\tilde{B}_n^{(\alpha)}(x+2) - \tilde{B}_n^{(\alpha)}(x)].$
- (ii)  $\tilde{B}_n^{(\alpha-1)}(x+1) = \frac{1}{n+1} \left[ \sum_{k=0}^{n+1} 2^{n+1-k} \binom{n+1}{k} \tilde{B}_k^{(\alpha)}(x) - \tilde{B}_{n+1}^{(\alpha)}(x) \right].$
- (iii)  $\tilde{B}_n^{(\alpha)}(x) = \sum_{k=0}^n \binom{n}{k} [2^{n-k} \tilde{B}_k^{(\alpha)}(x) - k\tilde{B}_{n-k}^{(\alpha-1)}(x)].$

*Proof.* (i) Using Corollary 2.3.(i), we can see that

$$\tilde{B}_n^{(\alpha)}(x+1) = \sum_{k=0}^n \binom{n}{k} \tilde{B}_k^{(\alpha)}(x), \tag{2.1}$$

and

$$\tilde{B}_n^{(\alpha)}(x+2) = \sum_{k=0}^n \binom{n}{k} 2^{n-k} \tilde{B}_k^{(\alpha)}(x). \tag{2.2}$$

If we combine Theorem 2.5 and (2.1), then we find that

$$\tilde{B}_n^{(\alpha)}(x+2) - \tilde{B}_n^{(\alpha)}(x) = n \sum_{k=0}^{n-1} \binom{n-1}{k} \tilde{B}_k^{(\alpha-1)}(x).$$

Therefore, we can see Theorem 2.6.(i).

(ii) From (2.2) we can transform Theorem 2.5 as following equation:

$$\sum_{k=0}^n \binom{n}{k} 2^{n-k} \tilde{B}_k^{(\alpha)}(x) - \tilde{B}_n^{(\alpha)}(x) = n \tilde{B}_{n-1}^{(\alpha-1)}(x+1).$$

Hence we have

$$\tilde{B}_n^{(\alpha-1)}(x+1) = \frac{1}{n+1} \left[ \sum_{k=0}^{n+1} 2^{n+1-k} \binom{n+1}{k} \tilde{B}_k^{(\alpha)}(x) - \tilde{B}_{n+1}^{(\alpha)}(x) \right].$$

(iii) This proof is very similar to the proof of (ii). We find the below equation from Theorem 2.5 and (2.2).

$$\sum_{k=0}^n \binom{n}{k} 2^{n-k} \tilde{B}_k^{(\alpha)}(x) - \tilde{B}_n^{(\alpha)}(x) = \sum_{k=0}^n \binom{n}{k} k \tilde{B}_{n-k}^{(\alpha-1)}(x).$$

From the above equation we know that

$$\tilde{B}_n^{(\alpha)}(x) = \sum_{k=0}^n \binom{n}{k} \left[ 2^{n-k} \tilde{B}_k^{(\alpha)}(x) - k \tilde{B}_{n-k}^{(\alpha-1)}(x) \right].$$

Thus we wind up the proof of Theorem 2.6. □

**Corollary 2.7.** Let  $n$  be non-negative integer. From Theorem 2.4.(ii), we can see that

$$x^n = \frac{1}{n+1} \sum_{k=0}^{n+1} 2^{n+1-k} \binom{n+1}{k} \tilde{B}_k(x-1).$$

**Theorem 2.8.** For  $w_1, w_2$  be positive integers and  $\alpha \in \mathbb{C}$ , we have

$$\begin{aligned} & \sum_{n=0}^l \binom{l}{n} \sum_{i=0}^{w_2-1} \sum_{j=0}^{w_1-1} w_1^{l-n} w_2^n \tilde{B}_{l-n}^{(\alpha)}(w_2x + 2j \frac{w_2}{w_1}) \tilde{B}_n^{(\alpha)}(w_1y + 2i \frac{w_1}{w_2}) \\ &= \sum_{n=0}^l \binom{l}{n} \sum_{i=0}^{w_1-1} \sum_{j=0}^{w_2-1} w_2^{l-n} w_1^n \tilde{B}_{l-n}^{(\alpha)}(w_1x + 2j \frac{w_1}{w_2}) \tilde{B}_n^{(\alpha)}(w_2y + 2i \frac{w_2}{w_1}). \end{aligned}$$

*Proof.* We can consider that

$$A(t) := \frac{(e^{2w_1w_2t} - 1)^2 (w_1w_2t^2 e^{t(w_1+w_2)})^\alpha e^{w_1w_2t(x+y)}}{(e^{2w_1t} - 1)^{\alpha+1} (e^{2w_2t} - 1)^{\alpha+1}}.$$

Then we are able to express  $A(t)$  as follows:

$$\begin{aligned} A(t) &:= \left(\frac{w_1te^{w_1t}}{e^{2w_1t} - 1}\right)^\alpha e^{w_1w_2tx} \sum_{i=0}^{w_2-1} e^{2iw_1t} \left(\frac{w_2te^{w_2t}}{e^{2w_2t} - 1}\right)^\alpha e^{w_1w_2ty} \sum_{j=0}^{w_1-1} e^{2jw_2t} \\ &= \sum_{l=0}^{\infty} \sum_{n=0}^l \binom{l}{n} \sum_{i=0}^{w_2-1} \sum_{j=0}^{w_1-1} w_1^{l-n} w_2^n \tilde{B}_{l-n}^{(\alpha)}(w_2x + 2j\frac{w_2}{w_1}) \tilde{B}_n^{(\alpha)}(w_1y + 2i\frac{w_1}{w_2}) \frac{t^l}{l!}. \end{aligned} \tag{2.3}$$

We also represent  $A(t)$  as the following form.

$$\begin{aligned} A(t) &:= \left(\frac{w_2te^{w_2t}}{e^{2w_2t} - 1}\right)^\alpha e^{w_1w_2tx} \sum_{i=0}^{w_1-1} e^{2iw_2t} \left(\frac{w_1te^{w_1t}}{e^{2w_1t} - 1}\right)^\alpha e^{w_1w_2ty} \sum_{j=0}^{w_2-1} e^{2jw_1t} \\ &= \sum_{l=0}^{\infty} \sum_{n=0}^l \binom{l}{n} \sum_{i=0}^{w_1-1} \sum_{j=0}^{w_2-1} w_2^{l-n} w_1^n \tilde{B}_{l-n}^{(\alpha)}(w_1x + 2j\frac{w_1}{w_2}) \tilde{B}_n^{(\alpha)}(w_2y + 2i\frac{w_2}{w_1}) \frac{t^l}{l!}. \end{aligned} \tag{2.4}$$

By using the coefficient comparison in (2.3) and (2.4), we proved Theorem 2.8.

□

**Corollary 2.9.** By substituting  $w_1 = 1$  from Theorem 2.8, we easily see that

$$\begin{aligned} &\sum_{n=0}^l \binom{l}{n} \sum_{i=0}^{w_2-1} w_2^n \tilde{B}_{l-n}^{(\alpha)}(w_2x) \tilde{B}_n^{(\alpha)}(y + \frac{2i}{w_2}) \\ &= \sum_{n=0}^l \binom{l}{n} \sum_{j=0}^{w_2-1} w_2^{l-n} \tilde{B}_n^{(\alpha)}(w_2y) \tilde{B}_{l-n}^{(\alpha)}(x + \frac{2j}{w_2}). \end{aligned}$$

**Theorem 2.10.** For  $w_1, w_2$  be positive integers,  $\alpha \in \mathbb{C}$  and  $n \in \mathbb{N}$ , we obtain

$$\begin{aligned} &\sum_{l=0}^n \binom{n}{l} l w_1^{n+1-l} w_2^{l-1} \tilde{B}_{n-l}^{(\alpha-1)}(w_2x) \sum_{i=0}^{w_2-1} \tilde{B}_{l-1}^{(\alpha)}(w_1y + 2i\frac{w_1}{w_2} + \frac{w_1}{w_2}) \\ &= \sum_{l=0}^n \binom{n}{l} l w_2^{n+1-l} w_1^{l-1} \tilde{B}_{n-l}^{(\alpha-1)}(w_1x) \sum_{j=0}^{w_1-1} \tilde{B}_{l-1}^{(\alpha)}(w_2y + 2j\frac{w_2}{w_1} + \frac{w_2}{w_1}). \end{aligned}$$

*Proof.* Assume that

$$B(t) := \frac{(e^{2w_1w_2t} - 1)(w_1w_2t^2 e^{t(w_1+w_2)})^\alpha e^{w_1w_2t(x+y)}}{(e^{2w_1t} - 1)^\alpha (e^{2w_2t} - 1)^\alpha}.$$

Then we express  $B(t)$  as follows:

$$\begin{aligned}
 B(t) &:= \left(\frac{w_1 t e^{w_1 t}}{e^{2w_1 t} - 1}\right)^{\alpha-1} e^{w_1 w_2 t x} w_1 t e^{w_1 t} \sum_{i=0}^{w_2-1} e^{2i w_1 t} \left(\frac{w_2 t e^{w_2 t}}{e^{2w_2 t} - 1}\right)^{\alpha} e^{w_1 w_2 t y} \\
 &= \sum_{n=1}^{\infty} \sum_{l=0}^n \binom{n}{l} l \tilde{B}_{n-l}^{(\alpha-1)}(w_2 x) w_1^{n+1-l} w_2^{l-1} \sum_{i=0}^{w_2-1} \tilde{B}_{l-1}^{(\alpha)}\left(w_1 y + 2i \frac{w_1}{w_2} + \frac{w_1}{w_2}\right) \frac{t^n}{n!}.
 \end{aligned} \tag{2.5}$$

We can also represent  $B(t)$  as the following equation.

$$\begin{aligned}
 B(t) &:= \left(\frac{w_2 t e^{w_2 t}}{e^{2w_2 t} - 1}\right)^{\alpha-1} e^{w_1 w_2 t x} w_2 t e^{w_2 t} \sum_{j=0}^{w_1-1} e^{2j w_2 t} \left(\frac{w_1 t e^{w_1 t}}{e^{2w_1 t} - 1}\right)^{\alpha} e^{w_1 w_2 t y} \\
 &= \sum_{n=1}^{\infty} \sum_{l=0}^n \binom{n}{l} l \tilde{B}_{n-l}^{(\alpha-1)}(w_1 x) w_2^{n+1-l} w_1^{l-1} \sum_{j=0}^{w_1-1} \tilde{B}_{l-1}^{(\alpha)}\left(w_2 y + 2j \frac{w_2}{w_1} + \frac{w_2}{w_1}\right) \frac{t^n}{n!}.
 \end{aligned} \tag{2.6}$$

We get the following symmetric equation from (2.5) and (2.6).

$$\begin{aligned}
 &\sum_{l=0}^n \binom{n}{l} l w_1^{n+1-l} w_2^{l-1} \tilde{B}_{n-l}^{(\alpha-1)}(w_2 x) \sum_{i=0}^{w_2-1} \tilde{B}_{l-1}^{(\alpha)}\left(w_1 y + 2i \frac{w_1}{w_2} + \frac{w_1}{w_2}\right) \\
 &= \sum_{l=0}^n \binom{n}{l} l w_2^{n+1-l} w_1^{l-1} \tilde{B}_{n-l}^{(\alpha-1)}(w_1 x) \sum_{j=0}^{w_1-1} \tilde{B}_{l-1}^{(\alpha)}\left(w_2 y + 2j \frac{w_2}{w_1} + \frac{w_2}{w_1}\right).
 \end{aligned}$$

Thus, we complete the proof of Theorem 2.10. □

**Corollary 2.11.** When  $w_1 = 1$  from Theorem 2.10, we can see that

$$\begin{aligned}
 &\sum_{l=0}^n \binom{n}{l} l w_2^{l-1} \tilde{B}_{n-l}^{(\alpha-1)}(w_2 x) \sum_{i=0}^{w_2-1} \tilde{B}_{l-1}^{(\alpha)}\left(y + \frac{2i}{w_2} + \frac{1}{w_2}\right) \\
 &= \sum_{l=0}^n \binom{n}{l} l w_2^{n+1-l} \tilde{B}_{n-l}^{(\alpha-1)}(x) \tilde{B}_{l-1}^{(\alpha)}(w_2 y + w_2).
 \end{aligned}$$

3. Some relations of the second kind Bernoulli polynomials of order  $\alpha$ , the various kinds of numbers and polynomials

In this section, we investigate some interesting relations between the second kind Bernoulli polynomials of order  $\alpha$  and the Euler polynomials of the second kind of order  $\alpha$ . We find some relations between the second kind Bernoulli polynomials of order  $\alpha$ , stirring numbers, the associated stirring numbers of the second kind and central factorial numbers. We also derive the analogue of Srivastava-Pintér addition theorem(see [14,20]).

**Theorem 3.1.** Let  $n \in \mathbb{N}_0$  and  $\alpha \in \mathbb{C}$ . Then one has

$$\tilde{B}_n^{(\alpha)}(x) = \left(\frac{1}{2}\right)^n \sum_{k=0}^n \binom{n}{k} \tilde{B}_{n-k}^{(\alpha)}(x) \tilde{E}_k^{(\alpha)}(x) = \left(\frac{1}{2}\right)^n \sum_{k=0}^n \binom{n}{k} \tilde{B}_{n-k}^{(\alpha)} \tilde{E}_k^{(\alpha)}(2x).$$

*Proof.* We can find the following equation by using generating function.

$$\sum_{n=0}^{\infty} \tilde{B}_n^{(\alpha)}(x) \frac{t^n}{n!} = \left(\frac{te^t}{e^{2t}-1}\right)^\alpha e^{tx} = \left(\frac{te^{\frac{1}{2}t}}{e^t-1}\right)^\alpha \left(\frac{e^{\frac{1}{2}t}}{e^t+1}\right)^\alpha e^{tx}.$$

From the above equation we can combine the Euler polynomials of the second kind of order  $\alpha$ ,  $\tilde{E}_n^{(\alpha)}$ , by using Cauchy product as follows:

$$\sum_{n=0}^{\infty} \tilde{B}_n^{(\alpha)}(x) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \sum_{k=0}^n \binom{n}{k} \left(\frac{1}{2}\right)^n \tilde{B}_{n-k}^{(\alpha)} \tilde{E}_k^{(\alpha)}(2x) \frac{t^n}{n!}.$$

Hence, the proof of Theorem 3.1 is clear. □

From simple transformation, we are able to represent to the following equation from Theorem 3.1.

$$2^n \tilde{B}_n^{(\alpha)}\left(\frac{x}{2}\right) = \sum_{k=0}^n \binom{n}{k} \tilde{B}_{n-k}^{(\alpha)}\left(\frac{x}{2}\right) \tilde{E}_k^{(\alpha)}\left(\frac{x}{2}\right) = \sum_{k=0}^n \binom{n}{k} \tilde{B}_{n-k}^{(\alpha)} \tilde{E}_k^{(\alpha)}(x).$$

By using addition theorem of the Euler polynomials of the second kind of order  $\alpha$ , we find the analogue of Srivastava-Pintér addition theorem as the following theorem 3.2.

**Theorem 3.2.** Let  $\alpha \in \mathbb{C}$  and  $n \in \mathbb{N}_0$ . Then we obtain that

$$\tilde{B}_n^{(\alpha)}(x+y) = \frac{1}{2} \sum_{k=0}^n \binom{n}{k} \left[ \tilde{B}_k^{(\alpha)}(y+2) + \tilde{B}_k^{(\alpha)}(y) \right] \tilde{E}_{n-k}(x-1).$$

*Proof.* In [6], we can make the following equation by using simple calculation.

$$x^n = \frac{1}{2} \left[ \sum_{k=0}^n \binom{n}{k} 2^{n-k} \tilde{E}_k(x-1) + \tilde{E}_n(x-1) \right]. \tag{3.1}$$

Substituting (3.1) in Corollary 2.3.(i), inverting the order of summation and using the elementary combinatorial, we get

$$\begin{aligned} \tilde{B}_n^{(\alpha)}(x+y) &= \sum_{k=0}^n \binom{n}{k} \tilde{B}_k^{(\alpha)}(y) \left[ \frac{1}{2} \left\{ \sum_{j=0}^{n-k} \binom{n-k}{j} 2^{n-k-j} \tilde{E}_j(x-1) + \tilde{E}_{n-k}(x-1) \right\} \right] \\ &= \frac{1}{2} \left[ \sum_{j=0}^n \binom{n}{j} \tilde{E}_j(x-1) \sum_{k=0}^{n-j} \binom{n-j}{k} 2^{n-k-j} \tilde{B}_k^{(\alpha)}(y) + \sum_{k=0}^n \binom{n}{k} \tilde{B}_k^{(\alpha)}(y) \tilde{E}_{n-k}(x-1) \right]. \end{aligned}$$

Note that

$$\tilde{B}_{n-j}^{(\alpha)}(y+2) = \sum_{k=0}^{n-j} \binom{n-j}{k} 2^{n-k-j} \tilde{B}_k^{(\alpha)}(y).$$

By applying the above form, we derive that

$$\tilde{B}_n^{(\alpha)}(x+y) = \frac{1}{2} \sum_{k=0}^n \binom{n}{k} \left[ \tilde{B}_k^{(\alpha)}(y+2) + \tilde{B}_k^{(\alpha)}(y) \right] \tilde{E}_{n-k}(x-1).$$

Hence, we complete the proof of Theorem 3.2. □

**Corollary 3.3.** When  $\alpha = 1$  from Theorem 3.2, we see that

$$\tilde{B}_n(x+y) = \frac{1}{2} \sum_{k=0}^n \binom{n}{k} \left[ \tilde{B}_k(y+2) + \tilde{B}_k(y) \right] \tilde{E}_{n-k}(x-1).$$

**Corollary 3.4.** By setting  $y = 0$  from Theorem 3.2, we have

$$\tilde{B}_n^{(\alpha)}(x) = \frac{1}{2} \sum_{k=0}^n \binom{n}{k} \left[ \tilde{B}_k^{(\alpha)}(2) + \tilde{B}_k^{(\alpha)} \right] \tilde{E}_{n-k}(x-1).$$

**Theorem 3.5.** Let  $n \in \mathbb{N}_0$  and  $m \in \mathbb{N}$ . Then we get

$$\begin{aligned} \tilde{B}_n^{(m)} &= \frac{n!}{2^m} \sum_{\substack{v_1, \dots, v_m=0 \\ v_1 + \dots + v_m = n}} \frac{\left( \sum_{k=0}^{v_1} \binom{v_1}{k} \left( \tilde{B}_{v_1-k}(1) \tilde{E}_k + \tilde{B}_{v_1-k} \tilde{E}_k(-1) \right) \right) \cdots}{v_1 \cdots} \\ &\times \frac{\left( \sum_{k=0}^{v_m} \binom{v_m}{k} \left( \tilde{B}_{v_m-k}(1) \tilde{E}_k + \tilde{B}_{v_m-k} \tilde{E}_k(-1) \right) \right)}{v_m!}. \end{aligned}$$

*Proof.* Suppose that  $\alpha = 1$ . That is,

$$\begin{aligned} \sum_{n=0}^{\infty} \tilde{B}_n \frac{t^n}{n!} &= \left( \frac{te^t}{e^{2t}-1} \right) = \frac{te^{3t}}{(e^{2t}-1)(e^{2t}+1)} + \frac{te^t}{(e^{2t}-1)(e^{2t}+1)} \\ &= \sum_{n=0}^{\infty} \tilde{B}_n(1) \sum_{n=0}^{\infty} \frac{1}{2} \tilde{E}_n \frac{t^n}{n!} + \sum_{n=0}^{\infty} \tilde{B}_n \sum_{n=0}^{\infty} \frac{1}{2} \tilde{E}_n(-1) \frac{t^n}{n!} \\ &= \frac{1}{2} \sum_{n=0}^{\infty} \sum_{k=0}^n \binom{n}{k} \left( \tilde{B}_{n-k}(1) \tilde{E}_k + \tilde{B}_{n-k} \tilde{E}_k(-1) \right) \frac{t^n}{n!}. \end{aligned}$$

Consider that  $\alpha = 2$ . Then we have

$$\begin{aligned} \sum_{n=0}^{\infty} \tilde{B}_n^{(2)} \frac{t^n}{n!} &= \left( \frac{1}{2} \sum_{n=0}^{\infty} \left\{ \sum_{k=0}^n \binom{n}{k} \left( \tilde{B}_{n-k}(1)\tilde{E}_k + \tilde{B}_{n-k}\tilde{E}_k(-1) \right) \right\} \frac{t^n}{n!} \right) \\ &\quad \times \left( \frac{1}{2} \sum_{n=0}^{\infty} \left\{ \sum_{k=0}^n \binom{n}{k} \left( \tilde{B}_{n-k}(1)\tilde{E}_k + \tilde{B}_{n-k}\tilde{E}_k(-1) \right) \right\} \frac{t^n}{n!} \right) \\ &= \frac{1}{2^2} \sum_{n=0}^{\infty} \sum_{\substack{v_1, v_2=0 \\ v_1+v_2=2}} \frac{\sum_{k=0}^{v_1} \binom{v_1}{k} \left( \tilde{B}_{v_1-k}(1)\tilde{E}_k + \tilde{B}_{v_1-k}\tilde{E}_k(-1) \right)}{v_1!} \\ &\quad \times \frac{\sum_{k=0}^{v_2} \binom{v_2}{k} \left( \tilde{B}_{v_2-k}(1)\tilde{E}_k + \tilde{B}_{v_2-k}\tilde{E}_k(-1) \right)}{v_2!}. \end{aligned}$$

Therefore, we can derive the below equation when  $\alpha = m$ .

$$\begin{aligned} \tilde{B}_n^{(m)} &= \frac{n!}{2^m} \sum_{\substack{v_1, \dots, v_m=0 \\ v_1 + \dots + v_m = n}} \frac{\left( \sum_{k=0}^{v_1} \binom{v_1}{k} \left( \tilde{B}_{v_1-k}(1)\tilde{E}_k + \tilde{B}_{v_1-k}\tilde{E}_k(-1) \right) \right) \cdots}{v_1 \cdots} \\ &\quad \times \frac{\left( \sum_{k=0}^{v_m} \binom{v_m}{k} \left( \tilde{B}_{v_m-k}(1)\tilde{E}_k + \tilde{B}_{v_m-k}\tilde{E}_k(-1) \right) \right)}{v_m!}. \end{aligned}$$

□

**Theorem 3.6.** Let  $n \in \mathbb{N}$  and  $\alpha \in \mathbb{C}$ . Then we have

$$\tilde{B}_n^{(\alpha)}(x+y) = \sum_{j=0}^n \binom{n}{j} \sum_{k=0}^{n-j} \binom{x}{k} k! \tilde{B}_j^{(\alpha)}(y) S_2(n-j, k),$$

where  $S_2(n-j, k)$  is the stirling numbers of the second kind.

*Proof.* From Definition which is the stirling numbers of the second kind we can find this theorem. Note that

$$x^n = \sum_{k=0}^n \binom{x}{k} k! S_2(n, k),$$

where  $S_2(n, k)$  are the stirling numbers of the second kind.

By combining the above equation and Corollary 2.3.(i), we can easily see that

$$\tilde{B}_n^{(\alpha)}(x+y) = \sum_{j=0}^n \binom{n}{j} \sum_{k=0}^{n-j} \binom{x}{k} k! \tilde{B}_j^{(\alpha)}(y) S_2(n-j, k),$$

Therefore, we complete the proof.

□

**Theorem 3.7.** Let  $n \in \mathbb{N}_0$  and  $\alpha \in \mathbb{C}$ . Then we derive

$$\tilde{B}_n^{(\alpha)} = \sum_{k=0}^n \binom{n}{k} 2^{n-\alpha-k} \sum_{j=0}^{n-k} \frac{(-1)^j}{\binom{n+j-k}{j}} \binom{\alpha+j-1}{j} b(n-k+j, j) \alpha^k,$$

where  $b(n-k+j, j)$  is the associated stirling numbers of the second kind.

*Proof.* By using the associated stirling numbers of the second kind, we find that

$$\begin{aligned} \sum_{n=0}^{\infty} \tilde{B}_n^{(\alpha)} \frac{t^n}{n!} &= \left( \frac{te^t}{e^{2t}-1} \right)^\alpha = \frac{1}{2^\alpha} \left( \frac{1}{1 + \frac{1}{2t}(e^{2t}-2t-1)} \right)^\alpha e^{\alpha t} \\ &= \frac{1}{2^\alpha} \sum_{j=0}^{\infty} \binom{\alpha+j-1}{j} \left( -\frac{1}{2} \right)^j (e^{2t}-2t-1)^j t^{-j} e^{\alpha t} \\ &= \frac{1}{2^\alpha} \sum_{j=0}^{\infty} (-1)^j \binom{\alpha+j-1}{j} j! \sum_{n=j}^{\infty} 2^n b(n+j, j) \frac{t^n}{(n+j)!} e^{\alpha t} \\ &= \frac{1}{2^\alpha} \sum_{n=0}^{\infty} \sum_{k=0}^n \binom{n}{k} 2^{n-k} \sum_{j=0}^{n-k} \frac{(-1)^j}{\binom{n+j-k}{j}} \binom{\alpha+j-1}{j} b(n+j-k, j) \alpha^k \frac{t^n}{n!}. \end{aligned}$$

Therefore we express  $\tilde{B}_n^{(\alpha)}$  as follows.

$$\tilde{B}_n^{(\alpha)} = \sum_{k=0}^n \binom{n}{k} 2^{n-\alpha-k} \sum_{j=0}^{n-k} \frac{(-1)^j}{\binom{n+j-k}{j}} \binom{\alpha+j-1}{j} b(n-k+j, j) \alpha^k.$$

□

**Theorem 3.8.** Let  $n \in \mathbb{N}_0$  and  $\alpha \in \mathbb{C}$ . Then we obtain

$$\tilde{B}_n^{(\alpha)}(x) = \sum_{l=0}^n \sum_{k=0}^{n-l} \binom{n-k}{l} 2^{n-\alpha-l-k} \alpha^k \sum_{j=0}^{n-l-k} \binom{n+j}{k} \frac{(-1)^j}{\binom{n+j}{j}} \binom{\alpha+j-1}{j} b(n+j-l-k, j) x^l.$$

*Proof.* By using the associated stirling numbers of the second kind, we express the polynomials  $\tilde{B}_n^{(\alpha)}(x)$  as follows.

$$\begin{aligned} \sum_{n=0}^{\infty} \tilde{B}_n^{(\alpha)}(x) \frac{t^n}{n!} &= \left( \frac{te^t}{e^{2t}-1} \right)^\alpha e^{tx} = \sum_{n=0}^{\infty} \tilde{B}_n^{(\alpha)} \frac{t^n}{n!} \sum_{n=0}^{\infty} x^n \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \sum_{l=0}^n \sum_{k=0}^{n-l} \binom{n-k}{l} 2^{n-\alpha-l-k} \alpha^k \sum_{j=0}^{n-l-k} \binom{n+j}{k} \frac{(-1)^j}{\binom{n+j}{j}} \binom{\alpha+j-1}{j} \\ &\quad \times b(n+j-l-k, j) x^l \frac{t^n}{n!}. \end{aligned}$$



From the above equation, we have the Theorem 3.8 as follows:

$$\tilde{B}_n^{(\alpha)}(x) = \sum_{l=0}^n \sum_{k=0}^{n-l} \binom{n-k}{l} 2^{n-\alpha-l-k} \alpha^k \sum_{j=0}^{n-l-k} \binom{n+j}{k} \frac{(-1)^j}{\binom{n+j}{j}} \binom{\alpha+j-1}{j} b(n+j-l-k, j) x^l.$$

Thus, we complete the proof of Theorem 3.8. □

**Theorem 3.9.** Let  $n \in \mathbb{N}_0$  and  $\alpha \in \mathbb{C}$ . Then we obtain that

$$\left(\tilde{B} + \tilde{B}^{(\alpha)}(1 + \alpha)\right)^n = \frac{2^n S_n(\alpha)}{2^{\alpha+1}},$$

where  $S_n(\alpha)$  is the stirling polynomials.

*Proof.* Consider that  $\tilde{B}^n = \tilde{B}_n$ . Then we get the following equation from binomial operation and definition of stirling polynomials.

$$\begin{aligned} 2^{\alpha+1} \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n \left(\tilde{B} + \tilde{B}^{(\alpha)}(1 + \alpha)\right)^n \frac{t^n}{n!} &= 2 \sum_{n=0}^{\infty} \tilde{B}_n \frac{\left(\frac{1}{2}t\right)^n}{n!} 2^\alpha \sum_{n=0}^{\infty} \tilde{B}_n^{(\alpha)}(1 + \alpha) \frac{\left(\frac{1}{2}t\right)^n}{n!} \\ &= \left(\frac{te^t}{e^t - 1}\right)^{\alpha+1} = \sum_{n=0}^{\infty} S_n(\alpha) \frac{t^n}{n!}. \end{aligned}$$

Hence we can finish the proof. □

**Theorem 3.10.** Let  $n \in \mathbb{N}_0$  and  $k \in \mathbb{N}$ . Then we derive that

$$\binom{n}{2k} \tilde{B}_{n-2k}^{(-2k)}(x) = \sum_{l=0}^{n-2k} 2^{2k+l} T(k+l, k) x^{n-(2k+l)},$$

where  $T(n, k)$  is the central factorial numbers.

*Proof.* Consider that

$$\sum_{n=k}^{\infty} T(n, k) \frac{t^{2n}}{(2n)!} = \frac{(e^t - 1)^{2k}}{(2k)! e^{tk}}.$$

In order to make relation of the second kind Bernoulli polynomials and the central factorial numbers, we can express as the follows:

$$\frac{1}{4^k (2k)!} \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n \tilde{B}_n^{(-2k)}(x) \frac{t^{2k+n}}{n!} = \sum_{n=k}^{\infty} T(n, k) \frac{t^{2n}}{(2n)!} \sum_{n=0}^{\infty} \left(\frac{1}{2}x\right)^n \frac{t^n}{n!}.$$

From the above equation, we obtain the below equation by using some calculation and comparing the both sides of  $\frac{t^{2k+n}}{(2k+n)!}$ .

$$\binom{n}{2k} \left(\frac{1}{2}\right)^n \tilde{B}_{n-2k}^{(-2k)}(x) = \sum_{l=0}^{n-2k} \frac{T(k+l, k)}{2^{n-2k-l}} x^{n-2k-l}.$$

Therefore the Theorem 3.10 is clear.

□

Let  $l \in \mathbb{N}_0$  and  $n$  be non-negative integers. Then we can define  $\tilde{P}_l(n)$  as follows:

$$\tilde{P}_l(n) = \tilde{B}_l(2n) - \tilde{B}_l = l \sum_{i=0}^{n-1} (1+2i)^{l-1}.$$

where  $\tilde{P}_l(n)$  is the symmetric power sum polynomials.

**Theorem 3.11.** For  $w_1, w_2$  be positive integers and  $\alpha \in \mathbb{C}$ , we get

$$\begin{aligned} & \sum_{l=0}^n \binom{n}{l} \tilde{B}_{n-l}^{(\alpha-1)}(w_1 y) \sum_{s=0}^l \binom{l}{s} w_1^{l-s} w_2^{n-l+s} \tilde{B}_{l-s}^{(\alpha)}(w_2 x) \tilde{P}_s(w_1) \\ &= \sum_{l=0}^n \binom{n}{l} \tilde{B}_{n-l}^{(\alpha-1)}(w_2 y) \sum_{s=0}^l \binom{l}{s} w_2^{l-s} w_1^{n-l+s} \tilde{B}_{l-s}^{(\alpha)}(w_1 x) \tilde{P}_s(w_2). \end{aligned}$$

*Proof.* Where

$$C(t) := \frac{(e^{2w_1 w_2 t} - 1)(w_1 w_2 t^2 e^{t(w_1+w_2)})^\alpha e^{w_1 w_2 t(x+y)}}{(e^{2w_1 t} - 1)^\alpha (e^{2w_2 t} - 1)^\alpha},$$

say that

$$\begin{aligned} C(t) &:= \left( \frac{w_1 t e^{w_1 t}}{e^{2w_1 t} - 1} \right)^\alpha e^{w_1 w_2 t x} \sum_{i=0}^{w_1-1} w_2 t e^{(1+2i)w_2 t} \left( \frac{w_2 t e^{w_2 t}}{e^{2w_2 t} - 1} \right)^{\alpha-1} e^{w_1 w_2 t y} \\ &= \sum_{n=0}^{\infty} \tilde{B}_n^{(\alpha)}(w_2 x) \frac{(w_1 t)^n}{n!} \sum_{s=0}^{\infty} \tilde{P}_s(w_1) \frac{(w_2 t)^s}{s!} \sum_{n=0}^{\infty} \tilde{B}_n^{(\alpha-1)}(w_1 y) \frac{(w_2 t)^n}{n!} \\ &= \sum_{n=0}^{\infty} \left( \sum_{l=0}^n \binom{n}{l} \tilde{B}_{n-l}^{(\alpha-1)}(w_1 y) \sum_{s=0}^l \binom{l}{s} w_1^{l-s} w_2^{n-l+s} \tilde{P}_s(w_1) \tilde{B}_{l-s}^{(\alpha)}(w_2 x) \right) \frac{t^n}{n!}, \end{aligned} \tag{3.2}$$

which is true.  $A(t)$  get transformed in the other following equation.

$$\begin{aligned} C(t) &:= \left( \frac{w_2 t e^{w_2 t}}{e^{2w_2 t} - 1} \right)^\alpha e^{w_1 w_2 t x} \sum_{i=0}^{w_2-1} 2w_1 t e^{(1+2i)w_1 t} \left( \frac{w_1 t e^{w_1 t}}{e^{2w_1 t} - 1} \right)^{\alpha-1} e^{w_1 w_2 t y} \\ &= \sum_{n=0}^{\infty} \left( \sum_{l=0}^n \binom{n}{l} \tilde{B}_{n-l}^{(\alpha-1)}(w_2 y) \sum_{s=0}^l \binom{l}{s} w_2^{l-s} w_1^{n-l+s} \tilde{P}_s(w_2) \tilde{B}_{l-s}^{(\alpha)}(w_1 x) \right) \frac{t^n}{n!}. \end{aligned} \tag{3.3}$$

Comparing the coefficients of  $\frac{t^n}{n!}$  in (3.2) and (3.3), we get the equation as:

$$\begin{aligned} & \sum_{l=0}^n \binom{n}{l} \tilde{B}_{n-l}^{(\alpha-1)}(w_1 y) \sum_{s=0}^l \binom{l}{s} w_1^{l-s} w_2^{n-l+s} \tilde{B}_{l-s}^{(\alpha)}(w_2 x) \tilde{P}_s(w_1) \\ &= \sum_{l=0}^n \binom{n}{l} \tilde{B}_{n-l}^{(\alpha-1)}(w_2 y) \sum_{s=0}^l \binom{l}{s} w_2^{l-s} w_1^{n-l+s} \tilde{B}_{l-s}^{(\alpha)}(w_1 x) \tilde{P}_s(w_2). \end{aligned}$$

Therefore we complete the proof.

□

By using Theorem 3.11, we also get symmetric property of the second kind Bernoulli polynomials of order  $\alpha$

**Corollary 3.12.** By substituting  $w_1 = 1$  from Theorem 3.11, we have

$$\begin{aligned} & \sum_{l=0}^n \binom{n}{l} \tilde{B}_{n-l}^{(\alpha-1)}(y) \sum_{s=0}^l s \binom{l}{s} w_2^{n-l+s} \tilde{B}_{l-s}^{(\alpha)}(w_2 x) \\ &= \sum_{l=0}^n \binom{n}{l} \tilde{B}_{n-l}^{(\alpha-1)}(w_2 y) \sum_{s=0}^l \binom{l}{s} w_2^{l-s} \tilde{B}_{l-s}^{(\alpha)}(x) \tilde{P}_s(w_2). \end{aligned}$$

#### Acknowledgements

This work was supported by NRF(National Research Foundation of Korea) Grant funded by the Korean Government(NRF-2013-Fostering Core Leaders of the Future Basic Science Program).

#### References

- [1] M. Alkan and Y. Simsek, Generating function for q-Eulerian polynomials and their decomposition and applications, *Fixed Point Theory and Applications*, (72)(2013).
- [2] Abdelmejid Bayad, Yilmaz Simsek, H. M. Srivastava, Some array type polynomials associated with special numbers and polynomials, *Applied Mathematics and Computation*, 244(2014), 149-157.
- [3] L. Comtet, *Advanced Combinatorics*, Reidel, Boston, Mass, 1974.
- [4] F. T. Howard, Congruences for stirling numbers and associated Stirling numbers, *Acta Arithmetica*, 55(1) (1990), 29-41.
- [5] Donald E. Knuth, Two Notes on Notation, arXiv:math/9205211v1 [math.HO] 1 May 1992.
- [6] T. Kim,  $q$ -Volkenborn integration, *Russ. J. Math. Phys.*, 9 (2002), 288-299.
- [7] Y. H. Kim, H. Y. Jung, C. S. Ryoo, On the generalized Euler polynomials of the second kind, *J. Appl. Math. and Informatics*, 31(5-6) (2013), 623-630.
- [8] J. Y. Kang, C. S. Ryoo, A research on the new polynomials involved with the central factorial numbers, Stirling numbers and others polynomials, *Journal of Inequalities and Applications*, 2014:16 (2014).
- [9] M-S. Kim, J-W. Son, Analytic properties of the  $q$ -volkenborn integral on the ring of  $p$ -adic integres, *Bull. Korean Math. Soc.*, 44 (2007), 1-12.

- [10] Hui Luo, Guodong Liu, An identity involving Norlund numbers and Stirling numbers of the first kind, *Scientia Magna*, 4(2) (2008), 45-48.
- [11] Guo-Dong Liu, H. M. Srivastava, Explicit formulas for the Norlund polynomials  $B_n^{(x)}$  and  $b_n^{(x)}$ , *Computers and Mathematics with Applications*, 51 (2006), 1377-1384.
- [12] Guodong Liu, Congruences for higher-order Euler numbers, *Proc. Japan Acad.*, 82, Ser. A (2006).
- [13] Guo Dong LIU, Wen Peng Zhang, Applications of an explicit formula for the generalized Euler numbers, *Acta Mathematica Sinica, English Series*, 24(2) (2008), 343-352.
- [14] Qiu-Ming Luo, H. M. Srivastava, Some relationships between the Apostol-Bernoulli and Apostol-Euler polynomials, *Computers and Mathematics with Applications*, 51 (2006), 631-642.
- [15] H. Ozden and Y. Simsek, Modification and unification of the Apostol-type numbers and polynomials and their applications, *Applied Mathematics and Computation*, 235 (2014), 338351.
- [16] C. S. Ryoo, Calculating zeros of  $q$ -extension of the second kind Bernoulli polynomials, *Journal of Computational Analysis and Applications*, 15(2) (2013), p248.
- [17] C. S. Ryoo, A note on the second kind Bernoulli polynomials, *Journal of Computational Analysis and Applications*, 12 (2010), 828-833.
- [18] C. S. Ryoo, A note on the second kind Genocchi polynomials, *J. Comput. Anal. Appl.*, 13 (2011), 986992.
- [19] S. Roman, *The Stirling Polynomials*, 4.8 in *The Umbral Calculus*. New York: Academic Press, 1984, 128-129.
- [20] H. M. Srivastava, Á. Pintér, Remarks on some relationships between the Bernoulli and Euler polynomial, *Appl. Math. Lett.*, 17(4) (2004), 375-380.
- [21] Y. Simesk, V. Kurt and D. Kim, New approach to the complete sum of products of the twisted  $(h, q)$ -Bernoulli numbers and polynomials, *J. Nonlinear Math. Phys.*, 14 (2007), 4456.
- [22] Y. Simesk, Special numbers on analytic functions, *Applied Mathematics*, 5 (2014), 1091-1098.
- [23] Y. Simesk, Generating functions for generalized Stirling type numbers, Array type polynomials, Eulerian type polynomials and their applications, *Fixed Point Theory and Applications*, (87) (2013).

# An oscillation of the solution for a nonlinear second-order stochastic differential equation

Iryna Komashynska<sup>a,\*</sup>, Mohammed AL-Smadi<sup>b</sup>, Ali Atewi<sup>c</sup>, Ayed Al e'damat<sup>c</sup>

<sup>a</sup>*Department of Mathematics, Faculty of Science, The University of Jordan, Amman 11942, Jordan*

<sup>b</sup>*Applied Science Department, Ajloun College, Al-Balqa Applied University, Ajloun 26816, Jordan*

<sup>c</sup>*Department of Mathematics, Faculty of Science, Al-Hussein Bin Talal University, P.O. Box 20, Ma'an-Jordan*

\*Corresponding author: e-mail: iryna\_kom@hotmail.com

**Abstract.** In this paper, we study the oscillatory properties for asymptotic behaviors of solutions of a class of nonlinear second-order stochastic Itô equations. Meanwhile, we investigate existence of zeros of its solutions with probability 1. Sufficient conditions for the oscillation and nonoscillation of solutions are obtained on the half-line  $[t_0, \infty)$  for every  $t_0 > 0$ .

**Keywords:** Oscillation, Stochastic differential equations, Zeros of solutions, Wiener process, Itô integral

**AMS Subject Classification:** 60H10; 60H25; 34K11

## 1 Introduction

During the past few decades, stochastic differential equations (SDEs) are becoming increasingly important as models of stochastic phenomena that play a prominent role in a diverse range of application areas, including mathematical modeling in engineering and physics, geophysical sciences, stochastic control, mechanics, environmental processes, mathematical biology, molecular dynamics for chemistry, epidemiology, economic modeling, industrial mathematics and mathematical finance [1-10]. Indeed, these models can be stochastic for different reasons. Therefore, numerous studies have been performed to understanding their dynamical behaviors, particularly in relation to problems of the specification of the stochastic processes governing the behaviors of an underlying quantity, as well as fundamental microscopic laws generate stochastic behaviors in the case of coarse-graining and modeling error and so on [11-16]. However, a complete understanding of SDEs theory requires familiarity with advanced probability and stochastic processes, whereas solutions of such models are themselves stochastic processes.

Further, in particular, second-order differential equations with random coefficients have found wide variety applications in branches of science. Typically, they are mathematical models of objects under the influence of random forces such that the presence of infinite set of zeros of solutions for these equations indicates that the evolution of investigated objects is oscillatory. Recently, research work about oscillation phenomenon occupies an important place in differential stochastic theory due to the sensitivity of stochastic forces and behaviors. Moreover, the stochastic theory for these equations, as well as the theory of oscillatory solutions of deterministic equations have been studied extensively and are well-developed. Oscillation and nonoscillation conditions for both linear and nonlinear differential equations, difference equations and delay equations have been investigated in [17-22]. The oscillating properties for solutions of difference equations can be found in the excellent monograph of Agarwal et al. [23]. Besides, the authors in the monograph [24] were devoted to the problem of relationship between oscillation behavior of solutions for differential equations and the corresponding difference equations. On the contrary, the theory of the oscillation of stochastic system is not well-developed.

Incidentally, Mao in [25] considered the stochastic equation of the following form

$$\ddot{x} + kx = h\dot{W}(t),$$

where  $\dot{W}(t)$  is a Wiener process which is nowhere differentiable. It was proved that the solution with initial values  $x(0) = 1, \dot{x}(0) = 0$  has infinitely many zeros, all simple, on each half-line  $[t_0, \infty)$  for every  $t_0 \geq 0$ . The first two moments of the first zero were estimated.

In contrast, the more general equation of the form

$$\ddot{x} + k(t, x, \dot{x}) = h\dot{W}(t),$$

was studied in [26]. The author there demonstrated that this equation has infinitely many zeros with probability 1. Consequently, the explicit upper and lower estimates for the expected values of these zeros were obtained. However, the Itô stochastic equations of the form

$$\ddot{x} + (p(t) + q(t)\dot{W}(t))x = 0$$

was considered by method of asymptotic equivalence in [27,28], whereas the oscillation of solutions was analyzed. In the monographs [29,30], the oscillatory properties of solutions for both linear and nonlinear stochastic delay differential equations with multiplicative noise are given. It was shown that such noise induces an oscillation in solutions. Besides, the oscillation of solutions of first order nonlinear stochastic difference equations is investigated in [31].

The purpose of this paper is to study an asymptotic behavior, as  $t \rightarrow \infty$ , of solutions of a second order stochastic Itô equation. Meanwhile, we investigate existence of zeros of its solutions with probability 1. In the sequel, unless otherwise specified, we say that a solution is oscillatory if it has infinitely many zeros with probability 1 on the half-line  $[0, \infty)$ . A solution which is not oscillatory is called nonoscillatory.

This paper is organized in five sections including the introduction. In the next section, we present some necessary definitions and preliminary results that will be used in this work. In the same time, statement of a second order SDEs is introduced. In Section 3, the discussion of a solution for linear case of second-order SDEs is presented, as well as the conditions of nonoscillatory behavior of its solutions for nonlinear case of SDEs are constructed. Finally, the conclusions are drawn in Section 4.

## 2 Statement of the problem and auxiliary results

The material in this section is basic in some sense. For the reader's convenience, we present some necessary definitions and auxiliary results related to the SDEs theory that will be used in the remainder of this paper.

Let us consider a nonlinear second-order stochastic equation of the following form

$$\ddot{x} + p(t, x, \dot{x}) + q(t, x, \dot{x})\dot{W}(t) = 0, t \geq 0. \tag{1}$$

While the corresponding system of stochastic Itô equations will be written as

$$\begin{aligned} dx_1 &= x_2 dt, \\ dx_2 &= -p(t, x_1, x_2)dt - q(t, x_1, x_2)dW(t), \end{aligned} \tag{2}$$

where  $x \in R^1, t \geq 0, W(t)$  is a standard Wiener process defined on the probability space  $(\Omega, F, P), \{F_t, t \geq 0\}$  is the family of  $\sigma$ -algebras adapted to  $W(t)$ , and the functions  $p(t, x_1, x_2)$  and  $q(t, x_1, x_2)$  are continuous with respect to  $x_1, x_2 \in R^1$  for  $t \geq 0$ , as well as satisfy the Lipschitz condition with respect to  $x_1, x_2$  together with linear growth condition. Without loss of generality, we assume that  $p(t, 0, 0) = q(t, 0, 0) = 0$ .

It should be noted that the presence of stochastic in equation (1) causes new difficulties in studying the oscillation of solutions. In this regard, we mention here the following remark: Firstly, solutions of equation (1) are random processes, so their zeros are random variables with certain properties. As a consequence, we need to introduce a new definition of zero which is different of the deterministic case ( $q = 0$ ). Secondly, from the Strook-Varadham support theorem, it follows that solutions of equation (1) can be nonoscillatory on finite intervals. Therefore, the oscillatory solutions should be considered only on infinite intervals. Thirdly, since solutions of equation (1) have only first derivative, so we can not use a second derivative to apply the concavity property of the solution between two successive zeros. It is well known that this method is used in the deterministic case.

Nevertheless, system (2) is a particular case of general second-order system, it would seem that this simplifies its investigation, as well it is the system with a degenerate diffusion that completes its investigation by probability methods. Subsequently, under the above assumptions of equation (1) and the corresponding system (2), we assume that the solution  $\bar{x}(t) = (x_1(t), x_2(t))$  of system (2) subject to the initial condition  $\bar{x}(t_0) = \bar{x}_0$  satisfy all necessary requirements of the existence of a unique solution for  $t \geq t_0$ , whereas  $\bar{x}_0$  is an  $F_{t_0}$ -measurable random variable. In addition, the process  $\bar{x}(t)$  will never reach the origin  $(0,0)$ , for more details see Lemma 2.3 in [32]. In our notation, let  $x_1(t) = x(t)$ . Throughout this paper, a solution  $x(t)$  of equation (1) is called a nontrivial solution if it satisfy the following condition

$$P \{x(t) = 0, t > t_0\} = 0$$

On the other hand, for any nontrivial solution  $x(t)$  of equation (1), where  $t \geq t_0 \geq 0$ , the random variable  $\tau_1$  can be defined as follows

$$\tau_1 = \begin{cases} \inf \{t > t_0 \mid x_1(t) = 0\}, & \text{if } \{t > t_0 \mid x_1(t) = 0\} \neq \emptyset, \\ \infty, & \text{otherwise.} \end{cases} \tag{3}$$

Now, we will introduce the definition of zeros of a solution  $x(t)$  on the half-line  $t > 0$ .

**Definition 2.1** The random variable  $\tau_1$  is called the first zero of a solution  $x(t)$  on the interval  $t \geq t_0$ , if  $\tau_1 < \infty$  with probability 1.

In consequence, one can define another random variable  $\tau_2$  as follows

$$\tau_2 = \begin{cases} \inf \{t > \tau_1 \mid x_1(t) = 0\}, & \text{if } \{t > \tau_1 \mid x_1(t) = 0\} \neq \emptyset, \\ \infty, & \text{otherwise.} \end{cases} \tag{4}$$

Here, the random variable  $\tau_2$  is called the second zero of a solution  $x(t)$  on the interval  $t \geq t_0$ , if  $\tau_2 < \infty$  with probability 1.

Correspondingly, one can define by induction a sequence of zeros  $\{\tau_n\}$  of a solution  $x(t)$  on the interval  $t \geq t_0$ . Particularly, if  $t_0 = 0$ . Then, we deal with zeros on the half-line  $t > 0$ .

**Definition 2.2** A nontrivial solution  $x(t)$  of equation (1) is called oscillatory on the half-line  $t > 0$ , if it has infinitely many zeros there. Otherwise, it is called nonoscillatory.

### 3 Main results and behavior solutions of the SDEs

In this section, some definitions and results are briefly reviewed to establish and generalize the results to the main equation in this work. Meanwhile, we study the behavior of the zeros of solutions for a class of second-order SDEs subject to some initial conditions, as well as we detect the conditions of nonoscillatory behavior of its solutions.

#### 3.1 Linear stochastic Itô equation

Consider the following equation

$$\ddot{x} + x = f(t)\dot{W}(t), \tag{5}$$

subject to the initial conditions

$$x(0) = 1, \dot{x}(0) = 0, \tag{6}$$

where  $f(t)$  is a nonrandom function defined on  $t \geq 0$  such that a stochastic Itô integral

$$\int_0^t f(s)dW(s)$$

is defined for any  $t > 0$ .

Note that equation (5) is special case of equation (1), so it satisfies all arguments mentioned in the previous part of our work. Further, we give the following theorem regarding to study the behavior of the zeros of solutions for equation (5) with initial conditions (6).

**Theorem 3.1** Assume that  $f(t)$  satisfies the following conditions:

1.  $f(t)$  is differentiable function for  $t \geq 0$  such that  $f(0) \geq 0$ ,
2.  $(\sin(t - s)f(s))' \leq 0$  for  $0 \leq s \leq t \leq \frac{\pi}{2}$ .

Then, the solution of equation (5) subject to initial conditions (6) oscillates on the half-line  $t \geq 0$ . Besides, a mathematical expectation  $\tau_1$  of the first zero satisfies the estimation

$$E\tau_1 \geq 2t^* \Phi\left(\frac{1}{\sqrt{t^*}}\right), \tag{7}$$

where  $t^*$  is the solution of the equation

$$f(0) = \cot(t) \tag{8}$$

on  $[0, \frac{\pi}{2}]$  and  $\Phi(z) = \frac{1}{2\pi} \int_0^z e^{-\frac{u^2}{2}} du$ .

**Proof.** From Itô formula, the representation of the solution of (5) with initial conditions (6) is given by

$$x(t) = \cos(t) + \int_0^t f(s) \sin(t - s)dW(s). \tag{9}$$

Which implies that

$$x(t) = \cos(t) + \sin(t) \int_0^t f(s) \cos(s)dW(s) - \cos(t) \int_0^t f(s) \sin(s)dW(s).$$

Accordingly, the process  $x(t)$  can be written as

$$x(t) = \cos(t) + \bar{W}_1(p(t)) \sin(t) + \bar{W}_2(q(t)) \cos(t),$$

where

$$p(t) = \int_0^t f^2(s) \cos^2(s)ds, \quad q(t) = \int_0^t f^2(s) \sin^2(s)ds,$$

and  $\bar{W}_1, \bar{W}_2$  are Wiener processes.

In contrast, if we consider  $x(t)$  at the times  $t_m = (2m + \frac{1}{2})\pi$  for  $m = 1, 2, 3, \dots$ , and define a sequence

$$\{Y_m\} \text{ by } Y_m = x((2m + \frac{1}{2})\pi) - x((2(m - 1) + \frac{1}{2})\pi), \text{ whereas } x((2m + \frac{1}{2})\pi) = \bar{W}_1 \left( \int_0^{(2m + \frac{1}{2})\pi} \cos^2(s)f^2(s)ds \right).$$



Then,  $Y_0 = \bar{W}(\frac{\pi}{4})$ ,  $Y_1 = \bar{W}_1(\frac{5\pi}{4}) - \bar{W}_1(\frac{\pi}{4})$ , ..., is a sequence of random variables with mean zero and variance

$$\int_{(2(m-1)+\frac{1}{2})\pi}^{(2m+\frac{1}{2})\pi} \cos^2(s)f^2(s)ds.$$

Here, it is worth to mention that

$$x((2m + \frac{1}{2})\pi) = Y_0 + Y_1 + \dots + Y_m. \tag{10}$$

By the familiar theorems on the limits of sums of independent random variables (e.g. the law of the iterated logarithm), it follows that the sequence  $\{x((2m + \frac{1}{2})\pi)\}$  has infinitely many switches of sign. Since  $x(t)$  is continuous on  $[0, \infty)$ , so it has infinitely many zeros on  $[0, \infty)$ . Therefore, it oscillates on  $[0, \infty)$ .

Now, let us prove the estimation (7) for the first zero of the oscillation. By applying the integration-by-parts formula to (9), we obtain

$$x(t) = \cos(t) - \int_0^t (\sin(t-s)f(s))'dW(s) \geq \cos(t) + \int_0^t (\sin(t-s)f(s))'ds, \tag{11}$$

for  $\omega \in \Omega$ , where  $W(t) \geq -1$  and  $0 \leq s \leq t \leq \frac{\pi}{2}$ .

From properties of a Wiener process, it follows that

$$P \left\{ \omega \mid \max_{t \in [0, T]} W(t) > -1 \right\} = 2\Phi\left(\frac{1}{2\sqrt{T}}\right), \tag{12}$$

where  $\Phi(z) = \frac{1}{2\sqrt{2\pi}} \int_0^z e^{-\frac{u^2}{2}} du$ .

As a result, from equation (11), we obtain the estimate

$$x(t) \geq \cos(t) - f(0) \sin(t) > 0, \tag{13}$$

for  $t \in [0, t^*)$ , where  $t^*$  is solution of equation (8).

Hence, from equations (12) and (13), we have

$$P \left\{ \tau \geq \cot^{-1}(f(0)) \right\} \geq 2\Phi \left( \frac{1}{\sqrt{\cot^{-1}(f(0))}} \right), \tag{14}$$

for the first zero  $\tau_1$ . By using Chebyshev's inequality, it yields that  $E\tau_1 \geq 2t^* \Phi(\frac{1}{\sqrt{t^*}})$ . The proof is complete. ■

### 3.2 Nonlinear stochastic Itô equation

Let  $P(a, n) = \{x \in R^m \mid (x - a, n) \geq 0\}$ , where  $a, n \in R^m$ , and  $(\cdot, \cdot)$  is the usual scalar product. Thus, a polyhedron is any set of the form

$$\bigcap_{\alpha \in I} P(a_\alpha, n_\alpha), \tag{15}$$

where  $I = \{1, \dots, N\}$  is a finite subset of  $N$ .

Now, consider a system of SDEs

$$dx = f(t, x)dt + g(t, x)dW(t) \tag{16}$$

where  $x \in R^m$ ,  $f : [0, \infty) \times R^m \rightarrow R^m$ ,  $g = [g_{ij}] : [0, \infty) \times R^m \rightarrow R^m \times R^r$  are mappings, and  $W(t)$  is an  $r$ -dimensional Wiener process.

**Definition 3.1** A set  $K \in R^m$  is said to be stochastically invariant for system (16), if for any  $x(0) \in K$  and every solution  $x(t)$  of equation (1), then  $P\{x(t) \in K, t > 0\} = 1$ .

The next theorem states conditions of an invariance of the set (15) for system (16).

**Theorem 3.2** [33] Let  $K = \bigcap_{\alpha \in I} P(a_\alpha, n_\alpha)$  be a polyhedron in  $R^m$ . Suppose that the coefficients  $f(t, x)$  and  $g(t, x)$  of system (16) are defined for  $t \geq 0$ ,  $x \in R^m$ , and satisfy the following conditions:

1. for each  $T > 0$ , there exists a constant  $K_T > 0$  such that for all  $x \in K$  and  $t \in [0, T)$ ,

$$\|f(t, x)\|^2 + \|g(t, x)\|^2 \leq K_T(1 + |x|^2);$$

2. for all  $T > 0$ ,  $x \in K$ ,  $y \in K$  and  $t \in [0, T)$ ,

$$\|f(t, x) - f(t, y)\| + \|g(t, x) - g(t, y)\| \leq K_T |x - y|;$$

3. for each  $x \in K$ , the functions  $f(\cdot, x)$  and  $g(\cdot, x)$ , defined for  $t \geq 0$ , are continuous.

Then, the set  $K$  is invariant for the system (16) if and only if the following condition holds:

- (a) for all  $\alpha \in I$  and  $x \in K$  such that  $(x - a_\alpha, n_\alpha) = 0$ , we have

$$(f(t, x), n_\alpha) \geq 0 \text{ and } (g_j(t, x), n_\alpha) = 0,$$

where  $t \geq 0$ ,  $j = \overline{1, r}$ , and  $g_j$  is the  $j$ -th column of the matrix  $g = [g_{ij}]$ .

Now, we use the above theorem to find the conditions of nonoscillatory behavior of the solutions of equation (1). As well, we state the following theorem:

**Theorem 3.3** Suppose that the functions  $p$  and  $q$  in equation (1) satisfy the conditions (1)-(3) of Theorem 3.2. Moreover, if

- (1)  $p(t, x_1, 0) \geq 0$ ,  $x_1 < 0$ ,  $t \geq 0$ ;
  - (2)  $p(t, x_1, 0) \leq 0$ ,  $x_1 > 0$ ,  $t \geq 0$ ;
  - (3)  $q(t, x_1, 0) = 0$ ,  $t \geq 0, x_1 \in R^1$ .
- (17)

Then, all solutions of equation (1) with nonrandom initial values such that  $x(0) > 0$ ,  $\dot{x}(0) \geq 0$  or  $x(0) < 0$ ,  $\dot{x}(0) \leq 0$  are not oscillate on the half-line  $[0, \infty)$ .

**Proof.** We consider any solution of equation (1) with initial values  $x(0) > 0$ ,  $\dot{x}(0) \geq 0$ . It corresponds to the solution  $(x_1, x_2)$  of system (2) with initial values  $x_1(0) > 0$ ,  $x_2(0) \geq 0$ . Obviously, there exists  $\epsilon > 0$  such that  $0 < \epsilon \leq x_1(0)$ .

Let  $M$  be a set such that  $M = \{(x_1, x_2) \mid x_1 \geq \epsilon, x_2 \geq 0\}$ . It is a polyhedron, if we set  $a_1 = \epsilon$ ,  $n_1 = l_1 = (1, 0)^T$ ,  $a_2 = 0$ ,  $n_2 = l_2 = (0, 1)^T$ . Then,  $M = \bigcap_{\alpha \in I} P(a_\alpha, n_\alpha)$ , where  $I = \{1, 2\}$ . Consequently, the boundaries of this polyhedron are lines

$$\gamma_1 = \{(x_1, x_2) \mid x_1 = \epsilon, x_2 \geq 0\},$$

$$\gamma_2 = \{(x_1, x_2) \mid x_1 \geq \epsilon, x_2 = 0\}.$$

Therefore, by using Theorem 3.2, the functions  $f$  and  $g$  have the form

$$f(t, x_1, x_2) = (x_2, -p(t, x_1, x_2))^T,$$

$$g(t, x_1, x_2) = (0, -q(t, x_1, x_2))^T,$$

Next, we verify the conditions of Theorem 3.2. On the boundary  $\gamma_1$ , we have

$$(f, n_1) = (f, l_1) = x_2 \geq 0, \text{ and } (g, n_1) = (g, l_1) = 0.$$

From condition (3) of equation (17), we have

$$(f, n_2) = (f, l_2) = -p(t, x_1, 0) \geq 0,$$

and

$$(g, n_2) = (g, l_2) = -q(t, x_1, 0) = 0.$$

on the boundary  $\gamma_2$ .

Again from Theorem 3.2, it follows that the set  $M$  is the invariant set for the solutions of system (2). Thus, the curve  $(x_1(t), x_2(t))$  does not intersect with probability 1 the line  $x_1 = 0$ . This means that the solutions of equation (1) with initial values  $x(0) > 0$ ,  $\dot{x}(0) \geq 0$  do not oscillate.

It remains to consider the case with initial values  $x(0) < 0$ ,  $\dot{x}(0) \leq 0$ . We only introduce the polyhedron  $M_1 = \{(x_1, x_2) \mid x_1 \leq -\epsilon, x_2 \leq 0\}$  instead the set  $M$ . Hence, the proof is complete. ■

#### 4 Concluding remarks

The use of SDEs is a natural way to model real-world phenomena under stochastic processes. In this paper, we study the qualitative behavior of nonlinear second order stochastic differential equations. Interest focuses on solutions of such equations which are oscillatory. A nontrivial solution is called oscillatory if it has infinitely many zeros with probability 1 on half-line. Otherwise, it is called nonoscillatory. The sufficient conditions for the oscillation and nonoscillation of solutions are obtained.

#### References

- [1] L.J.S. Allen, An Introduction to Stochastic Processes With Applications to Biology, Pearson Education Inc., Upper Saddle River, New Jersey, 2003.
- [2] P. Greenwood, L.F. Gordillo, R. Kuske, Autonomous stochastic resonance produces epidemic oscillations of fluctuating size, Proceedings of Prague Stochastics, 2006.
- [3] J. Golec, S. Sathananthan, Stability analysis of a stochastic logistic model, Mathematical and Computer Modelling, 38 (2003), 585–593.
- [4] E. Dogan, E.J. Allen, Derivation of Stochastic Partial Differential Equations for Reaction-Diffusion Processes, Stochastic Analysis and Applications, 29(3) (2011), 424–443.
- [5] W.D. Sharp, E.J. Allen, Stochastic neutron transport equations for rod and plane geometries, Annals of Nuclear Energy, 27 (2000), 99–116.
- [6] K.E. Emmert, L.J.S. Allen, Population extinction in deterministic and stochastic discrete-time epidemic models with periodic coefficients with applications to amphibians, Natural Resource Modeling, 19 (2006), 117–164.
- [7] D.T. Gillespie, Approximate accelerated stochastic simulation of chemically reacting systems, Journal of Chemical Physics, 115 (2001), 1716–1733.

- [8] G.N. Milstein, M.V. Tretyakov, *Stochastic Numerics for Mathematical Physics*, Springer-Verlag, Berlin, 2004.
- [9] J.G. Hayes, E.J. Allen, Stochastic point-kinetics equations in nuclear reactor dynamics, *Annals of Nuclear Energy*, 32 (2005), 572–587.
- [10] E.J. Allen, L.J.S. Allen, A. Arciniega, P. Greenwood, Construction of equivalent stochastic differential equation models, *Stochastic Analysis and Applications*, 26 (2008), 274–297.
- [11] B. Øksendal, *Stochastic Differential Equations*, 6th edition, Springer-Verlag, Berlin, 2002.
- [12] A.M. Atewi, I.V. Komashynska, The existence and unbounded extension of the solutions of the systems with stochastic impulse action, *Journal of Mathematics and Statistics*, 2(4) (2006), 460-463.
- [13] A.M. Atewi, Dissipativity of the systems of differential equations with stochastic impulse action, *International journal of applied mathematics*, 21 (2008), 73-81.
- [14] A.M. Samoilenko, O.M. Stanzhytskyi, A.M. Atewi, On invariant tori for a stochastic Itô system, *Journal of dynamics and differential equations*, 17(4) (2005), 737-758.
- [15] D.J. Higham, Mean-square and asymptotic stability of the stochastic theta method, *SIAM J. Numer. Anal.*, 38 (2000), 753–769.
- [16] I.V. Komashynska, Existence and uniqueness of solutions for a class of nonlinear stochastic differential equations, *Abstract and Applied Analysis*, 2013 (2013), <http://dx.doi.org/10.1155/2013/256809>, 7p (Article ID 256809).
- [17] J.A. Appleby, C. Kelly, Oscillation and non-oscillation in solutions of nonlinear stochastic delay differential equations, *Elect. Comm. in Probab.*, 9 (2004), 106-118.
- [18] I.T. Kiguradze, *Asymptotic properties of solutions of nonautonomous ordinary differential equations*, Nauka, Moscow, 1990.
- [19] A. Lomtadze, N. Partsvania, Oscillation and non-oscillation criteria for two-dimensional systems of first order linear ordinary differential equations, *Georgian Math. Journ.* 6(3) (1999), 285-298.
- [20] V. A. Căuş, Minimal quadratic oscillation for cubic splines, *Journal of Computational Analysis and Applications*, 9 (2007), 85-92.
- [21] Ercan Tunç, Ahmet Eroğlu, Oscillation Results for Second Order Half-Linear Nonhomogeneous Differential Equations with Damping, *Journal of Computational Analysis and Applications*, 15(2) (2013), 255-263.
- [22] A. M. Bica, V. A. Căuş, I. Fechet, S. Mureşan, Application of the Cauchy-Buniakovski-Schwarz's Inequality to an Optimal Property for Cubic Splines, *Journal of Computational Analysis and Applications*, 9 (2007), 43-53.
- [23] R.P. Agarwal, M. Bohner, S.R. Grace, D. O'Regan, *Discrete Oscillation Theory*. Hindawi Publishing Corporation, New York, USA, 2005.
- [24] O.V. Karpenko, V.I. Kravets', O.M. Stanzhytskyi, Oscillation of solutions of the second-order functional difference equations, *Ukr. Math. Journal*, 65(2) (2013), 249–259.
- [25] X. Mao, *Stochastic Differential Equations and Their Applications*, Horwood Publishing, Chichester, UK, 1997.
- [26] L. Markus, A. Weerasighe, Stochastic oscillators, *Journal of differential equations*, 71(3) (1998), 288-314.
- [27] A.M. Samoilenko, O.M. Stanzhytskyi, *Qualitative and asymptotic analysis of differential equations with random perturbations*, Singapore, New Jersey, London, Hong Kong: Word Scientific, 2011.
- [28] A.N. Stanzhytskii, A.P. Krenevich, I.G. Novak, Asymptotic equivalence of linear stochastic Itô systems and oscillation of solutions of linear second-order equations, *Differential Equations*, 47(6) (2011), 799-813.

- [29] J.A. Appleby, E. Buckwar, Noise induced oscillation in solutions of stochastic delay equations, *Dynamic Systems and Applications*, 14(2) (2005), 175-197.
- [30] J. Shao, F. Meng. Oscillations theorems for second -order forced neutral nonlinear differential equations with delayed argument, *Int. J. Diff. Eq.* 2010 (2010), <http://dx.doi.org/10.1155/2010/181784>, 15p (Article ID 181784).
- [31] J.A. Appleby, A. Rodkina, On the oscillation of solutions of stochastic difference equations with state-independent perturbations, *Int. J. Differ Equ.* 2(2) (2007), 139-164.
- [32] R.Z. Khasminskii, *Stochastic stability of differential equations*, Sijthoff and Noordhoff, 1980.
- [33] A. Milian, Stochastic viability and a comparison theorem, *Colloquium mathematicum*, Vol. LXVIII, (1995), 297-316.

# Fixed point theorems and $T$ -stability of Picard iteration for generalized Lipschitz mappings in cone metric spaces over Banach algebras

Huaping Huang<sup>1\*</sup>, Shaoyuan Xu<sup>2</sup>, Hao Liu<sup>1</sup>, Stojan Radenović<sup>3</sup>

1. School of Mathematics and Statistics, Hubei Normal University, Huangshi, 435002, China
2. Department of Mathematics and Statistics, Hanshan Normal University, Chaozhou, 521041, China
3. Faculty of Mathematics and Information Technology, Dong Thap University, Dong Thap, Việt Nam

**Abstract:** In this paper, we obtain the existence of non-normal solid cone and some fixed point theorems for generalized Lipschitz contractive mappings in cone metric spaces over Banach algebras. Our results greatly generalize the main work by Xu and Radenović (Fixed Point Theory and Applications, 2014, 2014: 102). Moreover, we verify the  $P$  property and  $T$ -stability of Picard's iteration. Further, we give an example to illustrate that our works are never a copy of metric results in the literature.

**MSC:** 47H10; 54H25

**Keywords:** Generalized Lipschitz constant,  $P$  property,  $T$  stability, Cone metric space over Banach algebra, Solid cone

## 1 Introduction

Since Huang and Zhang [1] introduced the concept of cone metric space, many scholars have focused on fixed point theorems in such spaces. There are lots of works on fixed point results in the setting of cone metric spaces (see [2-6]). It is said that [1] is well-known as a result of the fact that cone metric spaces generalize metric spaces and expands the famous Banach contraction principle. But recently, it had not yet been a hot topic since some authors appealed to the equivalence of some metric and cone metric fixed point results

---

\*Corresponding author: Huaping Huang. E-mail: mathhph@163.com

(see [7-12]). Owing to these reasons, people set out to lose interest in studying fixed point theorems in cone metric spaces. However, the present situation has gone better since, very recently, Liu and Xu [13] introduced the concept of cone metric space over Banach algebra and obtained some fixed point theorems in normal cone metric spaces over Banach algebras. Moreover, they gave an example to illustrate that the non-equivalence of versions of fixed point theorems between cone metric spaces over Banach algebras and (general) metric spaces (in usual sense), which shows that it is essentially necessary to investigate fixed points in cone metric spaces over Banach algebras. Lately, Xu and Radenović [15] delete the normality of cones and greatly generalize the main results of [13]. Throughout this paper, we obtain the existence of non-normal solid cone for generalized Lipschitz mappings in cone metric spaces over Banach algebras. Moreover, we present some fixed point theorems for such mappings in such setting by omitting the assumptions of normalities of cones. Our theorems include the main results of [13] and [15]. Furthermore, we consider the mapping's  $P$  property and  $T$ -stability of Picard's iteration. Our results greatly unite and extend the main work of [13-15] and [17-21]. In addition, we give an example to illustrate our results in cone metric spaces over Banach algebras are never equivalent to the counterpart of metric spaces.

Let  $\mathcal{A}$  be a Banach algebra with a unit  $e$ , and  $\theta$  the zero element of  $\mathcal{A}$ . A nonempty closed convex subset  $K$  of  $\mathcal{A}$  is called a cone if  $\{\theta, e\} \subset K$ ,  $K^2 = KK \subset K$ ,  $K \cap (-K) = \{\theta\}$  and  $\lambda K + \mu K \subset K$  for all  $\lambda, \mu \geq 0$ . On this basis, we define a partial ordering  $\preceq$  with respect to  $K$  by  $x \preceq y$  if and only if  $y - x \in K$ . We shall write  $x \prec y$  to indicate that  $x \preceq y$  but  $x \neq y$ , while  $x \ll y$  will indicate that  $y - x \in \text{int}K$ , where  $\text{int}K$  stands for the interior of  $K$ . If  $\text{int}K \neq \emptyset$ , then  $K$  is said to be a solid cone. Write  $\|\cdot\|$  as the norm on  $\mathcal{A}$ . A cone  $K$  is called normal if there is a number  $M > 0$  such that for all  $x, y \in \mathcal{A}$ ,  $\theta \preceq x \preceq y$  implies  $\|x\| \leq M\|y\|$ . The least positive number satisfying above is called the normal constant of  $K$ .

In the sequel we always suppose that  $\mathcal{A}$  is a Banach algebra with a unit  $e$ ,  $K$  is a solid cone in  $\mathcal{A}$ , and  $\preceq$  is a partial ordering with respect to  $K$ .

**Definition 1.1.**([13]) Let  $X$  be a nonempty set and  $\mathcal{A}$  a Banach algebra. Suppose that the mapping  $d : X \times X \rightarrow \mathcal{A}$  satisfies:

- (i)  $\theta \prec d(x, y)$  for all  $x, y \in X$  with  $x \neq y$  and  $d(x, y) = \theta$  if and only if  $x = y$ ;
- (ii)  $d(x, y) = d(y, x)$  for all  $x, y \in X$ ;

(iii)  $d(x, y) \preceq d(x, z) + d(z, y)$  for all  $x, y, z \in X$ .

Then  $d$  is called a cone metric on  $X$ , and  $(X, d)$  is called a cone metric space over Banach algebra.

**Definition 1.2.**([15]) Let  $(X, d)$  be a cone metric space,  $x \in X$  and  $\{x_n\}$  a sequence in  $X$ . Then

(i)  $\{x_n\}$  converges to  $x$  whenever for every  $c \gg \theta$  there is a natural number  $N$  such that  $d(x_n, x) \ll c$  for all  $n \geq N$ . We denote this by  $\lim_{n \rightarrow \infty} x_n = x$  or  $x_n \rightarrow x$  ( $n \rightarrow \infty$ ).

(ii)  $\{x_n\}$  is a Cauchy sequence whenever for each  $c \gg \theta$  there is a natural number  $N$  such that  $d(x_n, x_m) \ll c$  for all  $n, m \geq N$ .

(iii)  $(X, d)$  is a complete cone metric space if every Cauchy sequence is convergent.

**Definition 1.3.** Let  $(X, d)$  be a cone metric space,  $\{y_n\}$  a sequence in  $X$  and  $T$  a self-map of  $X$ . Let  $x_0$  be a point of  $X$ ,  $x_{n+1} = Tx_n$  a Picard's iteration in  $X$ . The iteration procedure  $x_{n+1} = Tx_n$  is said to be  $T$ -stable with respect to  $T$  if  $\{x_n\}$  converges to a fixed point  $q$  of  $T$ , and for each  $c \gg \theta$ , there exists a natural number  $N$  such that  $d(y_{n+1}, Ty_n) \ll c$  for all  $n > N$ , then  $\lim_{n \rightarrow \infty} y_n = q$ .

**Remark 1.4.** Comparing Definition 2.1 of [18] and Definition 1.3, we find that, the conditions of the former are stronger than the latter. Actually, if  $\lim_{n \rightarrow \infty} d(y_{n+1}, Ty_n) = \theta$ , then we must have that for each  $c \gg \theta$  there exists an integer  $N > 0$  such that  $d(y_{n+1}, Ty_n) \ll c$  for all  $n > N$ . But the contrary is not true (see [6]).

**Lemma 1.5.**([6]) Let  $u, v, w \in \mathcal{A}$ . If  $u \preceq v$  and  $v \ll w$ , then  $u \ll w$ .

**Lemma 1.6.**([6]) Let  $\mathcal{A}$  be a Banach algebra and  $\{a_n\}$  a sequence in  $\mathcal{A}$ . If  $a_n \rightarrow \theta$  ( $n \rightarrow \infty$ ), then for any  $c \gg \theta$ , there exists  $N$  such that for all  $n > N$ , one has  $a_n \ll c$ .

**Lemma 1.7.**([16]) Let  $\mathcal{A}$  be a Banach algebra with a unit  $e$ ,  $x \in \mathcal{A}$ , then the limit  $\lim_{n \rightarrow \infty} \|x^n\|^{\frac{1}{n}}$  exists and the spectral radius  $\rho(x)$  satisfies

$$\rho(x) = \lim_{n \rightarrow \infty} \|x^n\|^{\frac{1}{n}} = \inf \|x^n\|^{\frac{1}{n}}.$$

If  $\rho(x) < |\lambda|$ , then  $\lambda e - x$  is invertible in  $\mathcal{A}$ , moreover,

$$(\lambda e - x)^{-1} = \sum_{i=0}^{\infty} \frac{x^i}{\lambda^{i+1}},$$

where  $\lambda$  is a complex constant.

**Lemma 1.8.**([16]) Let  $\mathcal{A}$  be a Banach algebra with a unit  $e$ ,  $a, b \in \mathcal{A}$ . If  $a$  commutes with  $b$ , then

$$\rho(a + b) \leq \rho(a) + \rho(b), \quad \rho(ab) \leq \rho(a)\rho(b).$$



## 2 Main results

In this section we give some basic but important properties, which will be used constantly in the sequel. Moreover, we introduce a class of contractive mappings with some generalized Lipschitz constants and prove the existence of non-normal solid cone and several fixed point theorems based on them without the assumption of normalities of cones. In addition, we obtain the fixed point periodic property and  $T$ -stability of Picard's iteration. All results greatly generalize the main assertions of [13-15] and [17-21]. Further, we display an example to illustrate the applications. In the end, we give another example to claim that our results in the setting of cone metric spaces over Banach algebras are never equivalent to those in usual metric spaces.

**Lemma 2.1.** Let  $\mathcal{A}$  be a Banach algebra and  $k \in \mathcal{A}$ . If  $\rho(k) < 1$ , then  $\lim_{n \rightarrow \infty} \|k^n\| = 0$ .

**Proof.** Since  $\rho(k) = \lim_{n \rightarrow \infty} \|k^n\|^{\frac{1}{n}} < 1$ , then there exists  $\alpha > 0$  such that  $\lim_{n \rightarrow \infty} \|k^n\|^{\frac{1}{n}} < \alpha < 1$ . Letting  $n$  be big enough, we obtain  $\|k^n\|^{\frac{1}{n}} \leq \alpha$ , then  $\|k^n\| \leq \alpha^n \rightarrow 0$  ( $n \rightarrow \infty$ ). Hence  $\|k^n\| \rightarrow 0$  ( $n \rightarrow \infty$ ). □

**Lemma 2.2.** Let  $\mathcal{A}$  be a Banach algebra with a unit  $e$ ,  $\{x_n\}$  a sequence in  $\mathcal{A}$ . If  $x_n$  converges to  $x$  in  $\mathcal{A}$ , and for any  $n \geq 1$ ,  $x_n$  commutes with  $x$ , then  $\rho(x_n) \rightarrow \rho(x)$  as  $n \rightarrow \infty$ .

**Proof.** Since  $x_n$  commutes with  $x$ , then it follows by Lemma 1.8 that

$$\begin{aligned} \rho(x_n) &\leq \rho(x_n - x) + \rho(x) \Rightarrow \rho(x_n) - \rho(x) \leq \rho(x_n - x), \\ \rho(x) &\leq \rho(x - x_n) + \rho(x_n) \Rightarrow \rho(x) - \rho(x_n) \leq \rho(x - x_n), \end{aligned}$$

thus

$$|\rho(x_n) - \rho(x)| \leq \rho(x_n - x) \leq \|x_n - x\| \Rightarrow \rho(x_n) \rightarrow \rho(x) (n \rightarrow \infty).$$

□

**Lemma 2.3.** Let  $\mathcal{A}$  be a Banach algebra with a unit  $e$  and  $K$  be a solid cone in  $\mathcal{A}$ . Let  $\{a_n\}$  and  $\{c_n\}$  be two sequences in  $K$  satisfying the following inequality:

$$a_{n+1} \preceq ha_n + c_n, \tag{2.1}$$

where  $h \in K$  and  $\rho(h) < 1$ . If for each  $c \gg \theta$ , there exists  $N$  such that  $c_n \ll c$  for all  $n > N$ , then  $a_n \ll c$  ( $n > N$ ).

**Proof.** By virtue of  $\rho(h) < 1$ , it follows by Lemma 1.7,  $e - h$  is invertible and  $(e - h)^{-1} = \sum_{i=0}^{\infty} h^i$ . Moreover, by Lemma 2.1, it establishes  $\|h^n\| \rightarrow 0$  ( $n \rightarrow \infty$ ). Assume  $c \gg \theta$  be arbitrary. Then there exists  $N_1$  such that for all  $n > N_1$ , we have

$$c_n \ll \frac{(e - h)c}{2}. \tag{2.2}$$

Since

$$\|h^{n-N_1}a_{N_1+1}\| \leq \|h^{n-N_1}\| \|a_{N_1+1}\| \rightarrow 0 \quad (n \rightarrow \infty),$$

thus there is  $N_2$  such that for all  $n > N_2$ , it satisfies

$$h^{n-N_1}a_{N_1+1} \ll \frac{c}{2}. \tag{2.3}$$

Put  $N = \max\{N_1, N_2\}$ , then for all  $n > N$ , both (2.2) and (2.3) are satisfied. Taking advantage of (2.1), we speculate that

$$\begin{aligned} a_{n+1} - ha_n &\preceq c_n, \\ ha_n - h^2a_{n-1} &\preceq hc_{n-1}, \\ h^2a_{n-1} - h^3a_{n-2} &\preceq h^2c_{n-2}, \\ &\dots\dots\dots \end{aligned}$$

$$h^{n-N_1-1}a_{N_1+2} - h^{n-N_1}a_{N_1+1} \preceq h^{n-N_1-1}c_{N_1+1}.$$

Combine with the above terms, for all  $n > N$ , it follows that

$$\begin{aligned} a_{n+1} &\preceq h^{n-N_1}a_{N_1+1} + c_n + hc_{n-1} + h^2c_{n-2} + \dots + h^{n-N_1-1}c_{N_1+1} \\ &\ll \frac{c}{2} + (e + h + h^2 + \dots + h^{n-N_1-1}) \cdot \frac{(e - h)c}{2} \\ &\preceq \frac{c}{2} + (e - h)^{-1} \cdot \frac{(e - h)c}{2} = c. \end{aligned}$$

□

**Remark 2.4.** Lemma 2.3 greatly generalizes Lemma 1 of [17] and Lemma 1.5 of [18]. Virtually, we delete the normality of  $K$ . Moreover, our conditions are weaker than them. Indeed, if  $a_n \rightarrow \theta$  ( $n \rightarrow \infty$ ), then  $a_n \ll c$  ( $n > N$ ). But the converse is not true (see [6]). Further, if  $\|k\| < 1$ , it is natural that  $\rho(k) < 1$ . Yet, the converse is not true.

**Theorem 2.5.** Let  $(X, d)$  be a cone metric space over Banach algebra  $\mathcal{A}$  and  $K$  be a solid cone in  $\mathcal{A}$ . Suppose that the mapping  $T : X \rightarrow X$  satisfies the following contractive condition:

$$d(Tx, Ty) \preceq k_1d(x, y) + k_2d(x, Tx) + k_3d(y, Ty) + k_4d(x, Ty) + k_5d(y, Tx),$$

for all  $x, y \in X$ , where  $k_i \in K (i = 1, \dots, 5)$  are generalized Lipschitz constants with  $\rho(k_1) + \rho(k_2 + k_3 + k_4 + k_5) < 1$ . If  $k_1$  commutes with  $k_2 + k_3 + k_4 + k_5$ , then there exists a sequence  $\{x_n\}$  in  $X$  such that it is a Cauchy sequence. Moreover, if  $\{d(x_n, y_n)\}$  converges to some non-zero element in  $\mathcal{A}$  for any two different Cauchy sequence  $\{x_n\}$  and  $\{y_n\}$ , then  $K$  is a non-normal cone.

**Proof.** Fix  $x_0 \in X$  and set  $x_{n+1} = Tx_n = T^{n+1}x_0$ . Then we have

$$\begin{aligned} d(x_{n+1}, x_n) &= d(Tx_n, Tx_{n-1}) \\ &\preceq k_1d(x_n, x_{n-1}) + k_2d(x_n, Tx_n) + k_3d(x_{n-1}, Tx_{n-1}) \\ &\quad + k_4d(x_n, Tx_{n-1}) + k_5d(x_{n-1}, Tx_n) \\ &= k_1d(x_n, x_{n-1}) + k_2d(x_n, x_{n+1}) + k_3d(x_{n-1}, x_n) \\ &\quad + k_5d(x_{n-1}, x_{n+1}) \\ &\preceq (k_1 + k_3 + k_5)d(x_n, x_{n-1}) + (k_2 + k_5)d(x_{n+1}, x_n). \end{aligned} \tag{2.4}$$

We also have

$$\begin{aligned} d(x_{n+1}, x_n) &= d(Tx_n, Tx_{n-1}) = d(Tx_{n-1}, Tx_n) \\ &\preceq k_1d(x_{n-1}, x_n) + k_2d(x_{n-1}, Tx_{n-1}) + k_3d(x_n, Tx_n) \\ &\quad + k_4d(x_{n-1}, Tx_n) + k_5d(x_n, Tx_{n-1}) \\ &= k_1d(x_n, x_{n-1}) + k_2d(x_{n-1}, x_n) + k_3d(x_n, x_{n+1}) \\ &\quad + k_4d(x_{n-1}, x_{n+1}) \\ &\preceq (k_1 + k_2 + k_4)d(x_n, x_{n-1}) + (k_3 + k_4)d(x_{n+1}, x_n). \end{aligned} \tag{2.5}$$

Add up (2.4) and (2.5) yields that

$$\begin{aligned} 2d(x_{n+1}, x_n) &\preceq (2k_1 + k_2 + k_3 + k_4 + k_5)d(x_n, x_{n-1}) \\ &\quad + (k_2 + k_3 + k_4 + k_5)d(x_{n+1}, x_n), \end{aligned}$$

which establishes that

$$(2e - k_2 - k_3 - k_4 - k_5)d(x_{n+1}, x_n) \preceq (2k_1 + k_2 + k_3 + k_4 + k_5)d(x_n, x_{n-1}).$$

Put  $k = k_2 + k_3 + k_4 + k_5$ , then

$$(2e - k)d(x_{n+1}, x_n) \preceq (2k_1 + k)d(x_n, x_{n-1}). \tag{2.6}$$

Since  $\rho(k) \leq \rho(k_1) + \rho(k) < 1 < 2$ , then by Lemma 1.7 it follows that  $2e - k$  is invertible. Furthermore,

$$(2e - k)^{-1} = \sum_{i=0}^{\infty} \frac{k^i}{2^{i+1}}.$$

By multiplying in both sides of (2.6) by  $(2e - k)^{-1}$ , we arrive at

$$d(x_{n+1}, x_n) \preceq (2e - k)^{-1}(2k_1 + k)d(x_n, x_{n-1}). \tag{2.7}$$

Denote  $h = (2e - k)^{-1}(2k_1 + k)$ , then by (2.7) we get

$$d(x_{n+1}, x_n) \preceq hd(x_n, x_{n-1}) \preceq \cdots \preceq h^n d(x_1, x_0).$$

By Lemma 1.8 we conclude that

$$\rho\left(\sum_{i=0}^n \frac{k^i}{2^{i+1}}\right) \leq \sum_{i=0}^n \rho\left(\frac{k^i}{2^{i+1}}\right) \leq \sum_{i=0}^n \frac{[\rho(k)]^i}{2^{i+1}},$$

which implies by Lemma 2.2 that

$$\rho\left(\sum_{i=0}^{\infty} \frac{k^i}{2^{i+1}}\right) \leq \sum_{i=0}^{\infty} \frac{[\rho(k)]^i}{2^{i+1}}.$$

Since  $k_1$  commutes with  $k$ , it follows that

$$\begin{aligned} (2e - k)^{-1}(2k_1 + k) &= \left(\sum_{i=0}^{\infty} \frac{k^i}{2^{i+1}}\right)(2k_1 + k) = 2\left(\sum_{i=0}^{\infty} \frac{k^i}{2^{i+1}}\right)k_1 + \sum_{i=0}^{\infty} \frac{k^{i+1}}{2^{i+1}} \\ &= 2k_1\left(\sum_{i=0}^{\infty} \frac{k^i}{2^{i+1}}\right) + k\sum_{i=0}^{\infty} \frac{k^i}{2^{i+1}} = (2k_1 + k)\left(\sum_{i=0}^{\infty} \frac{k^i}{2^{i+1}}\right) = (2k_1 + k)(2e - k)^{-1}, \end{aligned}$$

that is to say,  $(2e - k)^{-1}$  commutes with  $2k_1 + k$ . Then by Lemma 1.8 we gain

$$\begin{aligned} \rho(h) &= \rho\left((2e - k)^{-1}(2k_1 + k)\right) \leq \rho\left((2e - k)^{-1}\right)\rho(2k_1 + k) \\ &\leq \rho\left(\sum_{i=0}^{\infty} \frac{k^i}{2^{i+1}}\right)[2\rho(k_1) + \rho(k)] \leq \left(\sum_{i=0}^{\infty} \frac{[\rho(k)]^i}{2^{i+1}}\right)[2\rho(k_1) + \rho(k)] \\ &= \frac{1}{2 - \rho(k)}[2\rho(k_1) + \rho(k)] < 1, \end{aligned}$$

which establishes that  $e - h$  is invertible and  $\|h^m\| \rightarrow 0$  ( $m \rightarrow \infty$ ). Thus for all  $n > m$ ,

$$\begin{aligned} d(x_n, x_m) &\preceq d(x_n, x_{n-1}) + d(x_{n-1}, x_{n-2}) + \cdots + d(x_{m+1}, x_m) \\ &\preceq (h^{n-1} + h^{n-2} + \cdots + h^m)d(x_1, x_0) \\ &= (h^{n-m-1} + h^{n-m-2} + \cdots + h + e)h^m d(x_1, x_0) \\ &\preceq \left(\sum_{i=0}^{\infty} h^i\right)h^m d(x_1, x_0) \\ &= (e - h)^{-1}h^m d(x_1, x_0). \end{aligned}$$

Owing to

$$\|(e - h)^{-1}h^m d(x_1, x_0)\| \leq \|(e - h)^{-1}\| \|h^m\| \|d(x_1, x_0)\| \rightarrow 0 \quad (m \rightarrow \infty),$$

we have  $(e - h)^{-1}h^m d(x_1, x_0) \rightarrow \theta$  ( $m \rightarrow \infty$ ). So by using Lemma 1.5 and 1.6, we easily see that  $\{x_n\}$  is a Cauchy sequence in  $X$ .

Now we take  $y_0 \in X$  such that  $y_0 \neq x_0$ . Using the same method as the above mentioned, we can show that  $\{y_n\}$  is also a Cauchy sequence if  $y_{n+1} = Ty_n = T^{n+1}y_0$ . In the following we suppose the contrary, that is,  $K$  is normal. We shall prove that  $\{d(x_n, y_n)\}$  is convergent in  $(\mathcal{A}, \|\cdot\|)$  if  $K$  is a normal cone with normal constant  $M$ . In fact, in view of the completeness of  $(\mathcal{A}, \|\cdot\|)$ , it will be enough to show that the sequence  $\{d(x_n, y_n)\}$  is a Cauchy sequence. To this end, let  $\varepsilon > 0$  and choose  $c \gg \theta$  and  $\|c\| < \frac{\varepsilon}{4M+2}$ . Since  $\{x_n\}$  and  $\{y_n\}$  are Cauchy sequences, there is  $N$  such that  $d(x_n, x_m) \ll c$  and  $d(y_n, y_m) \ll c$  for all  $n, m > N$ . It is clear that

$$d(x_n, y_n) \preceq d(x_n, x_m) + d(x_m, y_m) + d(y_m, y_n) \preceq d(x_m, y_m) + 2c, \tag{2.8}$$

$$d(x_m, y_m) \preceq d(x_m, x_n) + d(x_n, y_n) + d(y_n, y_m) \preceq d(x_n, y_n) + 2c. \tag{2.9}$$

It follows immediately from (2.8) and (2.9) that

$$\theta \preceq d(x_m, y_m) + 2c - d(x_n, y_n) \preceq d(x_n, y_n) + 2c + 2c - d(x_n, y_n) = 4c. \tag{2.10}$$

By virtue of the normality of  $K$ , (2.10) means that

$$\|d(x_m, y_m) + 2c - d(x_n, y_n)\| \leq 4M\|c\|.$$

Hence, it ensures us that

$$\|d(x_m, y_m) - d(x_n, y_n)\| \leq \|d(x_m, y_m) + 2c - d(x_n, y_n)\| + \|2c\| \leq (4M + 2)\|c\| < \varepsilon,$$

which implies that  $\{d(x_n, y_n)\}$  is Cauchy and hence convergent.

Next, put  $\lim_{n \rightarrow \infty} d(x_n, y_n) = a$ , it is evident that  $\theta \preceq a$ . Finally, we claim that  $a = \theta$ . Actually, if there exists  $n_0 \in \mathbb{N}$  such that  $x_{n_0} = y_{n_0}$ , the claim is clear. Without loss of

generality, we suppose that  $x_n \neq y_n$  for all  $n \in \mathbb{N}$ . Notice that

$$\begin{aligned} d(x_{n+1}, y_{n+1}) &= d(Tx_n, Ty_n) \\ &\preceq k_1d(x_n, y_n) + k_2d(x_n, Tx_n) + k_3d(y_n, Ty_n) \\ &\quad + k_4d(x_n, Ty_n) + k_5d(y_n, Tx_n) \\ &= k_1d(x_n, y_n) + k_2d(x_n, x_{n+1}) + k_3d(y_n, y_{n+1}) \\ &\quad + k_4d(x_n, y_{n+1}) + k_5d(y_n, x_{n+1}) \\ &\preceq (k_1 + k_4 + k_5)d(x_n, y_n) + (k_2 + k_5)d(x_n, x_{n+1}) + (k_3 + k_4)d(y_n, y_{n+1}). \end{aligned}$$

Taking the limit as  $n \rightarrow \infty$ , we obtain that

$$a \preceq (k_1 + k_4 + k_5)a.$$

Set  $\lambda = k_1 + k_4 + k_5$ , then it follows that

$$a \preceq \lambda a \preceq \cdots \preceq \lambda^n a.$$

Because  $\lambda \preceq k_1 + k$  leads to  $\lambda^n \preceq (k_1 + k)^n$ , moreover, by Lemma 2.1,  $\rho(k_1 + k) \leq \rho(k_1) + \rho(k) < 1$  leads to  $(k_1 + k)^n \rightarrow \theta$  ( $n \rightarrow \infty$ ), we claim that, for each  $c \gg \theta$ , there exists  $n_0(c)$  such that  $\lambda^n \ll c$  such that for all  $n > n_0(c)$ . Consequently,  $a = \theta$ , a contradiction.  $\square$

It is clear that if  $T$  is a map which has a fixed point  $u$ , then  $u$  is also a fixed point of  $T^n$  for each  $n \in \mathbb{N}$ . It is well known that the converse is not true. If a map  $T$  satisfies  $F(T) = F(T^n)$  for each  $n \in \mathbb{N}$ , where  $F(T)$  stands for the set of all fixed points of  $T$ , then it is said to have a property  $P$  (see [19-21]). The following results are generalizations of the corresponding results in metric and cone metric spaces (see [20-21]). It will be deduced also without using normality of the cone.

**Theorem 2.6.** Let  $(X, d)$  be a complete cone metric space over Banach algebra  $\mathcal{A}$  and  $K$  be a solid cone in  $\mathcal{A}$ . Let  $T : X \rightarrow X$  be a mapping such that  $F(T) \neq \emptyset$  and that

$$d(Tx, T^2x) \preceq kd(x, Tx) \tag{2.11}$$

for all  $x \in X$ , where  $k \in K$  is a generalized Lipschitz constant with  $\rho(k) < 1$ . Then  $T$  has a property  $P$ .

**Proof.** We will always assume that  $n > 1$ , since the statement for  $n = 1$  is trivial. Let  $z \in F(T^n)$ . By the hypotheses, it is clear that

$$\begin{aligned} d(z, Tz) &= d(TT^{n-1}z, T^2T^{n-1}z) \preceq kd(T^{n-1}z, T^n z) = kd(TT^{n-2}z, T^2T^{n-2}z) \\ &\preceq k^2d(T^{n-2}z, T^{n-1}z) \preceq \cdots \preceq k^nd(z, Tz). \end{aligned}$$

On account of  $\rho(k) < 1$ , it follows by Lemma 2.1 that  $\|k^n\| \rightarrow 0$  ( $n \rightarrow \infty$ ). Thus  $\|k^nd(z, Tz)\| \leq \|k^n\| \|d(z, Tz)\| \rightarrow 0$  ( $n \rightarrow \infty$ ). Hence  $d(z, Tz) = \theta$ , that is.,  $Tz = z$ .  $\square$

**Theorem 2.7.** Let  $(X, d)$  be a complete cone metric space over Banach algebra  $\mathcal{A}$  and  $K$  be a solid cone in  $\mathcal{A}$ . Suppose that the mapping  $T : X \rightarrow X$  satisfies the following contractive condition:

$$d(Tx, Ty) \preceq k_1d(x, y) + k_2d(x, Tx) + k_3d(y, Ty) + k_4d(x, Ty) + k_5d(y, Tx),$$

for all  $x, y \in X$ , where  $k_i \in K$  ( $i = 1, \dots, 5$ ) are generalized Lipschitz constants with  $\rho(k_1) + \rho(k_2 + k_3 + k_4 + k_5) < 1$ . If  $k_1$  commutes with  $k_2 + k_3 + k_4 + k_5$ , then  $T$  has a unique fixed point in  $X$ . Moreover, for arbitrary  $x \in X$ , iterative sequence  $\{T^n x\}$  converges to the fixed point. Further,  $T$  has a property  $P$ .

**Proof.** By using Theorem 2.5, we obtain  $\{x_n\}$  is a Cauchy sequence in  $X$ . Since  $X$  is complete, there exists  $x^* \in X$  such that  $x_n \rightarrow x^*$  as  $n \rightarrow \infty$ . We shall prove  $x^*$  is the fixed point of  $T$ . To this end, on the one hand, we have

$$\begin{aligned} d(Tx^*, x^*) &\preceq d(Tx^*, Tx_n) + d(Tx_n, x^*) \\ &\preceq k_1d(x^*, x_n) + k_2d(x^*, Tx^*) + k_3d(x_n, Tx_n) \\ &\quad + k_4d(x^*, Tx_n) + k_5d(x_n, Tx^*) + d(Tx_n, x^*) \\ &= k_1d(x^*, x_n) + k_2d(x^*, Tx^*) + k_3d(x_n, x_{n+1}) \\ &\quad + (e + k_4)d(x^*, x_{n+1}) + k_5d(x_n, Tx^*) \\ &\preceq (k_1 + k_3 + k_5)d(x_n, x^*) + (k_2 + k_5)d(Tx^*, x^*) \\ &\quad + (e + k_3 + k_4)d(x_{n+1}, x^*), \end{aligned}$$

which implies that

$$(e - k_2 - k_5)d(Tx^*, x^*) \preceq (k_1 + k_3 + k_5)d(x_n, x^*) + (e + k_3 + k_4)d(x_{n+1}, x^*). \tag{2.12}$$

On the other hand, we obtain

$$\begin{aligned}
 d(Tx^*, x^*) &\preceq d(Tx_n, Tx^*) + d(Tx_n, x^*) \\
 &\preceq k_1d(x_n, x^*) + k_2d(x_n, Tx_n) + k_3d(x^*, Tx^*) \\
 &\quad + k_4d(x_n, Tx^*) + k_5d(x^*, Tx_n) + d(Tx_n, x^*) \\
 &= k_1d(x_n, x^*) + k_2d(x_n, x_{n+1}) + k_3d(x^*, Tx^*) \\
 &\quad + (e + k_5)d(x^*, x_{n+1}) + k_4d(x_n, Tx^*) \\
 &\preceq (k_1 + k_2 + k_4)d(x_n, x^*) + (k_3 + k_4)d(Tx^*, x^*) \\
 &\quad + (e + k_2 + k_5)d(x_{n+1}, x^*),
 \end{aligned}$$

which means that

$$(e - k_3 - k_4)d(Tx^*, x^*) \preceq (k_1 + k_2 + k_4)d(x_n, x^*) + (e + k_2 + k_5)d(x_{n+1}, x^*). \tag{2.13}$$

Combining (2.12) and (2.13) yields that

$$\begin{aligned}
 &(2e - k_2 - k_3 - k_4 - k_5)d(Tx^*, x^*) \\
 &\preceq (2k_1 + k_2 + k_3 + k_4 + k_5)d(x_n, x^*) \\
 &\quad + (2e + k_2 + k_3 + k_4 + k_5)d(x_{n+1}, x^*).
 \end{aligned}$$

Denote  $k = k_2 + k_3 + k_4 + k_5$ , then

$$(2e - k)d(Tx^*, x^*) \preceq (2k_1 + k)d(x_n, x^*) + (2e + k)d(x_{n+1}, x^*).$$

Consequently, we deduce that

$$\begin{aligned}
 d(Tx^*, x^*) &\preceq (2e - k)^{-1}(2k_1 + k)d(x_n, x^*) \\
 &\quad + (2e - k)^{-1}(2e + k)d(x_{n+1}, x^*).
 \end{aligned}$$

In view of  $x_n \rightarrow x^*$  ( $n \rightarrow \infty$ ), then for each  $\frac{c}{m} \gg \theta$  ( $m = 1, 2, \dots$ ), there exists  $N_m$  such that for all  $n > N_m$ , one has  $d(x_n, x^*) \ll \frac{c}{m}$ . Hence we speculate that

$$\begin{aligned}
 d(Tx^*, x^*) &\preceq (2e - k)^{-1}(2k_1 + k)d(x_n, x^*) + (2e - k)^{-1}(2e + k)d(x_{n+1}, x^*) \\
 &\ll [(2e - k)^{-1}(2k_1 + k) + (2e - k)^{-1}(2e + k)]\frac{c}{m} \rightarrow \theta \quad (m \rightarrow \infty),
 \end{aligned}$$

which follows that  $Tx^* = x^*$ .



In the following we shall show the fixed point is unique. Actually, if there is another fixed point  $y^*$ , then

$$\begin{aligned} d(x^*, y^*) &\preceq k_1d(x^*, y^*) + k_2d(x^*, Tx^*) + k_3d(y^*, Ty^*) \\ &\quad + k_4d(x^*, Ty^*) + k_5d(y^*, Tx^*) \\ &= (k_1 + k_4 + k_5)d(x^*, y^*). \end{aligned}$$

Set  $\lambda = k_1 + k_4 + k_5$ , then it follows that

$$d(x^*, y^*) \preceq \lambda d(x^*, y^*) \preceq \dots \preceq \lambda^n d(x^*, y^*).$$

By the proof of Theorem 2.5, we claim that, for each  $c \gg \theta$ , there exists  $n_0(c)$  such that  $\lambda^n \ll c$  such that for all  $n > n_0(c)$ . Consequently,  $d(x^*, y^*) = \theta$ , that is,  $x^* = y^*$ .

Finally, we shall prove that  $T$  has a property  $P$ . We have to show that the mapping  $T$  satisfies the condition (2.11). Indeed, firstly we have

$$\begin{aligned} d(Tx, T^2x) &= d(Tx, TTx) \\ &\preceq k_1d(x, Tx) + k_2d(x, Tx) + k_3d(Tx, T^2x) \\ &\quad + k_4d(x, T^2x) + k_5d(Tx, Tx) \\ &\preceq k_1d(x, Tx) + k_2d(x, Tx) + k_3d(Tx, T^2x) \\ &\quad + k_4d(x, Tx) + k_4d(Tx, T^2x), \end{aligned}$$

that is,

$$(e - k_3 - k_4)d(Tx, T^2x) \preceq (k_1 + k_2 + k_4)d(x, Tx). \tag{2.14}$$

Also, we have

$$\begin{aligned} d(Tx, T^2x) &= d(TTx, Tx) \\ &\preceq k_1d(Tx, x) + k_2d(Tx, T^2x) + k_3d(x, Tx) \\ &\quad + k_4d(Tx, Tx) + k_5d(x, T^2x) \\ &\preceq k_1d(x, Tx) + k_2d(Tx, T^2x) + k_3d(x, Tx) \\ &\quad + k_5d(x, Tx) + k_5d(Tx, T^2x), \end{aligned}$$

that is,

$$(e - k_2 - k_5)d(Tx, T^2x) \preceq (k_1 + k_3 + k_5)d(x, Tx). \tag{2.15}$$

Adding (2.14) and (2.15) we obtain

$$(2e - k)d(Tx, T^2x) \preceq (2k_1 + k)d(x, Tx).$$

According to the above proof, we demonstrate that

$$d(Tx, T^2x) \preceq hd(x, Tx),$$

where  $h = (2e - k)^{-1}(2k_1 + k)$  and  $\rho(h) < 1$ . Therefore, by Theorem 2.6,  $T$  has a property  $P$ . □

**Theorem 2.8.** Let  $(X, d)$  be a complete cone metric space over Banach algebra  $\mathcal{A}$  and  $K$  be a solid cone in  $\mathcal{A}$ . Suppose that the mapping  $T : X \rightarrow X$  satisfies the following contractive condition:

$$d(Tx, Ty) \preceq k_1d(x, y) + k_2d(x, Tx) + k_3d(y, Ty) + k_4d(x, Ty) + k_5d(y, Tx),$$

for all  $x, y \in X$ , where  $k_i \in K (i = 1, \dots, 5)$  are generalized Lipschitz constants with  $\rho(k_1) + \rho(k_2 + k_3 + k_4 + k_5) < 1$ . If  $k_1$  commutes with  $k_2 + k_3 + k_4 + k_5$ , then Picard's iteration is  $T$ -stable.

**Proof.** By utilizing Theorem 2.7, we obtain that  $T$  has a unique fixed point  $q$  in  $X$ . Assume that  $\{y_n\} \subset X$  satisfies the following condition: for each  $c \gg \theta$ , there exists  $N$  such that for all  $n > N$ ,  $d(y_{n+1}, Ty_n) \ll c$ . Firstly we have

$$\begin{aligned} d(Ty_n, q) &= d(Ty_n, Tq) \\ &\preceq k_1d(y_n, q) + k_2d(y_n, Ty_n) + k_3d(q, Tq) \\ &\quad + k_4d(y_n, Tq) + k_5d(q, Ty_n) \\ &= k_1d(y_n, q) + k_2d(y_n, Ty_n) + k_4d(y_n, q) + k_5d(q, Ty_n) \\ &\preceq (k_1 + k_4)d(y_n, q) + k_2[d(y_n, q) + d(q, Ty_n)] + k_5d(q, Ty_n) \\ &= (k_1 + k_2 + k_4)d(y_n, q) + (k_2 + k_5)d(q, Ty_n). \end{aligned} \tag{2.16}$$

Secondly, we arrive at

$$\begin{aligned} d(Ty_n, q) &= d(q, Ty_n) = d(Tq, Ty_n) \\ &\preceq k_1d(q, y_n) + k_2d(q, Tq) + k_3d(y_n, Ty_n) \\ &\quad + k_4d(q, Ty_n) + k_5d(y_n, Tq) \\ &= k_1d(y_n, q) + k_3d(y_n, Ty_n) + k_4d(q, Ty_n) + k_5d(y_n, q) \\ &\preceq (k_1 + k_5)d(y_n, q) + k_3[d(y_n, q) + d(q, Ty_n)] + k_4d(q, Ty_n) \\ &= (k_1 + k_3 + k_5)d(y_n, q) + (k_3 + k_4)d(q, Ty_n). \end{aligned} \tag{2.17}$$

Adding up (2.16) and (2.17) yields that

$$2d(Ty_n, q) \preceq (2k_1 + k_2 + k_3 + k_4 + k_5)d(y_n, q) + (k_2 + k_3 + k_4 + k_5)d(q, Ty_n),$$

Denote  $k = k_2 + k_3 + k_4 + k_5$ , then we get

$$(2e - k)d(Ty_n, q) \preceq (2k_1 + k)d(y_n, q).$$

Based on the proof of Theorem 2.5, it is not hard to verify that

$$d(Ty_n, q) \preceq hd(y_n, q),$$

where  $h = (2e - k)^{-1}(2k_1 + k)$  and  $\rho(h) < 1$ .

Setting  $a_n = d(y_n, q)$  and  $c_n = d(y_{n+1}, Ty_n)$ , we claim that

$$a_{n+1} = d(y_{n+1}, q) \preceq d(y_{n+1}, Ty_n) + d(Ty_n, q) \preceq c_n + ha_n.$$

If for each  $c \gg \theta$ , there exists  $N$  such that for all  $n > N$ ,  $c_n = d(y_{n+1}, Ty_n) \ll c$ . Then, making full use of Lemma 2.3, we get  $a_n = d(y_n, q) \ll c$ , which leads to  $y_n \rightarrow q$  as  $n \rightarrow \infty$ . That is to say, the Picard's iteration is  $T$ -stable.

**Corollary 2.9.** Let  $(X, d)$  be a complete cone metric space over Banach algebra  $\mathcal{A}$  and  $K$  be a solid cone in  $\mathcal{A}$ . Suppose that the mapping  $T : X \rightarrow X$  satisfies the following contractive condition:

$$d(Tx, Ty) \preceq kd(x, y),$$

for all  $x, y \in X$ , where  $k \in K$  is a generalized Lipschitz constant with  $\rho(k) < 1$ . Then  $T$  has a unique fixed point in  $X$ . Moreover, for any  $x \in X$ , iterative sequence  $\{T^n x\}$  converges to the fixed point. Further,  $T$  has a property  $P$  and Picard's iteration is  $T$ -stable.

**Corollary 2.10.** Let  $(X, d)$  be a complete cone metric space over Banach algebra  $\mathcal{A}$  and  $K$  be a solid cone in  $\mathcal{A}$ . Suppose that the mapping  $T : X \rightarrow X$  satisfies the following contractive condition:

$$d(Tx, Ty) \preceq \frac{k}{2}[d(x, Tx) + d(y, Ty)],$$

for all  $x, y \in X$ , where  $k \in K$  is a generalized Lipschitz constant with  $\rho(k) < 1$ . Then  $T$  has a unique fixed point in  $X$ . Moreover, for every  $x \in X$ , iterative sequence  $\{T^n x\}$  converges to the fixed point. Further,  $T$  has a property  $P$  and Picard's iteration is  $T$ -stable.

**Corollary 2.11.** Let  $(X, d)$  be a complete cone metric space over Banach algebra  $\mathcal{A}$  and  $K$  be a solid cone in  $\mathcal{A}$ . Suppose that the mapping  $T : X \rightarrow X$  satisfies the following contractive condition:

$$d(Tx, Ty) \preceq \frac{k}{2}[d(x, Ty) + d(y, Tx)],$$

for all  $x, y \in X$ , where  $k \in K$  is a generalized Lipschitz constant with  $\rho(k) < 1$ . Then  $T$  has a unique fixed point in  $X$ . Moreover, for each  $x \in X$ , iterative sequence  $\{T^n x\}$  converges to the fixed point. Further,  $T$  has a property  $P$  and Picard's iteration is  $T$ -stable.

**Remark 2.12.** Throughout the conclusions above, we focus on fixed point theorems in cone metric spaces over Banach algebras instead of the theorems only in cone metric spaces. All the coefficients are vector elements and the multiplications such as  $kd(x, y)$  are vector multiplications instead of usual scalar ones, which may bring us more convenience in applications.

**Remark 2.13.** In our results such as Corollary 2.9, we only suppose the spectral radius of  $k$  is less than 1, while  $\|k\| < 1$  is not assumed. Generally speaking, it is meaningful since by Remark 2.4, the condition  $\rho(k) < 1$  is weaker than that  $\|k\| < 1$ .

**Remark 2.14.** Compared with the main results of [13] and [15], our main results in this paper deal not only with the fixed point theorems for generalized Lipschitz mappings, but also with  $P$  property and  $T$ -stability of Picard's iteration, all in the setting of cone metric spaces under the condition that the underlying cones are solid without assumption of normality. These results may be more valuable to put into use since the cones discussed are not necessarily normal under ordinary conditions. Therefore, it is an interesting thing to discuss the fixed point results in cone metric spaces over Banach algebras without the assumption that the underlying cones are normal. The following examples show that our main results will be very useful.

**Example 2.15.** Let  $\mathcal{A} = C_{\mathbb{R}}^1[0, 1]$  and define a norm on  $\mathcal{A}$  by  $\|x\| = \|x\|_{\infty} + \|x'\|_{\infty}$ . Define multiplication in  $\mathcal{A}$  as just pointwise multiplication. Then  $\mathcal{A}$  is a real Banach algebra with a unit  $e = 1$  ( $e(t) = 1$  for all  $t \in [0, 1]$ ). The set  $K = \{x \in \mathcal{A} : x(t) \geq 0 \text{ for all } t \in [0, 1]\}$  is a cone in  $\mathcal{A}$ . Moreover,  $K$  is a non-normal solid cone (see [6]). Let  $X = \{a, b, c\}$ . Define  $d : X \times X$  by  $d(a, b)(t) = d(b, a)(t) = e^t$ ,  $d(b, c)(t) = d(c, b)(t) = 2e^t$ ,  $d(c, a)(t) = d(a, c)(t) = 3e^t$  and  $d(x, x)(t) = \theta$  for all  $t \in [0, 1]$  and each  $x \in X$ . We have that  $(X, d)$  is a solid cone metric space over Banach algebra  $\mathcal{A}$ . Further, let  $T : X \rightarrow X$  be a mapping defined with  $Ta = Tb = b$ ,  $Tc = a$  and let  $k_1, k_2, k_3, k_4, k_5 \in K$  defined with  $k_1(t) = \frac{1}{3}t + \frac{1}{2}$ ,  $k_2(t) = k_3(t) = k_4(t) = k_5(t) = \frac{1}{25}$  for all  $t \in [0, 1]$ . By careful calculations one can get

that  $T$  is not a Banach contraction and all the conditions of Theorems 2.7 are fulfilled. The point  $x = b$  is the unique fixed point of  $T$ . By using Theorem 2.7 and Theorem 2.8, we can also conclude that  $T$  has a  $P$  property and Picard’s iteration is  $T$ -stable.

**Example 2.16.** Let  $\mathcal{A} = \mathbb{R}^2$  and the norm be  $\|(x_1, x_2)\| = |x_1| + |x_2|$ . Define the multiplication by

$$xy = (x_1, x_2)(y_1, y_2) = (x_1y_1, x_1y_2 + x_2y_1),$$

where  $x = (x_1, x_2), y = (y_1, y_2) \in \mathcal{A}$ . Then  $\mathcal{A}$  is a Banach algebra with a unit  $e = (1, 0)$ . Taking  $X = [0, 0.55] \times (-\infty, +\infty)$ ,  $K = \{(x_1, x_2) \in \mathcal{A} : x_1, x_2 \geq 0\}$  and

$$d(x, y) = (|x_1 - y_1|, |x_2 - y_2|) \text{ for all } x = (x_1, x_2), y = (y_1, y_2) \in X,$$

we claim that  $(X, d)$  is a cone metric space over  $\mathcal{A}$  and  $K$  is a normal solid cone with normal constant  $M = 1$ .

Define a mapping  $T : X \rightarrow X$  as

$$Tx = T(x_1, x_2) = \left( \frac{1}{2} \left( \cos \frac{x_1}{2} - |x_1 - \frac{1}{2}| \right), \arctan(1 + |x_2|) + \ln(x_1 + 1) \right).$$

By using mean value theorem of differentials, it follows that

$$\begin{aligned} d(Tx, Ty) &= d(T(x_1, x_2), T(y_1, y_2)) \\ &= \left( \left| \frac{1}{2} \left( \cos \frac{x_1}{2} - \cos \frac{y_1}{2} - |x_1 - \frac{1}{2}| + |y_1 - \frac{1}{2}| \right) \right|, \right. \\ &\quad \left. \left| \arctan(1 + |x_2|) - \arctan(1 + |y_2|) + \ln(x_1 + 1) - \ln(y_1 + 1) \right| \right) \\ &\preceq \left( \left| \frac{x_1 + y_1}{4} \right| \left| \frac{x_1 - y_1}{4} \right| + \frac{1}{2} |x_1 - y_1|, \frac{1}{2} |x_2 - y_2| + |x_1 - y_1| \right) \\ &\preceq \left( \frac{5}{8}, 1 \right) (|x_1 - y_1|, |x_2 - y_2|) \\ &= \left( \frac{5}{8}, 1 \right) d(x, y) \end{aligned}$$

for all  $x, y \in X$ . Put  $k = (\frac{5}{8}, 1)$ . Simple calculations show that all conditions of Corollary 2.9 are satisfied. Thus by Corollary 2.9,  $T$  has a unique fixed point in  $X$ . Further,  $T$  has a property  $P$  and Picard’s iteration is  $T$ -stable.

The following statement indicates our fixed point results in cone metric space over Banach algebra  $\mathcal{A}$  are not equivalent to those in metric spaces. In order to end this, put

$$d_1(x, y) = \inf_{\{u \in P : d(x, y) \preceq u\}} \|u\|, \quad d_2(x, y) = \inf \{r \in \mathbb{R} : d(x, y) \preceq re\},$$

where  $x, y \in X$  and  $e = (e_1, e_2) \in \text{int}K$ . Then by Theorem 2.2 of [10],  $d_1$  and  $d_2$  are both equivalent metrics. Hence we need to consider only one of them. Let us refer to the

metric  $d_2$ . We shall prove our conclusions are not equivalent to the well-known Banach contraction principle, which means Theorem 2.4 of [8] does not hold in the setting of cone metric spaces over Banach algebras. As a matter of fact, taking  $x' = (\frac{1}{2}, 0)$ ,  $y' = (0, 0)$ ,  $e = (1, \frac{1}{2})$ , we have

$$\begin{aligned} d_2(Tx', Ty') &= \inf \left\{ r \in \mathbb{R} : \left( \frac{1}{2} \cos \frac{1}{4} - \frac{1}{4}, \ln \frac{3}{2} \right) \preceq r \left( 1, \frac{1}{2} \right) \right\} \\ &= \max \left\{ \frac{1}{2} \cos \frac{1}{4} - \frac{1}{4}, 2 \ln \frac{3}{2} \right\} = 2 \ln \frac{3}{2} \\ &\geq \frac{1}{2} = d_2(x', y'), \end{aligned}$$

which implies that there does not exist  $\lambda \in [0, 1)$  such that

$$d_2(Tx, Ty) \leq \lambda d_2(x, y)$$

for all  $x, y \in X$ . Thus it does not satisfy the contractive condition of Banach contraction principle. That is to say, Theorem 2.4 of [8] is unsuitable for cone metric spaces over Banach algebras.

**Remark 2.17.** Since the contractive mapping in Example 2.15 is generalized Lipschitz mapping, we are easy to make a conclusion that Corollary 2.1 in [5] cannot cope with Example 2.15, which infers that the main results in the setting of cone metric spaces over Banach algebras are very meaningful.

**Remark 2.18.** In Example 2.16, we are not hard to see that the main results in this paper are indeed more different from the standard results of cone metric spaces presented in the literature. Also, Example 2.16 shows that cone metric space over Banach algebra do be a real generalization of metric space even if some works with the assumption of normal cone.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

All authors contributed equally and significantly in writing this paper. All authors read and approved the final manuscript.

## Acknowledgements

The authors would like to express their thanks to the referees for their helpful comments and suggestions. The research is partially supported by Doctoral Initial Foundation of Hanshan Normal University, China (no. QD20110920).

## References

- [1] L.-G. Huang, X. Zhang, Cone metric spaces and fixed point theorems of contractive mappings, *Journal of Mathematical Analysis and Applications*, 332(2)(2007), 1468-1476.
- [2] Z. Kadelburg, S. Radenović, V. Rakočević, Remarks on “Quasi-contraction on a cone metric space”, *Applied Mathematics Letters*, 22 (2009), 1674-1679.
- [3] H. K. Nashine, Z. Kadelburg, R. P. Pathak, S. Radenović, Coincidence and fixed point results in ordered  $G$ -cone metric spaces, *Mathematical and Computer Modelling*, 57 (2013), 701-709.
- [4] L. Gajić, V. Rakočević, Quasi-contractions on a non-normal cone metric space, *Functional Analysis and Its Applications*, 46(1)(2012), 62-65.
- [5] G.-X. Song, X. Sun, Y. Zhao, G.-T Wang, New common fixed point theorems for maps on cone metric spaces, *Applied Mathematics Letters*, 23 (2010), 1033-1037.
- [6] S. Janković, Z. Kadelburg, S. Radenović, On cone metric spaces: A survey, *Nonlinear Analysis*, 74(2011), 2591-2601.
- [7] Y.-Q Feng, W. Mao, The equivalence of cone metric spaces and metric spaces, *Fixed Point Theory*, 11(2)(2010), 259-264.
- [8] W.-S. Du, A note on cone metric fixed point theory and its equivalence, *Nonlinear Analysis*, 72 (2010), 2259-2261.
- [9] Z. Kadelburg, S. Radenović, V. Rakočević, A note on the equivalence of some metric and cone metric fixed point results, *Applied Mathematics Letters*, 24(2011), 370-374.

- [10] M. Asadi, B. E. Rhoades, H. Soleimani, Some notes on the paper “The equivalence of cone metric spaces and metric spaces”, *Fixed Point Theory and Applications*, 2012, 2012: 87.
- [11] W.-S. Du, E. Karapinar, A note on cone  $b$ -metric and its related results: generalizations or equivalence? *Fixed Point Theory and Applications*, 2013, 2013: 210.
- [12] Z. Ercan, On the end of the cone metric spaces, *Topology and its Applications*, 166(2014), 10-14.
- [13] H. Liu, S.-Y. Xu, Cone metric spaces over Banach algebras and fixed point theorems of generalized Lipschitz mappings, *Fixed Point Theory and Applications*, 2013, 2013: 320.
- [14] H. Liu, S.-Y. Xu, Fixed point theorems of quasi-contractions on cone metric spaces with Banach algebras, *Abstract and Applied Analysis*, Volume 2013, Article ID 187348, 5 pages, <http://dx.doi.org/10.1155/2013/187348>.
- [15] S.-Y. Xu, S. Radenović, Fixed point theorems of generalized Lipschitz mappings on cone metric spaces over Banach algebras without assumption of normality, *Fixed Point Theory and Applications*, 2014, 2014: 102.
- [16] W. Rudin, *Functional Analysis*, McGraw-Hill, New York, NY, USA, 2nd edition, 1991.
- [17] Y. Qing, B. E. Rhoades,  $T$ -stability of Picard iteration in metric spaces, *Fixed Point Theory and Applications*, Volume 2008, Article ID 418971, 4 pages, doi:10.1155/2008/418971.
- [18] M. Asadi, H. Soleimani, S. M. Vaezpour, B. E. Rhoades, On  $T$ -stability of Picard iteration in cone metric spaces, *Fixed Point Theory and Applications*, Volume 2009, Article ID 751090, 6 pages, doi:10.1155/2009/751090.
- [19] A. G. B. Ahmad, Z. M. Fadail, M. Abbas, Z. Kadelburg, S. Radenović, Some fixed and periodic points in abstract metric spaces, *Abstract and Applied Analysis*, Volume 2012, Article ID 908423, 15 pages, doi:10.1155/2012/908423.
- [20] G. S. Jeong, B. E. Rhoades, Maps for which  $F(T) = F(T^n)$ , *Fixed Point Theory and Applications*, 6 (2005), 87-131.



- [21] M. Abbas, B. E. Rhoades, Fixed and periodic point results in cone metric spaces, *Applied Mathematics Letters*, 22 (2009), 511-515.
- [22] S. Shukla, S. Balasubramanian, M. Pavlović, A Generalized Banach Fixed Point Theorem, *Bulletin of Malaysian Mathematical Society*, 2014, in press.

## Int-soft filters of MTL-algebras

Young Bae Jun<sup>1</sup>, Seok Zun Song<sup>2</sup>, Eun Hwan Roh<sup>3</sup> and Sun Shin Ahn<sup>4,\*</sup>

<sup>1</sup> *Department of Mathematics Education, Gyeongsang National University, Jinju 660-701, Korea*

<sup>2</sup> *Department of Mathematics, Jeju National University, Jeju 690-756, Korea*

<sup>3</sup> *Department of Mathematics Education, Chinju National University of Education, Jinju 660-756, Korea*

<sup>4</sup> *Department of Mathematics Education, Dongguk University, Seoul 100-715, Korea*

**Abstract.** The notions of (Boolean) int-soft filters in MTL-algebras are introduced, and several properties are investigated. Characterizations of (Boolean) int-soft filters are discussed, and a condition for an int-soft filter to be Boolean is provided. The extension property for a Boolean int-soft filter is constructed, and the least int-soft filter containing a given soft set is established.

### 1. Introduction

The logic MTL, Monoidal  $t$ -norm based logic, was introduced by Esteva and Godo in [3]. This logic is very interesting from many points of view. From the logic point of view, it can be regarded as a weak system of Fuzzy Logic. Indeed, it arises from Hájek's Basic Logic BL [4] by replacing the axiom

$$(A \hat{\wedge} (A \rightarrow B)) \leftrightarrow (A \wedge B)$$

by the weaker axiom

$$(A \hat{\wedge} (A \rightarrow B)) \rightarrow (A \wedge B).$$

In connection with the logic MTL, Esteva and Godo [3] introduced a new algebra, called a *MTL-algebra*, and studied several basic properties. They also introduced the notion of (prime) filters in MTL-algebras. Vetterlein [8] studied MTL-algebras arising from partially ordered groups. Borzooei, Khosravi Shoar and Americ [1] discussed some types of filters in MTL-algebras. Morton and van Alten [6] considered the algebraic semantics of the monoidal  $t$ -norm logic(MTL) with unary operations (modalities).

In this paper, we introduce the notion of (Boolean) int-soft filters in MTL-algebras, and investigate several properties. We discuss characterizations of (Boolean) int-soft filters, and provide a condition for an int-soft filter to be Boolean. We establish the extension property for a Boolean int-soft filter. We also construct the least int-soft filter containing a given soft set.

### 2. Preliminaries

---

<sup>0</sup>**2010 Mathematics Subject Classification:** 06F35; 03G25; 06D72.

<sup>0</sup>**Keywords:** (Boolean) filter; (Boolean) int-soft filter.

\* The corresponding author.

<sup>0</sup>E-mail: skywine@gmail.com (Y. B. Jun); szsong@jejunu.ac.kr (S. Z. Song); idealmath@gmail.com (E. H. Roh); sunshine@dongguk.edu (S. S. Ahn)

Young Bae Jun, Seok Zun, Eun Hwan Roh and Sun Shin Ahn

By a *residuated lattice* we shall mean a lattice  $L = (L, \leq, \wedge, \vee, \odot, \rightarrow, 0, 1)$  containing the least element 0 and the largest element 1, and endowed with two binary operations  $\odot$  (called *product*) and  $\rightarrow$  (called *residuum*) such that

- $\odot$  is associative, commutative and isotone.
- $(\forall x \in L) (x \odot 1 = x)$ .
- The Galois correspondence holds, that is,

$$(\forall x, y, z \in L) (x \odot y \leq z \iff x \leq y \rightarrow z).$$

In a residuated lattice, the following are true (see [7]):

- (a1)  $x \leq y \iff x \rightarrow y = 1$ .
- (a2)  $0 \rightarrow x = 1, 1 \rightarrow x = x, x \rightarrow (y \rightarrow x) = 1$ .
- (a3)  $y \leq (y \rightarrow x) \rightarrow x$ .
- (a4)  $x \rightarrow (y \rightarrow z) = (x \odot y) \rightarrow z = y \rightarrow (x \rightarrow z)$ .
- (a5)  $x \rightarrow y \leq (z \rightarrow x) \rightarrow (z \rightarrow y), x \rightarrow y \leq (y \rightarrow z) \rightarrow (x \rightarrow z)$ .
- (a6)  $y \leq x \implies x \rightarrow z \leq y \rightarrow z, z \rightarrow y \leq z \rightarrow x$ .
- (a7)  $\left( \bigvee_{i \in \Gamma} y_i \right) \rightarrow x = \bigwedge_{i \in \Gamma} (y_i \rightarrow x)$ .

We define  $x^* = \bigvee \{y \in L \mid x \odot y = 0\}$ , equivalently,  $x^* = x \rightarrow 0$ . Then

- (a8)  $0^* = 1, 1^* = 0, x \leq x^{**}, \text{ and } x^* = x^{***}$ .

Based on the Hájek's results [4], Axioms of MTL and Formulas which are provable in MTL, Esteva and Godo [3] defined the algebras, so called MTL-algebras, corresponding to the MTL-logic in the following way.

**Definition 2.1.** An *MTL-algebra* is a residuated lattice  $L = (L, \leq, \wedge, \vee, \odot, \rightarrow, 0, 1)$  satisfying the pre-linearity equation:

$$(x \rightarrow y) \vee (y \rightarrow x) = 1.$$

In an MTL-algebra, the following are true:

- (a9)  $x \rightarrow (y \vee z) = (x \rightarrow y) \vee (x \rightarrow z)$ .
- (a10)  $x \odot y \leq x \wedge y$ .

**Definition 2.2** ([3]). Let  $L$  be an MTL-algebra. A nonempty subset  $F$  of  $L$  is called a *filter* of  $L$  if it satisfies

- (b1)  $(\forall x, y \in F) (x \odot y \in F)$ .
- (b2)  $(\forall x \in F) (\forall y \in L) (x \leq y \implies y \in F)$ .

Since  $\wedge$  is not definable from  $\odot$  and  $\rightarrow$  in a MTL-algebra, one could consider that the further condition

- (b3)  $(\forall x, y \in F) (x \wedge y \in F)$

Int-soft filters of MTL-algebras

should be also required for a filter. However the condition (b3) is indeed redundant because it is a consequence of conditions (b1) and (b2). Namely, since  $x \odot y \leq x \wedge y$ , if  $x, y \in F$  then  $x \odot y \in F$  and thus  $x \wedge y \in F$  as well.

**Proposition 2.3.** *A nonempty subset  $F$  of an MTL-algebra  $L$  is a filter of  $L$  if and only if it satisfies:*

- (b4)  $1 \in F$ .
- (b5)  $(\forall x \in F) (\forall y \in L) (x \rightarrow y \in F \Rightarrow y \in F)$ .

A soft set theory is introduced by Molodtsov [5], and Çağman et al. [2] provided new definitions and various results on soft set theory.

Let  $\mathcal{P}(U)$  denote the power set of an initial universe set  $U$  and  $A, B, C, \dots \subseteq E$  where  $E$  is a set of parameters.

**Definition 2.4** ([2, 5]). *A soft set  $(\tilde{f}, A)$  over  $U$  is defined to be the set of ordered pairs*

$$(\tilde{f}, A) := \left\{ (x, \tilde{f}(x)) : x \in E, \tilde{f}(x) \in \mathcal{P}(U) \right\},$$

where  $\tilde{f} : E \rightarrow \mathcal{P}(U)$  such that  $\tilde{f}(x) = \emptyset$  if  $x \notin A$ .

For a soft set  $(\tilde{f}, L)$  over  $U$ , the set

$$i_L(\tilde{f}; \gamma) = \left\{ x \in L \mid \gamma \subseteq \tilde{f}(x) \right\}$$

is called the  $\gamma$ -inclusive set of  $(\tilde{f}, L)$ .

3. Int-soft filters

In what follows, we take an MTL-algebra  $L$  as a set of parameters.

**Definition 3.1.** A soft set  $(\tilde{f}, L)$  over  $U$  is called an *int-soft filter* of  $L$  if it satisfies:

$$(\forall x, y \in L) (\tilde{f}(x \odot y) \supseteq \tilde{f}(x) \cap \tilde{f}(y)). \tag{3.1}$$

$$(\forall x, y \in L) (x \leq y \Rightarrow \tilde{f}(x) \subseteq \tilde{f}(y)). \tag{3.2}$$

**Example 3.2.** Let  $L = [0, 1]$  and define a product  $\odot$  and a residuum  $\rightarrow$  on  $L$  as follows:

$$x \odot y := \begin{cases} x \wedge y & \text{if } x + y > \frac{1}{2}, \\ 0 & \text{otherwise} \end{cases}, \quad x \rightarrow y := \begin{cases} 1 & \text{if } x \leq y, \\ (0.5 - x) \vee y & \text{if } x > y \end{cases}$$

for all  $x, y \in L$ . Then  $L$  is an MTL-algebra. Let  $(\tilde{f}, L)$  be a soft set over  $U$  in which

$$\tilde{f}(x) := \begin{cases} \alpha & \text{if } x \in (0.5, 1], \\ \beta & \text{otherwise,} \end{cases}$$

where  $\alpha \supseteq \beta$  in  $\mathcal{P}(U)$ . Then it is routine to verify that  $(\tilde{f}, L)$  is an int-soft filter of  $L$ .

Young Bae Jun, Seok Zun, Eun Hwan Roh and Sun Shin Ahn

We provide characterizations of an int-soft filter.

**Theorem 3.3.** *A soft set  $(\tilde{f}, L)$  over  $U$  is an int-soft filter of  $L$  if and only if it satisfies:*

$$(\forall x \in L) \left( \tilde{f}(1) \supseteq \tilde{f}(x) \right), \tag{3.3}$$

$$(\forall x, y \in L) \left( \tilde{f}(y) \supseteq \tilde{f}(x) \cap \tilde{f}(x \rightarrow y) \right). \tag{3.4}$$

*Proof.* Assume that  $(\tilde{f}, L)$  is an int-soft filter of  $L$ . Since  $x \leq 1$  for all  $x \in L$ , it follows from (3.2) that  $\tilde{f}(x) \subseteq \tilde{f}(1)$  for all  $x \in L$ . Since  $x \leq (x \rightarrow y) \rightarrow y$ , we have  $x \odot (x \rightarrow y) \leq y$  for all  $x, y \in L$  by the Galois correspondence. It follows from (3.2) and (3.1) that

$$\tilde{f}(y) \supseteq \tilde{f}(x \odot (x \rightarrow y)) \supseteq \tilde{f}(x) \cap \tilde{f}(x \rightarrow y)$$

for all  $x, y \in L$ .

Conversely, let  $(\tilde{f}, L)$  be a soft set over  $U$  which satisfy two conditions (3.3) and (3.4). Let  $x, y \in L$  be such that  $x \leq y$ . Then  $x \rightarrow y = 1$ , and so

$$\tilde{f}(y) \supseteq \tilde{f}(x) \cap \tilde{f}(x \rightarrow y) = \tilde{f}(x) \cap \tilde{f}(1) = \tilde{f}(x),$$

for all  $x \in L$ . This proves (3.2). Using (a4), we know that

$$x \rightarrow (y \rightarrow (x \odot y)) = (x \odot y) \rightarrow (x \odot y) = 1.$$

Using (3.3) and (3.4), we have

$$\begin{aligned} \tilde{f}(x \odot y) &\supseteq \tilde{f}(y) \cap \tilde{f}(y \rightarrow (x \odot y)) \\ &\supseteq \tilde{f}(y) \cap \left( \tilde{f}(x) \cap \tilde{f}(x \rightarrow (y \rightarrow (x \odot y))) \right) \\ &= \tilde{f}(y) \cap \left( \tilde{f}(x) \cap \tilde{f}(1) \right) \\ &= \tilde{f}(x) \cap \tilde{f}(y) \end{aligned}$$

for all  $x, y \in L$ . Therefore  $(\tilde{f}, L)$  is an int-soft filter of  $L$ . □

**Theorem 3.4.** *A soft set  $(\tilde{f}, L)$  over  $U$  is an int-soft filter of  $L$  if and only if it satisfies:*

$$(\forall a, b, c \in L) \left( a \leq b \rightarrow c \Rightarrow \tilde{f}(c) \supseteq \tilde{f}(a) \cap \tilde{f}(b) \right). \tag{3.5}$$

*Proof.* Assume that  $(\tilde{f}, L)$  is an int-soft filter of  $L$ . Let  $a, b, c \in L$  be such that  $a \leq b \rightarrow c$ . Then  $\tilde{f}(a) \subseteq \tilde{f}(b \rightarrow c)$  by (3.2), and so

$$\tilde{f}(c) \supseteq \tilde{f}(b) \cap \tilde{f}(b \rightarrow c) \supseteq \tilde{f}(b) \cap \tilde{f}(a).$$

Int-soft filters of MTL-algebras

Conversely, let  $(\tilde{f}, L)$  be a soft set over  $U$  satisfying (3.5). Since  $x \leq x \rightarrow 1$  for all  $x \in L$ , it follows from (3.5) that

$$\tilde{f}(1) \supseteq \tilde{f}(x) \cap \tilde{f}(x) = \tilde{f}(x)$$

for all  $x \in L$ . Since  $x \rightarrow y \leq x \rightarrow y$  for all  $x, y \in L$ , we have

$$\tilde{f}(y) \supseteq \tilde{f}(x) \cap \tilde{f}(x \rightarrow y)$$

for all  $x, y \in L$ . Therefore  $(\tilde{f}, L)$  is an int-soft filter of  $L$ . □

**Corollary 3.5.** *A soft set  $(\tilde{f}, L)$  over  $U$  is an int-soft filter of  $L$  if and only if it satisfies the following assertion:*

$$\tilde{f}(x) \supseteq \bigcap_{k=1}^n \tilde{f}(a_k) \tag{3.6}$$

whenever  $a_n \rightarrow (\dots \rightarrow (a_2 \rightarrow (a_1 \rightarrow x)) \dots) = 1$  for every  $a_1, a_2, \dots, a_n \in L$ .

*Proof.* It is by induction. □

**Theorem 3.6.** *For a filter  $F$  of  $L$  and  $a \in L$ , let  $(\tilde{f}, L)$  be a soft set over  $U$  defined by*

$$\tilde{f}(x) := \begin{cases} \gamma_1 & \text{if } x \in \{z \in L \mid a \vee z \in F\}, \\ \gamma_2 & \text{otherwise,} \end{cases}$$

for all  $x \in L$  where  $\gamma_2 \subsetneq \gamma_1$  in  $\mathcal{P}(U)$ . Then  $(\tilde{f}, L)$  is an int-soft filter of  $L$ .

*Proof.* Since  $a \vee 1 \in F$ , we have  $1 \in \{z \in L \mid a \vee z \in F\}$  and so  $\tilde{f}(1) = \gamma_1 \supseteq \tilde{f}(x)$  for all  $x \in L$ . Now if  $y \in \{z \in L \mid a \vee z \in F\}$ , then clearly

$$\tilde{f}(y) = \gamma_1 \supseteq \tilde{f}(x) \cap \tilde{f}(x \rightarrow y).$$

Suppose that  $y \notin \{z \in L \mid a \vee z \in F\}$ . Then at least one of  $x$  and  $x \rightarrow y$  does not belong to  $\{z \in L \mid a \vee z \in F\}$ . Hence

$$\tilde{f}(y) = \gamma_2 = \tilde{f}(x) \cap \tilde{f}(x \rightarrow y),$$

and therefore  $(\tilde{f}, L)$  is an int-soft filter of  $L$ . □

**Theorem 3.7.** *A soft set  $(\tilde{f}, L)$  over  $U$  is an int-soft filter of  $L$  if and only if the nonempty  $\gamma$ -inclusive set  $i_L(\tilde{f}; \gamma)$  is a filter of  $L$  for all  $\gamma \in \mathcal{P}(U)$ .*

*Proof.* Assume that the nonempty  $\gamma$ -inclusive set  $i_L(\tilde{f}; \gamma)$  is a filter of  $L$  for all  $\gamma \in \mathcal{P}(U)$ . For any  $x \in L$ , let  $\tilde{f}(x) = \gamma$ . Then  $x \in i_L(\tilde{f}; \gamma)$ . Since  $i_L(\tilde{f}; \gamma)$  is a filter of  $L$ , we have  $1 \in i_L(\tilde{f}; \gamma)$  and so  $\tilde{f}(x) = \gamma \subseteq \tilde{f}(1)$ . For any  $x, y \in L$ , let  $\tilde{f}(x \rightarrow y) \cap \tilde{f}(x) = \gamma$ . Then  $x \rightarrow y \in i_L(\tilde{f}; \gamma)$

Young Bae Jun, Seok Zun, Eun Hwan Roh and Sun Shin Ahn

and  $x \in i_L(\tilde{f}; \gamma)$ . It follows from (b5) that  $y \in i_L(\tilde{f}; \gamma)$ . Hence  $\tilde{f}(y) \supseteq \gamma = \tilde{f}(x \rightarrow y) \cap \tilde{f}(x)$ . Therefore  $(\tilde{f}, L)$  is an int-soft filter of  $L$  by Theorem 3.3.

Conversely, suppose that  $(\tilde{f}, L)$  is an int-soft filter of  $L$ . Let  $\gamma \in \mathcal{P}(U)$  be such that  $i_L(\tilde{f}; \gamma) \neq \emptyset$ . Then there exists  $a \in i_L(\tilde{f}; \gamma)$ , and so  $\gamma \subseteq \tilde{f}(a)$ . It follows from (3.3) that  $\gamma \subseteq \tilde{f}(a) \subseteq \tilde{f}(1)$ . Thus  $1 \in i_L(\tilde{f}; \gamma)$ . Let  $x, y \in L$  be such that  $x \rightarrow y \in i_L(\tilde{f}; \gamma)$  and  $x \in i_L(\tilde{f}; \gamma)$ . Then  $\gamma \subseteq \tilde{f}(x \rightarrow y)$  and  $\gamma \subseteq \tilde{f}(x)$ . It follows from (3.4) that

$$\gamma \subseteq \tilde{f}(x \rightarrow y) \cap \tilde{f}(x) \subseteq \tilde{f}(y),$$

that is,  $y \in i_L(\tilde{f}; \gamma)$ . Thus  $i_L(\tilde{f}; \gamma) (\neq \emptyset)$  is a filter of  $L$  by Proposition 2.3. □

**Theorem 3.8.** *If  $(\tilde{f}, L)$  is an int-soft filter of  $L$ , then the set*

$$\Omega_a := \{x \in L \mid \tilde{f}(x) \supseteq \tilde{f}(a)\}$$

*is a filter of  $L$  for every  $a \in L$ .*

*Proof.* Since  $\tilde{f}(1) \supseteq \tilde{f}(x)$  for all  $x \in L$ , we have  $1 \in \Omega_a$ . Let  $x, y \in L$  be such that  $x \in \Omega_a$  and  $x \rightarrow y \in \Omega_a$ . Then  $\tilde{f}(x) \supseteq \tilde{f}(a)$  and  $\tilde{f}(x \rightarrow y) \supseteq \tilde{f}(a)$ . Since  $\tilde{f}$  is an int-soft filter of  $L$ , it follows from (3.4) that

$$\tilde{f}(y) \supseteq \tilde{f}(x) \cap \tilde{f}(x \rightarrow y) \supseteq \tilde{f}(a)$$

so that  $y \in \Omega_a$ . Hence  $\Omega_a$  is a filter of  $L$ . □

**Theorem 3.9.** *Let  $a \in L$  and let  $(\tilde{f}, L)$  be a soft set over  $U$ . Then*

(1) *If  $\Omega_a$  is a filter of  $L$ , then  $(\tilde{f}, L)$  satisfies the following implication:*

$$(\forall x, y \in L) (\tilde{f}(a) \subseteq \tilde{f}(x \rightarrow y) \cap \tilde{f}(x) \Rightarrow \tilde{f}(a) \subseteq \tilde{f}(y)). \tag{3.7}$$

(2) *If  $(\tilde{f}, L)$  satisfies (3.3) and (3.7), then  $\Omega_a$  is a filter of  $L$ .*

*Proof.* (1) Assume that  $\Omega_a$  is a filter of  $L$ . Let  $x, y \in L$  be such that

$$\tilde{f}(a) \subseteq \tilde{f}(x \rightarrow y) \cap \tilde{f}(x).$$

Then  $x \rightarrow y \in \Omega_a$  and  $x \in \Omega_a$ . Using (b5), we have  $y \in \Omega_a$  and so  $\tilde{f}(y) \supseteq \tilde{f}(a)$ .

(2) Suppose that  $\tilde{f}$  satisfies (3.3) and (3.7). From (3.3) it follows that  $1 \in \Omega_a$ . Let  $x, y \in L$  be such that  $x \in \Omega_a$  and  $x \rightarrow y \in \Omega_a$ . Then  $\tilde{f}(a) \subseteq \tilde{f}(x)$  and  $\tilde{f}(a) \subseteq \tilde{f}(x \rightarrow y)$ , which imply that  $\tilde{f}(a) \subseteq \tilde{f}(x) \cap \tilde{f}(x \rightarrow y)$ . Thus  $\tilde{f}(a) \subseteq \tilde{f}(y)$  by (3.7), and so  $y \in \Omega_a$ . Therefore  $\Omega_a$  is a filter of  $L$ . □

**Proposition 3.10.** *Let  $(\tilde{f}, L)$  be an int-soft filter of  $L$ . Then the following are equivalent:*

(1)  $(\forall x, y, z \in L) (\tilde{f}(x \rightarrow z) \supseteq \tilde{f}(x \rightarrow (y \rightarrow z)) \cap \tilde{f}(x \rightarrow y))$ .

Int-soft filters of MTL-algebras

- (2)  $(\forall x, y \in L) (\tilde{f}(x \rightarrow y) \supseteq \tilde{f}(x \rightarrow (x \rightarrow y)))$ .
- (3)  $(\forall x, y, z \in L) (\tilde{f}((x \rightarrow y) \rightarrow (x \rightarrow z)) \supseteq \tilde{f}(x \rightarrow (y \rightarrow z)))$ .

*Proof.* (1)  $\Rightarrow$  (2). Suppose that  $(\tilde{f}, L)$  satisfies the condition (1). Taking  $z = y$  and  $y = x$  in (1) and using (3.3), we have

$$\begin{aligned} \tilde{f}(x \rightarrow y) &\supseteq \tilde{f}(x \rightarrow (x \rightarrow y)) \cap \tilde{f}(x \rightarrow x) \\ &= \tilde{f}(x \rightarrow (x \rightarrow y)) \cap \tilde{f}(1) \\ &= \tilde{f}(x \rightarrow (x \rightarrow y)) \end{aligned}$$

for all  $x, y, z \in L$ .

(2)  $\Rightarrow$  (3). Suppose that  $(\tilde{f}, L)$  satisfies the condition (2) and let  $x, y, z \in L$ . Since

$$x \rightarrow (y \rightarrow z) \leq x \rightarrow ((x \rightarrow y) \rightarrow (x \rightarrow z)),$$

it follows that

$$\begin{aligned} \tilde{f}((x \rightarrow y) \rightarrow (x \rightarrow z)) &= \tilde{f}(x \rightarrow ((x \rightarrow y) \rightarrow z)) \\ &\supseteq \tilde{f}(x \rightarrow (x \rightarrow ((x \rightarrow y) \rightarrow z))) \\ &= \tilde{f}(x \rightarrow ((x \rightarrow y) \rightarrow (x \rightarrow z))) \\ &\supseteq \tilde{f}(x \rightarrow (y \rightarrow z)). \end{aligned}$$

(3)  $\Rightarrow$  (1). If  $(\tilde{f}, L)$  satisfies the condition (3), then

$$\begin{aligned} \tilde{f}(x \rightarrow y) &\supseteq \tilde{f}((x \rightarrow y) \rightarrow (x \rightarrow z)) \cap \tilde{f}(x \rightarrow y) \\ &\supseteq \tilde{f}(x \rightarrow (y \rightarrow z)) \cap \tilde{f}(x \rightarrow y). \end{aligned}$$

This completes the proof. □

**Theorem 3.11.** For a fixed element  $a \in L$ , let  $(\tilde{f}_a, L)$  be a soft set over  $U$  defined by

$$\tilde{f}_a(x) := \begin{cases} \gamma_1 & \text{if } a \leq x, \\ \gamma_2 & \text{otherwise,} \end{cases}$$

where  $\gamma_1 \supsetneq \gamma_2$  in  $\mathcal{P}(U)$ . Then  $(\tilde{f}_a, L)$  is an int-soft filter of  $L$  if and only if it satisfies the following implication:

$$(\forall x, y \in L) (a \leq y \rightarrow x, a \leq y \Rightarrow a \leq x). \tag{3.8}$$

*Proof.* Assume that  $(\tilde{f}_a, L)$  is an int-soft filter of  $L$  and let  $x, y \in L$  be such that  $a \leq y \rightarrow x$  and  $a \leq y$ . Then  $\tilde{f}_a(y \rightarrow x) = \gamma_1 = \tilde{f}_a(y)$ , and thus

$$\tilde{f}_a(x) \supseteq \tilde{f}_a(y \rightarrow x) \cap \tilde{f}_a(y) = \gamma_1$$

which implies that  $\tilde{f}_a(x) = \gamma_1$  and so  $a \leq x$ .



Young Bae Jun, Seok Zun, Eun Hwan Roh and Sun Shin Ahn

Conversely, suppose that (3.8) is valid. Note that  $i_L(\tilde{f}_a; \gamma_2) = L$  and  $i_L(\tilde{f}_a; \gamma_1) = \{x \in L \mid a \leq x\}$ . Obviously  $1 \in i_L(\tilde{f}_a; \gamma_1)$ . Let  $x, y \in L$  be such that  $x \in i_L(\tilde{f}_a; \gamma_1)$  and  $x \rightarrow y \in i_L(\tilde{f}_a; \gamma_1)$ . Then  $a \leq x$  and  $a \leq x \rightarrow y$ , which imply from (3.8) that  $a \leq y$ , that is,  $y \in i_L(\tilde{f}_a; \gamma_1)$ . Hence  $i_L(\tilde{f}_a; \gamma_1)$  is a filter of  $L$ . Using Theorem 3.7, we know that  $(\tilde{f}_a, L)$  is an int-soft filter of  $L$ .  $\square$

**Definition 3.12.** An int-soft filter  $(\tilde{f}, L)$  of  $L$  is said to be *Boolean* if it satisfies the following identity

$$(\forall x \in L) (\tilde{f}(x \vee x^*) = \tilde{f}(1)). \tag{3.9}$$

**Proposition 3.13.** Every Boolean int-soft filter  $(\tilde{f}, L)$  of  $L$  satisfies the following inclusion:

$$(\forall x, y, z \in L) (\tilde{f}(x \rightarrow z) \supseteq \tilde{f}(x \rightarrow (z^* \rightarrow y)) \cap \tilde{f}(y \rightarrow z)). \tag{3.10}$$

*Proof.* Using (a5), we have

$$y \rightarrow z \leq (z^* \rightarrow y) \rightarrow (z^* \rightarrow z) \leq (x \rightarrow (z^* \rightarrow y)) \rightarrow (x \rightarrow (z^* \rightarrow z)).$$

It follows from (3.2) that

$$\tilde{f}(y \rightarrow z) \subseteq \tilde{f}((x \rightarrow (z^* \rightarrow y)) \rightarrow (x \rightarrow (z^* \rightarrow z)))$$

so from (3.4) that

$$\begin{aligned} \tilde{f}(x \rightarrow (z^* \rightarrow z)) &\supseteq \tilde{f}(x \rightarrow (z^* \rightarrow y)) \cap \tilde{f}((x \rightarrow (z^* \rightarrow y)) \rightarrow (x \rightarrow (z^* \rightarrow z))) \\ &\supseteq \tilde{f}(x \rightarrow (z^* \rightarrow y)) \cap \tilde{f}(y \rightarrow z). \end{aligned}$$

Since

$$z^* \vee z = ((z^* \rightarrow z) \rightarrow z) \wedge ((z \rightarrow z^*) \rightarrow z^*) \leq (z^* \rightarrow z) \rightarrow z,$$

we have  $\tilde{f}((z^* \rightarrow z) \rightarrow z) \supseteq \tilde{f}(z^* \vee z) = \tilde{f}(1)$ . Since

$$x \rightarrow (z^* \rightarrow z) \leq ((z^* \rightarrow z) \rightarrow z) \rightarrow (x \rightarrow z),$$

it follows from (3.2) that

$$\tilde{f}(x \rightarrow (z^* \rightarrow z)) \subseteq \tilde{f}(((z^* \rightarrow z) \rightarrow z) \rightarrow (x \rightarrow z)).$$

Thus

$$\begin{aligned} \tilde{f}(x \rightarrow z) &\supseteq \tilde{f}((z^* \rightarrow z) \rightarrow z) \cap \tilde{f}(((z^* \rightarrow z) \rightarrow z) \rightarrow (x \rightarrow z)) \\ &\supseteq \tilde{f}(1) \cap \tilde{f}(x \rightarrow (z^* \rightarrow z)) \\ &= \tilde{f}(x \rightarrow (z^* \rightarrow z)) \\ &\supseteq \tilde{f}(x \rightarrow (z^* \rightarrow y)) \cap \tilde{f}(y \rightarrow z). \end{aligned}$$

This completes the proof.  $\square$

Int-soft filters of MTL-algebras

We provide a condition for an int-soft filter to be Boolean.

**Proposition 3.14.** *If an int-soft filter  $(\tilde{f}, L)$  of  $L$  satisfies the following inclusion*

$$(\forall x, y \in L) \left( \tilde{f}(x) \supseteq \tilde{f}((x \rightarrow y) \rightarrow x) \right), \tag{3.11}$$

*then it is Boolean.*

*Proof.* Using (a2), (a4) and (a5), we have

$$\begin{aligned} 1 &= x \rightarrow ((x^* \rightarrow x) \rightarrow x) \\ &\leq ((x^* \rightarrow x) \rightarrow x)^* \rightarrow x^* \\ &\leq (x^* \rightarrow x) \rightarrow (((x^* \rightarrow x) \rightarrow x)^* \rightarrow x) \\ &= ((x^* \rightarrow x) \rightarrow x)^* \rightarrow ((x^* \rightarrow x) \rightarrow x) \\ &= (((x^* \rightarrow x) \rightarrow x) \rightarrow 0) \rightarrow ((x^* \rightarrow x) \rightarrow x). \end{aligned}$$

It follows from (3.2), (3.3) and (3.11) that

$$\tilde{f}((x^* \rightarrow x) \rightarrow x) \supseteq \tilde{f}(((x^* \rightarrow x) \rightarrow x) \rightarrow 0) \rightarrow ((x^* \rightarrow x) \rightarrow x) = \tilde{f}(1).$$

Using (a7) and (a9), since

$$\begin{aligned} (x^* \rightarrow x) \rightarrow x &\leq ((x^* \rightarrow x) \rightarrow x) \vee ((x^* \rightarrow x) \rightarrow x^*) \\ &= (x^* \rightarrow x) \rightarrow (x \vee x^*) \\ &= (1 \wedge (x^* \rightarrow x)) \rightarrow (x \vee x^*) \\ &= ((x \rightarrow x) \wedge (x^* \rightarrow x)) \rightarrow (x \vee x^*) \\ &= ((x \vee x^*) \rightarrow x) \rightarrow (x \vee x^*), \end{aligned}$$

we get

$$\begin{aligned} \tilde{f}(1) &= \tilde{f}((x^* \rightarrow x) \rightarrow x) \\ &\subseteq \tilde{f}(((x \vee x^*) \rightarrow x) \rightarrow (x \vee x^*)) \\ &\subseteq \tilde{f}(x \vee x^*), \end{aligned}$$

and so  $\tilde{f}(x \vee x^*) = \tilde{f}(1)$ . Therefore  $(\tilde{f}, L)$  is Boolean. □

**Proposition 3.15.** *If an int-soft filter  $(\tilde{f}, L)$  of  $L$  satisfies the condition (3.10), then it satisfies the condition (3.11).*

*Proof.* Since  $(x \rightarrow y) \rightarrow x \leq x^* \rightarrow x$ , it follows from (3.2) that

$$\begin{aligned} \tilde{f}(x) &= \tilde{f}(1 \rightarrow x) \\ &\supseteq \tilde{f}(1 \rightarrow (x^* \rightarrow x^*)) \cap \tilde{f}(x^* \rightarrow x) \\ &\supseteq \tilde{f}(1) \cap \tilde{f}((x \rightarrow y) \rightarrow x) \\ &= \tilde{f}((x \rightarrow y) \rightarrow x). \end{aligned}$$

Young Bae Jun, Seok Zun, Eun Hwan Roh and Sun Shin Ahn

Hence  $(\tilde{f}, L)$  satisfies the condition (3.11). □

**Proposition 3.16.** *If an int-soft filter  $(\tilde{f}, L)$  of  $L$  satisfies (3.11), then it satisfies the following inclusion:*

$$(\forall x, y, z \in L) \left( \tilde{f}(x \rightarrow z) \supseteq \tilde{f}(x \rightarrow (y \rightarrow z)) \cap \tilde{f}(x \rightarrow y) \right). \tag{3.12}$$

*Proof.* Since  $x \rightarrow (y \rightarrow z) = y \rightarrow (x \rightarrow z) \leq (x \rightarrow y) \rightarrow (x \rightarrow (x \rightarrow z))$ , it follows from (3.2) that

$$\tilde{f}(x \rightarrow (y \rightarrow z)) \subseteq \tilde{f}((x \rightarrow y) \rightarrow (x \rightarrow (x \rightarrow z)))$$

so from (3.4) that

$$\begin{aligned} \tilde{f}(x \rightarrow (x \rightarrow z)) &\supseteq \tilde{f}(x \rightarrow y) \cap \tilde{f}((x \rightarrow y) \rightarrow (x \rightarrow (x \rightarrow z))) \\ &\supseteq \tilde{f}(x \rightarrow y) \cap \tilde{f}(x \rightarrow (y \rightarrow z)). \end{aligned}$$

Since

$$x \rightarrow (x \rightarrow z) \leq x \rightarrow (((x \rightarrow z) \rightarrow z) \rightarrow z) = ((x \rightarrow z) \rightarrow z) \rightarrow (x \rightarrow z),$$

we have

$$\begin{aligned} \tilde{f}(x \rightarrow z) &\supseteq \tilde{f}(((x \rightarrow z) \rightarrow z) \rightarrow (x \rightarrow z)) \\ &\supseteq \tilde{f}(x \rightarrow (x \rightarrow z)) \\ &\supseteq \tilde{f}(x \rightarrow y) \cap \tilde{f}(x \rightarrow (y \rightarrow z)) \end{aligned}$$

by using (3.2) and (3.10). This completes the proof. □

**Proposition 3.17.** *Every Boolean int-soft filter  $(\tilde{f}, L)$  of  $L$  satisfies the following inclusion:*

$$(\forall x, y, z \in L) \left( \tilde{f}(x \rightarrow z) \supseteq \tilde{f}(x \rightarrow (y \rightarrow z)) \cap \tilde{f}(x \rightarrow y) \right).$$

*Proof.* Note that  $x \rightarrow (y \rightarrow z) = y \rightarrow (x \rightarrow z) \leq (x \rightarrow y) \rightarrow (x \rightarrow (x \rightarrow z))$  and

$$x \rightarrow (x \rightarrow z) \leq x \rightarrow (((x \rightarrow z) \rightarrow z) \rightarrow z) = ((x \rightarrow z) \rightarrow z) \rightarrow (x \rightarrow z)$$

for all  $x, y, z \in L$ . It follows from (3.2), (3.4), and Propositions 3.13 and 3.14 that

$$\begin{aligned} \tilde{f}(x \rightarrow z) &\supseteq \tilde{f}(((x \rightarrow z) \rightarrow z) \rightarrow (x \rightarrow z)) \\ &\supseteq \tilde{f}(x \rightarrow (x \rightarrow z)) \\ &\supseteq \tilde{f}(x \rightarrow y) \cap \tilde{f}((x \rightarrow y) \rightarrow (x \rightarrow (x \rightarrow z))) \\ &\supseteq \tilde{f}(x \rightarrow y) \cap \tilde{f}(x \rightarrow (y \rightarrow z)). \end{aligned}$$

This completes the proof. □

Combining Propositions 3.13, 3.14 and 3.15, we have a characterization of a Boolean int-soft filter.

**Theorem 3.18.** *Let  $(\tilde{f}, L)$  be an int-soft filter of  $L$ . Then the following assertions are equivalent:*

Int-soft filters of MTL-algebras

- (1)  $(\tilde{f}, L)$  is Boolean.
- (2)  $(\tilde{f}, L)$  satisfies the condition (3.10).
- (3)  $(\tilde{f}, L)$  satisfies the condition (3.11).

**Proposition 3.19.** *Every Boolean int-soft filter  $(\tilde{f}, L)$  of  $L$  satisfies:*

$$(\forall x, y \in L) \left( \tilde{f}(x \rightarrow y) \subseteq \tilde{f}(((y \rightarrow x) \rightarrow x) \rightarrow y) \right). \tag{3.13}$$

*Proof.* Let  $(\tilde{f}, L)$  be a Boolean int-soft filter of  $L$ . Since  $y \leq ((y \rightarrow x) \rightarrow x) \rightarrow y$ , we have

$$(((y \rightarrow x) \rightarrow x) \rightarrow y) \rightarrow x \leq y \rightarrow x \tag{3.14}$$

by (a6). Using (a4), (a5), (a6) and (3.14), we get

$$\begin{aligned} x \rightarrow y &\leq ((y \rightarrow x) \rightarrow x) \rightarrow ((y \rightarrow x) \rightarrow y) \\ &= (y \rightarrow x) \rightarrow (((y \rightarrow x) \rightarrow x) \rightarrow y) \\ &\leq (((y \rightarrow x) \rightarrow x) \rightarrow y) \rightarrow x \rightarrow (((y \rightarrow x) \rightarrow x) \rightarrow y) \end{aligned}$$

and so

$$\begin{aligned} \tilde{f}(((y \rightarrow x) \rightarrow x) \rightarrow y) &\supseteq \tilde{f}((((y \rightarrow x) \rightarrow x) \rightarrow y) \rightarrow x \rightarrow (((y \rightarrow x) \rightarrow x) \rightarrow y)) \\ &\supseteq \tilde{f}(x \rightarrow y) \end{aligned}$$

by Theorem 3.18(3) and (3.2). □

**Theorem 3.20.** (Extension Property for Boolean int-soft filter) *Let  $(\tilde{f}, L)$  and  $(\tilde{g}, L)$  be two int-soft filters of  $L$  such that  $\tilde{f}(1) = \tilde{g}(1)$  and  $\tilde{f}(x) \subseteq \tilde{g}(x)$  for all  $x \in L$ . If  $(\tilde{f}, L)$  is Boolean, then so is  $(\tilde{g}, L)$ .*

*Proof.* Assume that  $(\tilde{f}, L)$  is a Boolean int-soft filter of  $L$ . Then  $\tilde{f}(x \vee x^*) = \tilde{f}(1)$  for all  $x \in L$ . It follows from the hypothesis that

$$\tilde{g}(x \vee x^*) \supseteq \tilde{f}(x \vee x^*) = \tilde{f}(1) = \tilde{g}(1). \tag{3.15}$$

Combining (3.15) and (3.3), we have  $\tilde{g}(x \vee x^*) = \tilde{g}(1)$  for all  $x \in L$ . Hence  $(\tilde{g}, L)$  is a Boolean int-soft filter of  $L$ . □

For any soft set  $(\tilde{f}, L)$  over  $U$ , let  $(\tilde{g}, L)$  be a soft set over  $U$  in which

$$\tilde{g}(x) := \bigcup \left\{ \bigcap_{k=1}^n \tilde{f}(a_k) \mid \begin{array}{l} a_n \rightarrow (\cdots \rightarrow (a_2 \rightarrow (a_1 \rightarrow x)) \cdots) = 1, \\ a_1, a_2, \dots, a_n \in L \end{array} \right\} \tag{3.16}$$

Young Bae Jun, Seok Zun, Eun Hwan Roh and Sun Shin Ahn

for all  $x \in L$ . Let  $a, b, x \in L$  be such that  $a \leq b \rightarrow x$ . Take  $a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_m \in L$  such that

$$\begin{aligned} a_n &\rightarrow (\dots \rightarrow (a_2 \rightarrow (a_1 \rightarrow a)) \dots) = 1, \\ b_m &\rightarrow (\dots \rightarrow (b_2 \rightarrow (b_1 \rightarrow b)) \dots) = 1, \\ \tilde{g}(a) &= \bigcap_{k=1}^n \tilde{f}(a_k), \\ \tilde{g}(b) &= \bigcap_{j=1}^m \tilde{f}(b_j). \end{aligned}$$

Then

$$b_m \rightarrow (\dots \rightarrow (b_1 \rightarrow (a_n \rightarrow (\dots \rightarrow (a_2 \rightarrow (a_1 \rightarrow x)) \dots))) \dots) = 1,$$

and so

$$\begin{aligned} \tilde{g}(x) &\supseteq \tilde{f}(a_1) \cap \tilde{f}(a_2) \cap \dots \cap \tilde{f}(a_n) \cap \tilde{f}(b_1) \cap \tilde{f}(b_2) \cap \dots \cap \tilde{f}(b_m) \\ &= \left( \bigcap_{k=1}^n \tilde{f}(a_k) \right) \cap \left( \bigcap_{j=1}^m \tilde{f}(b_j) \right) \\ &= \tilde{g}(a) \cap \tilde{g}(b). \end{aligned}$$

Hence  $(\tilde{g}, L)$  is an int-soft filter of  $L$  by Theorem 3.4. Since  $x \rightarrow x = 1$  for all  $x \in L$ , we have  $\tilde{f}(x) \subseteq \tilde{g}(x)$  for all  $x \in L$ . Thus  $(\tilde{g}, L)$  contains  $(\tilde{f}, L)$ . Let  $(\tilde{h}, L)$  be an int-soft filter of  $L$  that contains  $(\tilde{f}, L)$ . Then

$$\begin{aligned} \tilde{g}(x) &= \bigcup \left\{ \bigcap_{k=1}^n \tilde{f}(a_k) \mid \begin{array}{l} a_n \rightarrow (\dots \rightarrow (a_2 \rightarrow (a_1 \rightarrow x)) \dots) = 1, \\ a_1, a_2, \dots, a_n \in L \end{array} \right\} \\ &\subseteq \bigcup \left\{ \bigcap_{k=1}^n \tilde{h}(a_k) \mid \begin{array}{l} a_n \rightarrow (\dots \rightarrow (a_2 \rightarrow (a_1 \rightarrow x)) \dots) = 1, \\ a_1, a_2, \dots, a_n \in L \end{array} \right\} \\ &\subseteq \bigcup \tilde{h}(x) = \tilde{h}(x) \end{aligned}$$

by Corollary 3.5, that is,  $(\tilde{g}, L)$  is contained in  $(\tilde{h}, L)$ .

We summarize this as follows:

**Theorem 3.21.** For any soft set  $(\tilde{f}, L)$  over  $U$ , the soft set  $(\tilde{g}, L)$  over  $U$  in which

$$\tilde{g}(x) := \bigcup \left\{ \bigcap_{k=1}^n \tilde{f}(a_k) \mid \begin{array}{l} a_n \rightarrow (\dots \rightarrow (a_2 \rightarrow (a_1 \rightarrow x)) \dots) = 1, \\ a_1, a_2, \dots, a_n \in L \end{array} \right\}$$

for all  $x \in L$  is the least int-soft filter of  $L$  that contains  $(\tilde{f}, L)$ .

## Int-soft filters of MTL-algebras

## CONCLUSION

Based on the soft set theory, we have introduced the notion of (Boolean) int-soft filters in MTL-algebras, and have investigated several properties. We have discussed characterizations of (Boolean) int-soft filters, and have provided a condition for an int-soft filter to be Boolean. We have established the extension property for a Boolean int-soft filter. We have also constructed the least int-soft filter containing a given soft set. Future research will focus on applying the notions and contents to other types of filters in related algebraic structures, and on studying it again by using Boolean algebra instead of  $\mathcal{P}(U)$ .

## REFERENCES

- [1] R. A. Borzooei, S. Khosravi Shoar and R. Ameri, Some types of filters in MTL-algebras, *Fuzzy Sets and Systems* 187 (2012), 92–102.
- [2] N. Çağman, F. Çitak and S. Enginoğlu, Soft set theory and uni-int decision making, *Eur. J. Oper. Res.* 207 (2010) 848–855.
- [3] F. Esteva and L. Godo, Monoidal  $t$ -norm based logic: towards a logic for left-continuous  $t$ -norms, *Fuzzy Sets and Systems* 124 (2001), 271–288.
- [4] P. Hájek, *Metamathematics of Fuzzy Logic*, Kluwer Academic Press, Dordrecht, 1998.
- [5] D. Molodtsov, Soft set theory - First results, *Comput. Math. Appl.* 37 (1999) 19–31.
- [6] W. Morton and C. J. van Alten, Modal MTL-algebras, *Fuzzy Sets and Systems* 222 (2013), 58–77.
- [7] E. Turunen, BL-algebras of basic fuzzy logic, *Mathware & Soft Computing* 6 (1999), 49–61.
- [8] T. Vetterlein, MTL-algebras arising from partially ordered groups, *Fuzzy Sets and Systems* 161 (2010), 433–443.

# Convergence Analysis of New Iterative Approximating Schemes with Errors for Total Asymptotically Nonexpansive Mappings in Hyperbolic Spaces

Ting-jian Xiong <sup>a</sup> and Heng-you Lan <sup>a, b \*</sup>

<sup>a</sup> Department of Mathematics, Sichuan University of Science & Engineering,  
Zigong, Sichuan 643000, PR China

<sup>b</sup> Key Laboratory of Higher Education of Sichuan Province for Enterprise Informationalization and  
Internet of Things, Zigong, Sichuan 643000, PR China

**Abstract.** The purpose of this paper is to introduce the concept of total asymptotically nonexpansive mappings and to prove some  $\Delta$ -convergence theorems of iteration processes with errors to approximating a common fixed point for four total asymptotically nonexpansive mappings in hyperbolic spaces. The results presented in the paper extend and improve some recent results announced in the current literature.

**Key Words and Phrases.** New iterative approximations with errors, asymptotically nonexpansive mapping, total asymptotically nonexpansive mapping, common fixed point, convergence analysis.

**AMS Subject Classification.** 47H09, 47H10, 54E70.

## 1 Introduction and preliminaries

Most of the problems in various disciplines of science are nonlinear in nature, whereas fixed point theory proposed in the setting of normed linear spaces or Banach spaces majorly depends on the linear structure of the underlying spaces. A nonlinear framework for fixed point theory is a metric space embedded with a 'convex structure'. The class of hyperbolic spaces, nonlinear in nature, is a general abstract theoretic setting with rich geometrical structure for metric fixed point theory. The study of hyperbolic spaces has been largely motivated and dominated by questions about hyperbolic groups, one of the main objects of study in geometric group theory.

Throughout this paper, we work in the setting of hyperbolic spaces due to Kohlenbach [1], defined below, which is more restrictive than the hyperbolic type introduced in [2] and more general than the concept of hyperbolic space in [3].

A hyperbolic spaces is a metric space  $(X, d)$  together with a mapping  $W : X^2 \times [0, 1] \rightarrow X$  satisfying

- (i)  $d(u, W(x, y, \alpha)) \leq \alpha d(u, x) + (1 - \alpha)d(u, y)$ ;
- (ii)  $d(W(x, y, \alpha), W(x, y, \beta)) = |\alpha - \beta|d(x, y)$ ;
- (iii)  $W(x, y, \alpha) = W(y, x, (1 - \alpha))$ ;
- (iv)  $d(W(x, z, \alpha), W(y, w, \alpha)) \leq \alpha d(x, y) + (1 - \alpha)d(z, w)$

for all  $u, x, y, z, w \in X$  and  $\alpha, \beta \in [0, 1]$  (see also [4]). A nonempty subset  $K$  of a hyperbolic space  $X$  is convex if  $W(x, y, \alpha) \in K$  for all  $x, y \in K$  and  $\alpha \in [0, 1]$ . The class of hyperbolic spaces contains normed spaces and convex subsets thereof, the Hilbert open unit ball equipped with the hyperbolic metric [5], Hadamard manifolds as well as CAT(0) spaces in the sense of Gromov (see [6]).

---

\*The corresponding author: hengyoulan@163.com (H.Y. Lan)

A hyperbolic space is uniformly convex [7, 8] if for any  $r > 0$  and  $\epsilon \in (0, 2]$  there exists  $\delta \in (0, 1]$  such that for all  $u, x, y \in X$ , we have

$$d(W(x, y, \frac{1}{2}), u) \leq (1 - \delta)r,$$

provided  $d(x, u) \leq r$ ,  $d(y, u) \leq r$  and  $d(x, y) \geq \epsilon r$ .

A map  $\eta : (0, +\infty) \times (0, 2] \rightarrow (0, 1]$ , which provides such  $\delta = \eta(r, \epsilon)$  for given  $r > 0$  and  $\epsilon \in (0, 2]$ , is known as a modulus of uniform convexity of  $X$ . We call  $\eta$  monotone if it decreases with  $r$  (for fixed  $\epsilon$ ), i.e.,  $\forall \epsilon > 0, \forall r_2 \geq r_1 > 0 (\eta(r_2, \epsilon) \leq \eta(r_1, \epsilon))$ .

In the sequel, let  $(X, d)$  be a metric space, and let  $K$  be a nonempty subset of  $X$ . We shall denote the fixed point set of a mapping  $T$  by  $F(T) = \{x \in K : Tx = x\}$ .

A mapping  $T : K \rightarrow K$  is said to be nonexpansive if

$$d(Tx, Ty) \leq d(x, y), \quad \forall x, y \in K.$$

A mapping  $T : K \rightarrow K$  is said to be asymptotically nonexpansive if there exists a sequence  $\{k_n\} \subset [0, +\infty)$  with  $k_n \rightarrow 0$  such that

$$d(T^n x, T^n y) \leq (1 + k_n)d(x, y), \quad \forall n \geq 1, x, y \in K.$$

A mapping  $T : K \rightarrow K$  is said to be uniformly  $L$ -Lipschitzian if there exists a constant  $L > 0$  such that

$$d(T^n x, T^n y) \leq Ld(x, y), \quad \forall n \geq 1, x, y \in K.$$

**Definition 1.1** A mapping  $T : K \rightarrow K$  is said to be  $(\{\mu_n\}, \{\xi_n\}, \rho)$ -total asymptotically nonexpansive if there exist nonnegative sequences  $\{\mu_n\}, \{\xi_n\}$  with  $\mu_n \rightarrow 0, \xi_n \rightarrow 0$  and a strictly increasing continuous function  $\rho : [0, +\infty) \rightarrow [0, +\infty)$  with  $\rho(0) = 0$  such that

$$d(T^n x, T^n y) \leq d(x, y) + \mu_n \rho(d(x, y)) + \xi_n, \quad \forall n \geq 1, x, y \in K.$$

**Remark 1.1** From the definitions, it is to know that each nonexpansive mapping is an asymptotically nonexpansive mapping with a sequence  $\{k_n = 0\}$ , and each asymptotically nonexpansive mapping is a  $(\{\mu_n\}, \{\xi_n\}, \rho)$ -total asymptotically nonexpansive mapping with  $\xi_n = 0$ , and  $\rho(t) = t, t \geq 0$ .

The existence of fixed points of various nonlinear mappings has relevant applications in many branches of nonlinear analysis and topology. On the other hand, there are certain situations where it is difficult to derive conditions for the existence of fixed points for certain types of nonlinear mappings. It is worth to mention that fixed point theory for nonexpansive mappings, a limit case of a contraction mapping when the Lipschitz constant is allowed to be 1, requires tools far beyond metric fixed point theory. Iteration schemas are the only main tool for analysis of generalized nonexpansive mappings. Fixed point theory has a computational flavor as one can define effective iteration schemas for the computation of fixed points of various nonlinear mappings. The problem of finding a common fixed point of some nonlinear mappings acting on a nonempty convex domain often arises in applied mathematics.

On the other hand, Zhao et al. [9] introduced a mixed type iteration for total asymptotically nonexpansive mappings in hyperbolic spaces, and prove some  $\Delta$ -convergence theorems for the iteration process approximating to a common fixed point; Zhao et al. [10] consider convergence theorems for total asymptotically nonexpansive mappings in hyperbolic spaces. Furthermore, Fukhar-ud-din and Kalsoom [11] extended iterative process with errors to asymptotically nonexpansive mappings in hyperbolic spaces, and obtained some convergence results.

Motivated and inspired by the above works, the purpose of this paper is to introduce the concepts of total asymptotically nonexpansive mappings and to prove some  $\Delta$ -convergence theorems of iteration process with errors for approximating a common fixed point of four total asymptotically nonexpansive mappings in hyperbolic spaces. The results presented in the paper extend and improve some recent results given in [9-25].



## 2 Preliminaries

In order to define the concept of  $\Delta$ -convergence in the general setup of hyperbolic spaces, we first collect some basic concepts.

Let  $\{x_n\}$  be a bounded sequence in a hyperbolic space  $X$ . For  $x \in X$ , we define a continuous functional  $r(\cdot, \{x_n\}) : X \rightarrow [0, +\infty)$  by

$$r(x, \{x_n\}) = \limsup_{n \rightarrow \infty} d(x, x_n).$$

The asymptotic radius  $r(\{x_n\})$  of  $\{x_n\}$  is given by

$$r(\{x_n\}) = \inf\{r(x, \{x_n\}) : x \in X\}.$$

The asymptotic center  $A_K(\{x_n\})$  of a bounded sequence  $\{x_n\}$  with respect to  $K \subset X$  is the set

$$A_K(\{x_n\}) = \{x \in X : r(x, \{x_n\}) \leq r(y, \{x_n\}), \forall y \in K\}.$$

This is the set of minimizers of the functional  $r(\cdot, \{x_n\})$ . If the asymptotic center is taken with respect to  $X$ , then it is simply denoted by  $A(\{x_n\})$ . It is known that uniformly convex Banach spaces and CAT(0) spaces enjoy the property that 'bounded sequences have unique asymptotic centers with respect to closed convex subsets'. The following lemma is due to Leustean [26] and ensures that this property also holds in a complete uniformly convex hyperbolic space.

**Lemma 2.1** ([26]) Let  $(X, d, W)$  be a complete uniformly convex hyperbolic space with monotone modulus of uniform convexity. Then every bounded sequence  $\{x_n\}$  in  $X$  has a unique asymptotic center with respect to any nonempty closed convex subset  $K$  of  $X$ .

Recall that a sequence  $\{x_n\}$  in  $X$  is said to  $\Delta$ -converge to  $x \in X$  if  $x$  is the unique asymptotic center of  $\{u_n\}$  for every subsequence  $\{u_n\}$  of  $\{x_n\}$ . In this case, we write  $\Delta\text{-}\lim_{n \rightarrow \infty} x_n = x$  and call  $x$  the  $\Delta$ -limit of  $\{x_n\}$ .

**Lemma 2.2** ([27]) Let  $\{a_n\}, \{b_n\}$  and  $\{\omega_n\}$  be sequences of nonnegative real numbers satisfying

$$a_{n+1} \leq (1 + \omega_n)a_n + b_n, \quad \forall n \geq 1.$$

If  $\sum_{n=1}^{\infty} \omega_n < +\infty$  and  $\sum_{n=1}^{\infty} b_n < +\infty$ , then  $\lim_{n \rightarrow \infty} a_n$  exists. If there exists a subsequence  $\{a_{n_i}\} \subset \{a_n\}$  such that  $a_{n_i} \rightarrow 0$ , then  $\lim_{n \rightarrow \infty} a_n = 0$ .

**Lemma 2.3** ([17]) Let  $(X, d, W)$  be a uniformly convex hyperbolic space with monotone modulus of uniform convexity  $\eta$ . Let  $x \in X$  and  $\{\alpha_n\}$  be a sequence in  $[a, b]$  for some  $a, b \in (0, 1)$ . If  $\{x_n\}$  and  $\{y_n\}$  are sequences in  $X$  such that

$$\begin{aligned} \limsup_{n \rightarrow \infty} d(x_n, x) \leq c, \quad \limsup_{n \rightarrow \infty} d(y_n, x) \leq c, \\ \lim_{n \rightarrow \infty} d(W(x_n, y_n, \alpha_n), x) = c \end{aligned}$$

for some  $c \geq 0$ , then  $\lim_{n \rightarrow \infty} d(x_n, y_n) = 0$ .

**Lemma 2.4** [17] Let  $K$  be a nonempty closed convex subset of uniformly convex hyperbolic space and  $\{x_n\}$  be a bounded sequence in  $K$  such that  $A(\{x_n\}) = \{y\}$  and  $r(\{x_n\}) = \zeta$ . If  $\{y_m\}$  is another sequence in  $K$  such that  $\lim_{m \rightarrow \infty} r(y_m, \{x_n\}) = \zeta$ , then  $\lim_{m \rightarrow \infty} y_m = y$ .

## 3 Main results

In this section, we give our main results.

**Theorem 3.1** Let  $K$  be a nonempty closed convex subset of a complete uniformly convex hyperbolic space  $X$  with monotone modulus of uniform convexity  $\eta$ . Let  $T_i : K \rightarrow K, i = 1, 2$ , be a uniformly  $L_i$ -Lipschitzian and  $(\{\mu_n^i\}, \{\xi_n^i\}, \rho^i)$ -total asymptotically nonexpansive mapping with  $\{\mu_n^i\}$  and  $\{\xi_n^i\}$  satisfying  $\lim_{n \rightarrow \infty} \mu_n^i = 0, \lim_{n \rightarrow \infty} \xi_n^i = 0$  and a strictly increasing continuous function  $\rho^i : [0, +\infty) \rightarrow [0, +\infty)$  with  $\rho^i(0) = 0, i = 1, 2$ , let  $S_i : K \rightarrow K, i = 1, 2$ , be a uniformly  $\hat{L}_i$ -Lipschitzian and  $(\{\hat{\mu}_n^i\}, \{\hat{\xi}_n^i\}, \hat{\rho}^i)$ -total asymptotically nonexpansive mapping with  $\{\hat{\mu}_n^i\}$

and  $\{\hat{\xi}_n^i\}$  satisfying  $\lim_{n \rightarrow \infty} \hat{\mu}_n^i = 0$ ,  $\lim_{n \rightarrow \infty} \hat{\xi}_n^i = 0$  and a strictly increasing continuous function  $\hat{\rho}^i : [0, +\infty) \rightarrow [0, +\infty)$  with  $\hat{\rho}^i(0) = 0, i = 1, 2$ . Assume that  $\mathcal{F} := \bigcap_{i=1}^2 (F(T_i) \cap F(S_i)) \neq \emptyset$ , and for arbitrarily chosen  $x_1 \in K$ , a new iterative approximating scheme  $\{x_n\}$  with errors is defined as follows:

$$\begin{aligned} x_{n+1} &= W(S_1^n x_n, W(T_1^n y_n, u_n, \theta_{n_1}), \alpha_n), \\ y_n &= W(S_2^n x_n, W(T_2^n x_n, v_n, \theta_{n_2}), \beta_n), \end{aligned} \tag{3.1}$$

where  $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}, \{\delta_n\}, \{\zeta_n\}, \{\lambda_n\}$  are sequences in  $[0, 1]$  and  $\{u_n\}, \{v_n\}$  are bounded sequences in  $K$  and  $\theta_{n_1} = 1 - \frac{\gamma_n}{1-\alpha_n} = \frac{\delta_n}{1-\alpha_n}$ ,  $\theta_{n_2} = 1 - \frac{\lambda_n}{1-\beta_n} = \frac{\zeta_n}{1-\beta_n}$ . Let  $\{\mu_n^i\}, \{\hat{\mu}_n^i\}, \rho^i, \{\hat{\rho}_n^i\}, \{\xi_n^i\}, \hat{\xi}_n^i, i = 1, 2, \{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}, \{\delta_n\}, \{\zeta_n\}$  and  $\{\lambda_n\}$  satisfy the following conditions:

- (i)  $\sum_{n=1}^{\infty} \mu_n^i < +\infty, \sum_{n=1}^{\infty} \hat{\mu}_n^i < +\infty, \sum_{n=1}^{\infty} \xi_n^i < +\infty, \sum_{n=1}^{\infty} \hat{\xi}_n^i < +\infty, \sum_{n=1}^{\infty} \gamma_n < +\infty, \sum_{n=1}^{\infty} \lambda_n < +\infty, i = 1, 2;$
- (ii) There exist constants  $a, b \in (0, 1)$  such that  $\{\alpha_n\} \subset [a, b], \{\beta_n\} \subset [a, b], \{\delta_n\} \subset [a, b], \{\zeta_n\} \subset [a, b]$  and  $\lim_{n \rightarrow \infty} \alpha_n = \alpha \in [a, b];$
- (iii) There exist a constant  $M^* > 0$  such that  $\rho^i(r) \leq M^*r$  and  $\hat{\rho}^i(r) \leq M^*r, r > 0, i = 1, 2;$
- (iv)  $d(x, y) \leq d(S_i x, y)$  for all  $x, y \in K$  and  $i = 1, 2.$

Then the iterative sequence  $\{x_n\}$  defined by (3.1)  $\Delta$ -converges to a common fixed point of  $\mathcal{F} := \bigcap_{i=1}^2 (F(T_i) \cap F(S_i)).$

*Proof.* Set  $L = \max\{L_i, \hat{L}_i, i = 1, 2\}, \mu_n = \max\{\mu_n^i, \hat{\mu}_n^i, i = 1, 2\},$  and  $\xi_n = \max\{\xi_n^i, \hat{\xi}_n^i, i = 1, 2\}, \rho = \max\{\rho^i, \hat{\rho}^i, i = 1, 2\}.$  By condition (i), we know that  $\sum_{n=1}^{\infty} \mu_n < +\infty, \sum_{n=1}^{\infty} \xi_n < +\infty.$  The proof of Theorem 3.1 is divided into three steps.

Step 1. We first prove that  $\lim_{n \rightarrow \infty} d(x_n, p)$  exists for each  $p \in \mathcal{F}.$

For any given  $p \in \mathcal{F},$  since  $T_i$  and  $S_i, i = 1, 2,$  are total asymptotically nonexpansive mappings, by condition (iii) and (3.1), we have

$$\begin{aligned} d(x_{n+1}, p) &= d(W(S_1^n x_n, W(T_1^n y_n, u_n, \theta_{n_1}), \alpha_n), p) \\ &\leq \alpha_n d(S_1^n x_n, p) + (1 - \alpha_n) d(W(T_1^n y_n, u_n, \theta_{n_1}), p) \\ &\leq \alpha_n d(S_1^n x_n, p) + (1 - \alpha_n) (\theta_{n_1} d(T_1^n y_n, p) + (1 - \theta_{n_1}) d(u_n, p)) \\ &\leq \alpha_n d(S_1^n x_n, p) + \delta_n d(T_1^n y_n, p) + \gamma_n d(u_n, p) \\ &\leq \alpha_n [d(x_n, p) + \mu_n \rho(d(x_n, p))] + \xi_n \\ &\quad + \delta_n [d(y_n, p) + \mu_n \rho(d(y_n, p))] + \xi_n + \gamma_n d(u_n, p) \\ &\leq \alpha_n [(1 + \mu_n M^*) d(x_n, p) + \xi_n] \\ &\quad + \delta_n [(1 + \mu_n M^*) d(y_n, p) + \xi_n] + \gamma_n d(u_n, p) \\ &\leq \alpha_n (1 + \mu_n M^*) d(x_n, p) + \delta_n (1 + \mu_n M^*) d(y_n, p) \\ &\quad + \gamma_n d(u_n, p) + (\alpha_n + \delta_n) \xi_n, \end{aligned} \tag{3.2}$$

where

$$\begin{aligned} d(y_n, p) &= d(W(S_2^n x_n, W(T_2^n x_n, v_n, \theta_{n_2}), \beta_n), p) \\ &\leq \beta_n d(S_2^n x_n, p) + (1 - \beta_n) d(W(T_2^n x_n, v_n, \theta_{n_2}), p) \\ &\leq \beta_n d(S_2^n x_n, p) + (1 - \beta_n) [\theta_{n_2} d(T_2^n x_n, p) + (1 - \theta_{n_2}) d(v_n, p)] \\ &\leq \beta_n d(S_2^n x_n, p) + \zeta_n d(T_2^n x_n, p) + \lambda_n d(v_n, p) \\ &\leq \beta_n [d(x_n, p) + \mu_n \rho(d(x_n, p))] + \xi_n \\ &\quad + \zeta_n [d(x_n, p) + \mu_n \rho(d(x_n, p))] + \xi_n + \lambda_n d(v_n, p) \\ &\leq (\beta_n + \zeta_n) [(1 + \mu_n M^*) d(x_n, p) + \xi_n] + \lambda_n d(v_n, p) \\ &= (\beta_n + \zeta_n) (1 + \mu_n M^*) d(x_n, p) + \lambda_n d(v_n, p) + (\beta_n + \zeta_n) \xi_n \\ &\leq (1 + \mu_n M^*) d(x_n, p) + \lambda_n d(v_n, p) + \xi_n. \end{aligned} \tag{3.3}$$

Substituting (3.3) into (3.2) and simplifying it, we have

$$\begin{aligned}
 d(x_{n+1}, p) &\leq \delta_n(1 + \mu_n M^*)[(1 + \mu_n M^*)d(x_n, p) + \lambda_n d(v_n, p) + \xi_n] \\
 &\quad + \alpha_n(1 + \mu_n M^*)d(x_n, p) + \gamma_n d(u_n, p) + (\alpha_n + \delta_n)\xi_n \\
 &= (\alpha_n + \delta_n + \alpha_n \mu_n M^* + 2\delta_n \mu_n M^* + \delta_n \mu_n^2 M^{*2})d(x_n, p) \\
 &\quad + \gamma_n d(u_n, p) + \lambda_n \delta_n(1 + \mu_n M^*)d(v_n, p) \\
 &\quad + [\alpha_n + \delta_n + \delta_n(1 + \mu_n M^*)]\xi_n \\
 &\leq [1 + \mu_n M^*(\alpha_n + 2\delta_n + \delta_n \mu_n M^*)]d(x_n, p) \\
 &\quad + \gamma_n d(u_n, p) + \lambda_n \delta_n(1 + \mu_n M^*)d(v_n, p) \\
 &\quad + [1 + \delta_n(1 + \mu_n M^*)]\xi_n \\
 &= (1 + \omega_n)d(x_n, p) + b_n,
 \end{aligned} \tag{3.4}$$

where  $\omega_n = \mu_n M^*(\alpha_n + 2\delta_n + \delta_n \mu_n M^*)$ ,  $b_n = \gamma_n d(u_n, p) + \lambda_n \delta_n(1 + \mu_n M^*)d(v_n, p) + [1 + \delta_n(1 + \mu_n M^*)]\xi_n$ . Since  $\sum_{n=1}^{\infty} \mu_n < +\infty$ ,  $\sum_{n=1}^{\infty} \xi_n < +\infty$  and condition (i),(ii), and  $\{u_n\}$ ,  $\{v_n\}$  are bounded sequences in  $K$ , it follows from Lemma 2.2 that  $\lim_{n \rightarrow \infty} d(x_n, p)$  exists for each  $p \in \mathcal{F}$ .

Step 2. We show that

$$\lim_{n \rightarrow \infty} d(x_n, T_i x_n) = 0, \quad \lim_{n \rightarrow \infty} d(x_n, S_i x_n) = 0, \quad i = 1, 2. \tag{3.5}$$

For each  $p \in \mathcal{F}$ , from the proof of Step 1, we know that  $\lim_{n \rightarrow \infty} d(x_n, p)$  exists. We may assume that  $\lim_{n \rightarrow \infty} d(x_n, p) = c \geq 0$ . If  $c = 0$ , then the conclusion is trivial. Next, we deal with the case  $c > 0$ . From (3.3), we have

$$d(y_n, p) \leq (1 + \mu_n M^*)d(x_n, p) + \lambda_n d(v_n, p) + \xi_n. \tag{3.6}$$

Taking lim sup on both sides in (3.6), we have

$$\limsup_{n \rightarrow \infty} d(y_n, p) \leq c. \tag{3.7}$$

In addition, since

$$d(T_1^n y_n, p) \leq d(y_n, p) + \mu_n \rho(d(y_n, p)) + \xi_n \leq (1 + \mu_n M^*)d(y_n, p) + \xi_n$$

and

$$d(S_1^n x_n, p) \leq (1 + \mu_n M^*)d(x_n, p) + \xi_n,$$

we have

$$\limsup_{n \rightarrow \infty} d(T_1^n y_n, p) \leq c \tag{3.8}$$

and

$$\limsup_{n \rightarrow \infty} d(S_1^n x_n, p) \leq c. \tag{3.9}$$

Also

$$\begin{aligned}
 &d(W(T_1^n y_n, u_n, \theta_{n_1}), p) \\
 &\leq \theta_{n_1} d(T_1^n y_n, p) + (1 - \theta_{n_1})d(u_n, p) \\
 &= \frac{\delta_n}{1 - \alpha_n} d(T_1^n y_n, p) + \frac{\gamma_n}{1 - \alpha_n} d(u_n, p) \\
 &\leq d(T_1^n y_n, p) + \frac{\gamma_n}{1 - b} d(u_n, p).
 \end{aligned} \tag{3.10}$$

Since  $\sum_{n=1}^{\infty} \gamma_n < +\infty$ ,  $\lim_{n \rightarrow \infty} \gamma_n = 0$ , by boundedness of  $\{u_n\}$  in  $K$  and (3.8), taking lim sup on both sides in (3.10), we have

$$\limsup_{n \rightarrow \infty} d(W(T_1^n y_n, u_n, \theta_{n_1}), p) \leq c. \tag{3.11}$$

By  $\lim_{n \rightarrow \infty} d(x_{n+1}, p) = c$ , it is easy to prove that

$$\lim_{n \rightarrow \infty} d(W(S_1^n x_n, W(T_1^n y_n, u_n, \theta_{n_1}), \alpha_n), p) = c. \tag{3.12}$$

It follows from (3.9), (3.11), (3.12) and Lemma 2.3 that

$$\lim_{n \rightarrow \infty} d(S_1^n x_n, W(T_1^n y_n, u_n, \theta_{n_1})) = 0. \tag{3.13}$$

Since

$$\begin{aligned} d(x_{n+1}, p) &= d(W(S_1^n x_n, W(T_1^n y_n, u_n, \theta_{n_1}), \alpha_n), p) \\ &\leq d(S_1^n x_n, p) + d(S_1^n x_n, x_{n+1}) \\ &\leq d(S_1^n x_n, p) + (1 - \alpha_n)d(S_1^n x_n, W(T_1^n y_n, u_n, \theta_{n_1})), \end{aligned}$$

with the help of (3.13), we have

$$\liminf_{n \rightarrow \infty} d(S_1^n x_n, p) \geq c.$$

Combined with (3.9), it yields that

$$\lim_{n \rightarrow \infty} d(S_1^n x_n, p) = c. \tag{3.14}$$

Since

$$\begin{aligned} d(x_{n+1}, p) &\leq \alpha_n d(S_1^n x_n, p) + \delta_n d(T_1^n y_n, p) + \gamma_n d(u_n, p) \\ &\leq \alpha_n d(S_1^n x_n, p) + \delta_n [d(y_n, p) + \mu_n \rho(d(y_n, p)) + \xi_n] \\ &\quad + \gamma_n d(u_n, p) \\ &\leq \alpha_n d(S_1^n x_n, p) + \delta_n (1 + \mu_n M^*) d(y_n, p) + \gamma_n d(u_n, p) + \delta_n \xi_n \\ &\leq \alpha_n d(S_1^n x_n, p) + (1 - \alpha_n)(1 + \mu_n M^*) d(y_n, p) \\ &\quad + \gamma_n d(u_n, p) + (1 - \alpha_n) \xi_n, \end{aligned}$$

we get

$$\begin{aligned} &\frac{d(x_{n+1}, p) - \alpha_n d(S_1^n x_n, p)}{1 - \alpha_n} \\ &\leq (1 + \mu_n M^*) d(y_n, p) + \frac{\gamma_n}{1 - \alpha_n} d(u_n, p) + \xi_n. \end{aligned}$$

By condition (ii), (3.12) and (3.14), we have

$$\liminf_{n \rightarrow \infty} d(y_n, p) \geq c.$$

Combined with (3.7), it yields that

$$\lim_{n \rightarrow \infty} d(y_n, p) = c. \tag{3.15}$$

By the same method and (3.15), we can also prove that

$$\lim_{n \rightarrow \infty} d(S_2^n x_n, W(T_2^n x_n, v_n, \theta_{n_2})) = 0. \tag{3.16}$$

It follows from virtue of condition (iv), (3.13), and (3.16) that

$$\lim_{n \rightarrow \infty} d(x_n, W(T_1^n y_n, u_n, \theta_{n_1})) \leq \lim_{n \rightarrow \infty} d(S_1^n x_n, W(T_1^n y_n, u_n, \theta_{n_1})) = 0, \tag{3.17}$$

and

$$\lim_{n \rightarrow \infty} d(x_n, W(T_2^n x_n, v_n, \theta_{n_2})) \leq \lim_{n \rightarrow \infty} d(S_2^n x_n, W(T_2^n x_n, v_n, \theta_{n_2})) = 0. \tag{3.18}$$

From (3.1) and (3.16), we have

$$\begin{aligned} d(y_n, S_2^n x_n) &= d(W(S_2^n x_n, W(T_2^n x_n, v_n, \theta_{n_2}), \beta_n), S_2^n x_n) \\ &\leq (1 - \beta_n)d(S_2^n x_n, W(T_2^n x_n, v_n, \theta_{n_2})) \rightarrow 0 \\ &\quad (\text{as } n \rightarrow \infty) \end{aligned} \tag{3.19}$$

and

$$d(x_n, y_n) = d(x_n, W(T_2^n x_n, v_n, \theta_{n_2})) + d(S_2^n x_n, W(T_2^n x_n, v_n, \theta_{n_2})) + d(S_2^n x_n, y_n).$$

It follows from (3.16), (3.18) and (3.19) that

$$\lim_{n \rightarrow \infty} d(x_n, y_n) = 0. \tag{3.20}$$

This together with (3.17) implies that

$$\begin{aligned} &d(x_n, W(T_1^n x_n, u_n, \theta_{n_1})) \\ &\leq d(x_n, W(T_1^n y_n, u_n, \theta_{n_1})) + d(W(T_1^n y_n, u_n, \theta_{n_1}), W(T_1^n x_n, u_n, \theta_{n_1})) \\ &\leq d(x_n, W(T_1^n y_n, u_n, \theta_{n_1})) + \theta_{n_1}d(T_1^n y_n, T_1^n x_n) \\ &\leq d(x_n, W(T_1^n y_n, u_n, \theta_{n_1})) + \frac{\delta_n}{1 - \alpha_n}Ld(y_n, x_n) \rightarrow 0 \quad (n \rightarrow \infty). \end{aligned} \tag{3.21}$$

On the other hand, from (3.13) and (3.20), we have

$$\begin{aligned} &d(S_1^n x_n, W(T_1^n x_n, u_n, \theta_{n_1})) \\ &\leq d(S_1^n x_n, W(T_1^n y_n, u_n, \theta_{n_1})) + d(W(T_1^n y_n, u_n, \theta_{n_1}), W(T_1^n x_n, u_n, \theta_{n_1})) \\ &\leq d(S_1^n x_n, W(T_1^n y_n, u_n, \theta_{n_1})) + \theta_{n_1}d(T_1^n y_n, T_1^n x_n) \\ &\leq d(S_1^n x_n, W(T_1^n y_n, u_n, \theta_{n_1})) + \frac{\delta_n}{1 - \alpha_n}Ld(y_n, x_n) \rightarrow 0 \quad (n \rightarrow \infty). \end{aligned} \tag{3.22}$$

From (3.21) and (3.22), we have that

$$\begin{aligned} d(S_1^n x_n, x_n) &\leq d(S_1^n x_n, W(T_1^n x_n, u_n, \theta_{n_1})) \\ &\quad + d(W(T_1^n x_n, u_n, \theta_{n_1}), x_n) \rightarrow 0 \quad (n \rightarrow \infty). \end{aligned} \tag{3.23}$$

In addition, since

$$\begin{aligned} d(x_{n+1}, x_n) &= d(W(S_1^n x_n, W(T_1^n y_n, u_n, \theta_{n_1}), \alpha_n), x_n) \\ &\leq \alpha_n d(S_1^n x_n, x_n) + (1 - \alpha_n)d(W(T_1^n y_n, u_n, \theta_{n_1}), x_n), \end{aligned}$$

it follows from (3.17) and (3.23) that

$$\lim_{n \rightarrow \infty} d(x_{n+1}, x_n) = 0. \tag{3.24}$$

Observe that

$$\begin{aligned} d(x_n, T_1^n x_n) &\leq d(x_n, W(T_1^n x_n, u_n, \theta_{n_1})) + d(W(T_1^n x_n, u_n, \theta_{n_1}), T_1^n x_n) \\ &\leq d(x_n, W(T_1^n x_n, u_n, \theta_{n_1})) + (1 - \theta_{n_1})d(T_1^n x_n, u_n) \\ &\leq d(x_n, W(T_1^n x_n, u_n, \theta_{n_1})) \\ &\quad + \frac{\gamma_n}{1 - \alpha_n}[d(T_1^n x_n, x_n) + d(x_n, p) + d(u_n, p)], \end{aligned}$$

then

$$\begin{aligned} d(x_n, T_1^n x_n) &\leq \frac{1 - \alpha_n}{1 - \alpha_n - \gamma_n}d(x_n, W(T_1^n x_n, u_n, \theta_{n_1})) \\ &\quad + \frac{\gamma_n}{1 - \alpha_n - \gamma_n}[d(x_n, p) + d(u_n, p)]. \end{aligned} \tag{3.25}$$

By boundedness of  $\{u_n\}$  in  $K$  and condition (i), (ii) and  $\lim_{n \rightarrow \infty} d(x_n, p)$  exists and (3.21), (3.25), we have

$$\lim_{n \rightarrow \infty} d(x_n, T_1^n x_n) = 0. \tag{3.26}$$

Similarly, we can also prove that

$$\lim_{n \rightarrow \infty} d(x_n, T_2^n x_n) = 0. \tag{3.27}$$

For all  $i = 1, 2$ , now we know

$$\begin{aligned} d(x_n, T_i x_n) &\leq d(x_n, x_{n+1}) + d(x_{n+1}, T_i^{n+1} x_{n+1}) \\ &\quad + d(T_i^{n+1} x_{n+1}, T_i^{n+1} x_n) + d(T_i^{n+1} x_n, T_i x_n) \\ &\leq (1 + L)d(x_n, x_{n+1}) + d(x_{n+1}, T_i^{n+1} x_{n+1}) + Ld(T_i^n x_n, x_n). \end{aligned}$$

It follows from (3.24), (3.26) and (3.27) that

$$\lim_{n \rightarrow \infty} d(x_n, T_i x_n) = 0, \quad i = 1, 2.$$

By virtue of condition (iv), i.e.,

$$d(S_1 x_n, W(T_1^n x_n, u_n, \theta_{n_1})) \leq d(S_1^n x_n, W(T_1^n x_n, u_n, \theta_{n_1})),$$

we have

$$\begin{aligned} d(x_n, S_1 x_n) &\leq d(x_n, W(T_1^n x_n, u_n, \theta_{n_1})) + d(S_1 x_n, W(T_1^n x_n, u_n, \theta_{n_1})) \\ &\leq d(x_n, W(T_1^n x_n, u_n, \theta_{n_1})) + d(S_1^n x_n, W(T_1^n x_n, u_n, \theta_{n_1})), \end{aligned}$$

from (3.21) and (3.22), which implies that

$$\lim_{n \rightarrow \infty} d(x_n, S_1 x_n) = 0.$$

By the same method, we can also prove that

$$\lim_{n \rightarrow \infty} d(x_n, S_2 x_n) = 0.$$

Step 3. We shall prove that the sequence  $\{x_n\}$   $\Delta$ -converges to a common fixed point of  $\mathcal{F} := \bigcap_{i=1}^2 (F(T_i) \cap F(S_i))$ .

In fact, for each  $p \in F$ ,  $\lim_{n \rightarrow \infty} d(x_n, p)$  exist. This implies that the sequence  $\{d(x_n, p)\}$  is bounded, so is the sequence  $\{x_n\}$ . Hence, by virtue of Lemma 2.1,  $\{x_n\}$  has a unique asymptotic center  $A_K(\{x_n\}) = \{x\}$ .

Let  $\{u_n\}$  be any subsequence of  $\{x_n\}$  with  $A_K(\{u_n\}) = \{u\}$ . It follows from (3.5) that

$$\lim_{n \rightarrow \infty} d(u_n, T_i u_n) = 0. \tag{3.28}$$

Next, we show that  $u \in F(T_i)$ , for all  $i = 1, 2$ . For this, we define a sequence  $\{z_n^i\}$  in  $K$  by  $z_n^i = T_i^m u$ , for all  $i = 1, 2$ . So we calculate

$$\begin{aligned} d(z_m^i, u_n) &\leq d(T_i^m u, T_i^m u_n) + d(T_i^m u_n, T_i^{m-1} u_n) + \dots + d(T_i u_n, u_n) \\ &\leq d(u, u_n) + \mu_m \rho(d(u, u_n)) + \xi_m + \sum_{k=1}^m d(T_i^k u_n, T_i^{k-1} u_n) \\ &\leq (1 + \mu_m M^*)d(u, u_n) + \xi_m + \sum_{k=1}^m d(T_i^k u_n, T_i^{k-1} u_n). \end{aligned} \tag{3.29}$$

Since  $T_i$  is uniformly  $L$ -Lipschitzian, it follows from (3.29) that

$$d(z_m^i, u_n) \leq (1 + \mu_m M^*)d(u, u_n) + \xi_m + mLd(T_i u_n, u_n).$$

Taking lim sup on both sides of the above estimate and using (3.28), we have

$$\begin{aligned} r(z_m^i, \{u_n\}) &= \limsup_{n \rightarrow \infty} d(z_m^i, u_n) \\ &\leq (1 + \mu_m M^*) \limsup_{n \rightarrow \infty} d(u, u_n) + \xi_m \\ &= (1 + \mu_m M^*)r(u, \{u_n\}) + \xi_m, \end{aligned}$$

and so

$$\limsup_{m \rightarrow \infty} r(z_m^i, \{u_n\}) \leq r(u, \{u_n\}).$$

Based on  $A_K(\{u_n\}) = \{u\}$  and the definition of asymptotic center  $A_K(\{u_n\})$  of a bounded sequence  $\{u_n\}$  with respect to  $K \subset X$ , we have

$$r(u, \{u_n\}) \leq r(y, \{u_n\}), \quad \forall y \in K.$$

This implies that

$$\liminf_{m \rightarrow \infty} r(z_m^i, \{u_n\}) \geq r(u, \{u_n\}).$$

Hence, we have

$$\lim_{m \rightarrow \infty} r(z_m^i, \{u_n\}) = r(u, \{u_n\}).$$

It follows from Lemma 2.4 that  $\lim_{m \rightarrow \infty} z_m^i = u$ , namely,  $\lim_{m \rightarrow \infty} T_i^m u = u$ . As  $T_i$  is uniformly continuous, so that  $T_i u = T_i(\lim_{m \rightarrow \infty} T_i^m u) = \lim_{m \rightarrow \infty} T_i^{m+1} u = u$ . That is,  $u \in F(T_i)$ . Similarly, we also can show that  $u \in F(S_i)$ , for all  $i = 1, 2$ . Hence,  $u$  is the common fixed point of  $T_i$  and  $S_i$ , for all  $i = 1, 2$ . And we want to show  $x = u$ , suppose  $x \neq u$ , by the uniqueness of asymptotic centers,

$$\begin{aligned} \limsup_{n \rightarrow \infty} d(u_n, u) &< \limsup_{n \rightarrow \infty} d(u_n, x) \\ &\leq \limsup_{n \rightarrow \infty} d(x_n, x) \\ &< \limsup_{n \rightarrow \infty} d(x_n, u) \\ &= \limsup_{n \rightarrow \infty} d(u_n, u), \end{aligned}$$

a contradiction. Thus we have  $x = u$ . Since  $\{u_n\}$  is an arbitrary subsequence of  $\{x_n\}$ ,  $A(\{u_n\}) = \{x\}$  for all subsequence  $\{u_n\}$  of  $\{x_n\}$ . This proves that  $\{x_n\}$   $\Delta$ -converges to a common fixed point of  $\mathcal{F} := \bigcap_{i=1}^2 (F(T_i) \cap F(S_i))$ . This completes the proof.  $\square$

The following theorem can be obtained from Theorem 3.1 immediately.

**Theorem 3.2** Let  $K$  be a nonempty closed convex subset of a complete uniformly convex hyperbolic space  $X$  with monotone modulus of uniform convexity  $\eta$ . Let  $T_i : K \rightarrow K, i = 1, 2$ , be a uniformly  $L_i$ -Lipschitzian and  $(\{\mu_n^i\}, \{\xi_n^i\}, \rho^i)$ -total asymptotically nonexpansive mapping with  $\{\mu_n^i\}$  and  $\{\xi_n^i\}$  satisfying  $\lim_{n \rightarrow \infty} \mu_n^i = 0, \lim_{n \rightarrow \infty} \xi_n^i = 0$  and a strictly increasing continuous function  $\rho^i : [0, +\infty) \rightarrow [0, +\infty)$  with  $\rho^i(0) = 0, i = 1, 2$ , let  $S_i : K \rightarrow K, i = 1, 2$ , be a uniformly  $\hat{L}_i$ -Lipschitzian and asymptotically nonexpansive mapping with  $\{k_n^i\} \subset [0, +\infty)$  satisfying  $\lim_{n \rightarrow \infty} k_n^i = 0$ . Assume that  $\mathcal{F} := \bigcap_{i=1}^2 (F(T_i) \cap F(S_i)) \neq \phi$ , and for arbitrarily chosen  $x_1 \in K, \{x_n\}$  is defined as follows:

$$\begin{aligned} x_{n+1} &= W(S_1^n x_n, W(T_1^n y_n, u_n, \theta_{n_1}), \alpha_n), \\ y_n &= W(S_2^n x_n, W(T_2^n x_n, v_n, \theta_{n_2}), \beta_n), \end{aligned} \tag{3.30}$$

where  $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}, \{\delta_n\}, \{\zeta_n\}, \{\lambda_n\}$  are sequences in  $[0, 1]$  and  $\{u_n\}, \{v_n\}$  are bounded sequences in  $K$  and  $\theta_{n1} = 1 - \frac{\gamma_n}{1-\alpha_n} = \frac{\delta_n}{1-\alpha_n}, \theta_{n2} = 1 - \frac{\lambda_n}{1-\beta_n} = \frac{\zeta_n}{1-\beta_n}$ . Let  $\{\mu_n^i\}, \{\xi_n^i\}, \rho^i, \{k_n^i\}, i = 1, 2, \{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}, \{\delta_n\}, \{\zeta_n\}$  and  $\{\lambda_n\}$  satisfy the following conditions:

- (i)  $\sum_{n=1}^{\infty} \mu_n^i < +\infty, \sum_{n=1}^{\infty} \xi_n^i < +\infty, \sum_{n=1}^{\infty} k_n^i < +\infty, \sum_{n=1}^{\infty} \gamma_n < +\infty, \sum_{n=1}^{\infty} \lambda_n < +\infty, i = 1, 2;$
- (ii) There exist constants  $a, b \in (0, 1)$  such that  $\{\alpha_n\} \subset [a, b], \{\beta_n\} \subset [a, b], \{\delta_n\} \subset [a, b], \{\zeta_n\} \subset [a, b]$  and  $\lim_{n \rightarrow \infty} \alpha_n = \alpha \in [a, b];$
- (iii) There exist a constant  $M^* > 0$  such that  $\rho^i(r) \leq M^*r, r > 0, i = 1, 2;$
- (iv)  $d(x, y) \leq d(S_i x, y)$  for all  $x, y \in K$  and  $i = 1, 2.$

Then the sequence  $\{x_n\}$  defined by (3.30)  $\Delta$ -converges to a common fixed point of  $\mathcal{F} := \bigcap_{i=1}^2 (F(T_i) \cap F(S_i)).$

*Proof.* Take  $\hat{\rho}^i(t) = t, t \geq 0, \hat{\xi}_n^i = 0, \hat{\mu}_n^i = k_n^i, i = 1, 2,$  in Theorem 3.1. Since all the conditions in Theorem 3.1 are satisfied, it follows from Theorem 3.1 that the sequence  $\{x_n\}$   $\Delta$ -converges to a common fixed point of  $\mathcal{F} := \bigcap_{i=1}^2 (F(T_i) \cap F(S_i)).$  This completes the proof of Theorem 3.2.  $\square$

**Theorem 3.3** Let  $K$  be a nonempty closed convex subset of a complete uniformly convex hyperbolic space  $X$  with monotone modulus of uniform convexity  $\eta.$  Let  $T_i : K \rightarrow K, i = 1, 2,$  be a uniformly  $L_i$ -Lipschitzian and  $(\{\mu_n^i\}, \{\xi_n^i\}, \rho^i)$ -total asymptotically nonexpansive mapping with  $\{\mu_n^i\}$  and  $\{\xi_n^i\}$  satisfying  $\lim_{n \rightarrow \infty} \mu_n^i = 0, \lim_{n \rightarrow \infty} \xi_n^i = 0$  and a strictly increasing continuous function  $\rho^i : [0, +\infty) \rightarrow [0, +\infty)$  with  $\rho^i(0) = 0, i = 1, 2,$  let  $S_i : K \rightarrow K, i = 1, 2,$  be a uniformly  $\tilde{L}_i$ -Lipschitzian and asymptotically nonexpansive mapping with  $\{k_n^i\} \subset [0, +\infty)$  satisfying  $\lim_{n \rightarrow \infty} k_n^i = 0.$  Assume that  $\mathcal{F} := \bigcap_{i=1}^2 (F(T_i) \cap F(S_i)) \neq \phi,$  and for arbitrarily chosen  $x_1 \in K, \{x_n\}$  is defined as follows:

$$\begin{aligned} x_{n+1} &= W(S_1^n x_n, T_1^n y_n, \alpha_n), \\ y_n &= W(S_2^n x_n, T_2^n x_n, \beta_n), \end{aligned} \tag{3.31}$$

where  $\{\alpha_n\}$  and  $\{\beta_n\}$  are sequences in  $[0, 1].$  Let  $\{\mu_n^i\}, \{\xi_n^i\}, \rho^i, \{k_n^i\}, i = 1, 2, \{\alpha_n\}, \{\beta_n\}$  satisfy the following conditions:

- (i)  $\sum_{n=1}^{\infty} \mu_n^i < +\infty, \sum_{n=1}^{\infty} \xi_n^i < +\infty, \sum_{n=1}^{\infty} k_n^i < +\infty;$
- (ii) There exist constants  $a, b \in (0, 1)$  such that  $\{\alpha_n\} \subset [a, b], \{\beta_n\} \subset [a, b], \lim_{n \rightarrow \infty} \alpha_n = \alpha \in [a, b];$
- (iii) There exist a constant  $M^* > 0$  such that  $\rho^i(r) \leq M^*r, r > 0, i = 1, 2;$
- (iv)  $d(x, y) \leq d(S_i x, y)$  for all  $x, y \in K$  and  $i = 1, 2.$

Then the sequence  $\{x_n\}$  defined by (3.31)  $\Delta$ -converges to a common fixed point of  $\mathcal{F} := \bigcap_{i=1}^2 (F(T_i) \cap F(S_i)).$

*Proof.* Take  $\hat{\rho}^i(t) = t, t \geq 0, \hat{\xi}_n^i = 0, \hat{\mu}_n^i = k_n^i, i = 1, 2$  and  $\gamma_n \equiv \lambda_n \equiv 0$  in Theorem 3.1. Since all the conditions in Theorem 3.1 are satisfied, it follows from Theorem 3.1 that the sequence  $\{x_n\}$   $\Delta$ -converges to a common fixed point of  $\mathcal{F} := \bigcap_{i=1}^2 (F(T_i) \cap F(S_i)).$  This completes the proof.  $\square$

**Theorem 3.4** Let  $K$  be a nonempty closed convex subset of a complete uniformly convex hyperbolic space  $X$  with monotone modulus of uniform convexity  $\eta.$  Let  $T_i : K \rightarrow K, i = 1, 2,$  be a uniformly  $L_i$ -Lipschitzian and  $(\{\mu_n^i\}, \{\xi_n^i\}, \rho^i)$ -total asymptotically nonexpansive mapping with  $\{\mu_n^i\}$  and  $\{\xi_n^i\}$  satisfying  $\lim_{n \rightarrow \infty} \mu_n^i = 0, \lim_{n \rightarrow \infty} \xi_n^i = 0$  and a strictly increasing continuous function  $\rho^i : [0, +\infty) \rightarrow [0, +\infty)$  with  $\rho^i(0) = 0, i = 1, 2,$  Suppose that  $\mathcal{F} := \bigcap_{i=1}^2 F(T_i) \neq \phi,$  and for arbitrarily chosen  $x_1 \in K, \{x_n\}$  is defined as follows:

$$\begin{aligned} x_{n+1} &= W(x_n, T_1^n y_n, \alpha_n), \\ y_n &= W(x_n, T_2^n x_n, \beta_n), \end{aligned} \tag{3.32}$$

where  $\{\alpha_n\}$  and  $\{\beta_n\}$  are sequences in  $[0, 1].$  Let  $\{\mu_n^i\}, \{\xi_n^i\}, \rho^i, i = 1, 2, \{\alpha_n\}, \{\beta_n\}$  satisfy the following conditions:

- (i)  $\sum_{n=1}^{\infty} \mu_n^i < +\infty, \sum_{n=1}^{\infty} \xi_n^i < +\infty;$
- (ii) There exist constants  $a, b \in (0, 1)$  such that  $\{\alpha_n\} \subset [a, b], \{\beta_n\} \subset [a, b];$
- (iii) There exist a constant  $M^* > 0$  such that  $\rho^i(r) \leq M^*r, r > 0, i = 1, 2.$

Then the sequence  $\{x_n\}$  defined by (3.32)  $\Delta$ -converges to a common fixed point of  $\mathcal{F} := \bigcap_{i=1}^2 F(T_i).$



*Proof.* Take  $\gamma_n \equiv \lambda_n \equiv 0$  and  $S_i = I, i = 1, 2$  in Theorem 3.1. Since all the conditions in Theorem 3.1 are satisfied, it follows from Theorem 3.1 that the sequence  $\{x_n\}$   $\Delta$ -converges to a common fixed point of  $\mathcal{F} := \bigcap_{i=1}^2 F(T_i)$ .  $\square$

**Remark 3.1** The results of Theorems 3.3 and 3.4 improve the corresponding results in Theorem 2.1 of [9] and Theorem 7 of [10], respectively.

## Acknowledgements

This work was partially supported by the Sichuan Province Cultivation Fund Project of Academic and Technical Leaders, and the Open Research Fund of Key Laboratory of Higher Education of Sichuan Province for Enterprise Informationalization and Internet of Things (2013WZJ01).

## References

- [1] U. Kohlenbach, Some logical metatheorems with applications in functional analysis, *Trans. Amer. Math. Soc.* **357**(1) (2004), 89-128.
- [2] P.K.F. Kuhfittig, Common fixed points of nonexpansive mappings by iteration. *Pacific J. Math.* **97**(1) (1981), 137-139.
- [3] S. Reich and I. Shafrir, Nonexpansive iterations in hyperbolic spaces, *Nonlinear Anal.* **15** (1990), 537-558.
- [4] M. Abbas, M.A. Khamsi and A.R. Khan, Common fixed point and invariant approximation in hyperbolic ordered metric spaces, *Fixed Point Theory Appl.* **2011**, 2011: 25, 14 pp.
- [5] K. Goebel and S. Reich, *Uniform Convexity, Hyperbolic Geometry and Nonexpansive Mappings*, Dekker, New York, 1984.
- [6] N. Bridson and A. Haefliger, *Metric Spaces of Non-Positive Curvature*, Springer, Berlin, 1999.
- [7] L. Leustean, A quadratic rate of asymptotic regularity for CAT(0) spaces, *J. Math. Anal. Appl.* **325** (2007), 386-399.
- [8] T. Shimizu and W. Takahashi, Fixed points of multivalued mappings in certain convex metric spaces, *Topol. Methods. Nonlinear. Anal.* **8** (1996), 197-203.
- [9] L.C. Zhao, S.S. Chang and J.K. Kim, Mixed type iteration for total asymptotically nonexpansive mappings in hyperbolic spaces, *Fixed Point Theory Appl.* **2013**, 2013: 353, 11 pp.
- [10] L.C. Zhao, S.S. Chang and X.R. Wang, Convergence theorems for total asymptotically nonexpansive mappings in hyperbolic spaces, *J. Appl. Math.* **2013**, Art. ID 689765, 5 pp.
- [11] H. Fukhar-ud-din and A. Kalsoom, Fixed point approximation of asymptotically nonexpansive mappings in hyperbolic spaces, *Fixed Point Theory Appl.* **2014**, 2014: 64, 15 pp.
- [12] R.P. Agarwal, D. O'Regan and D.R. Sahu, Iterative construction of fixed points of nearly asymptotically nonexpansive mappings, *J. Nonlinear convex Anal.* **8**(1) (2007), 61-79.
- [13] S.S. Chang, Y.J. Cho and H.Y. Zhou, Demiclosed principle and weak convergence problems for asymptotically nonexpansive mappings, *J. Korean. Math. Soc.* **38**(6) (2001), 1245-1260.
- [14] H. Fukhar-ud-din, Strong convergence of an Ishikawa-type algorithm in CAT(0) spaces, *Fixed Point Theory Appl.* **2013**, 2013: 207, 11 pp.
- [15] H. Fukhar-ud-din and A.R. Khan, Approximating common fixed points of asymptotically nonexpansive maps in uniformly convex Banach spaces, *Comput. Math. Appl.* **53** (2007), 1349-1360.
- [16] F. Gu and Q. Fu, Strong convergence theorems for common fixed points of multistep iterations with errors in Banach spaces, *J. Inequal. Appl.* **2009**, Art. ID 819036, 12 pp.

- [17] A.R. Khan, H. Fukhar-ud-din and M.A.A. Kuan, An implicit algorithm for two finite families of non-expansive maps in hyperbolic spaces, *Fixed Point Theory Appl.* **2012**, 2012: 54, 12 pp.
- [18] A.R. Khan, M.A. Khamsi and H. Fukhar-ud-din, Strong convergence of a general iteration scheme in CAT(0) spaces, *Nonlinear Anal.* **74** (2011), 783-791.
- [19] M.O. Osilike and S.C. Aniagbosor, Weak and strong convergence theorems for fixed points of asymptotically nonexpansive mappings, *Math. Compt. Model.* **32** (2000), 1181-1191.
- [20] A. Sahin and M. Basarir, On the strong convergence of a modified  $S$ -iteration process for asymptotically quasi-nonexpansive mappings in CAT(0) spaces, *Fixed Point Theory Appl.* **2013**, 2013: 12, 10 pp.
- [21] J. Schu, Weak and strong convergence to fixed points of asymptotically nonexpansive mappings, *Bull. Aust. Math. Soc.* **43** (1991), 153-159.
- [22] J. Schu, Iterative construction of fixed points of asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* **158** (1991), 407-413.
- [23] H. Fukhar-ud-din and M.A. Khamsi, Approximating common fixed points in hyperbolic spaces, *Fixed Point Theory Appl.* **2014**, 2014: 113, 15 pp.
- [24] K.K. Tan, H.K. Xu, Fixed point iteration process for asymptotically nonexpansive mappings, *Proc. Amer. Math. Soc.* **122(3)** (1994), 733-739.
- [25] Y. Yao, Y.C. Liou, New iterative schemes for asymptotically quasi-nonexpansive mappings, *J. Inequal. Appl.* **2010**, Art. ID 934692, 9 pp.
- [26] L. Leustean, Nonexpansive iteration in uniformly convex  $W$ -hyperbolic spaces, *Nonlinear analysis and optimization I*, Nonlinear analysis, 193–210, Contemp. Math., 513, Amer. Math. Soc., Providence, RI, 2010.
- [27] K.K. Tan and H.K. Xu, Approximating fixed point of nonexpansive mapping by the Ishikawa iteration process, *J. Math. Anal. Appl.* **178** (1993), 301-308.

## SOME OSTROWSKI TYPE INTEGRAL INEQUALITIES FOR DOUBLE INTEGRAL ON TIME SCALES

WAJEEHA IRSHAD, MUHAMMAD IQBAL BHATTI, AND MUHAMMAD MUDDASSAR\*

ABSTRACT. Weighted montgomery identity on time scales for functions of two variables is established. Corresponding discrete and continuous versions of montgomery identities for functions of two variables are obtained. By using the obtained weighted montgomery identity on time scales, an Ostrowski type inequality for double integrals on time scales is pointed out as well.

### 1. INTRODUCTION AND PRELIMINARY RESULTS

The Ostrowski type inequality, which was originally presented by Ostrowski in [14], can be used to estimate the absolute deviation of a function from its integral mean. In [6], Bohner and Matthews derived the Montgomery identity on time scales and established the following Ostrowski inequality on time scales, which unifies and extends corresponding discrete [7], continuous [13] and other cases.

**Theorem 1.** *Let  $a, b, s, t \in \mathbb{T}$  with  $a < b$   $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function with the property that,  $M = \sup_{a < t < b} |f \Delta(t)| < \infty$ , induces*

$$\left| f(t) - \frac{1}{(b-a)} \int_a^b f(\sigma(s)) \Delta s \right| \leq \frac{M}{b-a} + (h_2(t, a) + h_2(t, b)) \quad (1.1)$$

*is the best possible in the sense that rightside cannot be replaced by a smaller quantity.*

*Where  $h(\cdot, \cdot)$  is defined by definition (6) below.*

### 2. TIME SCALE ESSENTIALS

During the past decades, with the development of the theory of differential and integral equations as well as difference equations, a lot of integral and difference inequalities have

*Date:* September 15, 2014.

*2010 Mathematics Subject Classification.* 26D10, 26D15, 39A12.

*Key words and phrases.* Ostrowski Inequality, Weighted Montgomery Identity, Time Scales.

been discovered e.g., and the references therein, which play an important role in the research of boundedness, global existence, stability of solutions of differential and integral equations as well as difference equations. On the other hand, Hilger initiated the theory of time scales as a theory capable to contain both difference and differential calculus in a consistent way. Since then many authors have expounded on various aspects of the theory of dynamic equations on time scales including various inequalities on time scales. The existence of a derivative at a point in a time scale depends on the type of the point itself, because time scale may not be connected. Points are classified according to two major operators, the forward jump operators and the backward jump operators.

**Definition 1.** For an arbitrary time scale  $T$ , the forward jump operator,  $\sigma : T \rightarrow T$ , is defined to be

$$\sigma(t) := \inf\{s \in T : s > t\}$$

If the set of all points in  $T$  that are larger than  $t$  is empty, then  $\inf \emptyset = \sup T$ .

If  $T$  has a maximum  $t$ , then  $\sigma(t) = t$ .

The backward jump operator,  $\rho : T \rightarrow T$ , is

$$\rho(t) := \sup\{s \in T : s < t\}$$

If the set of all points in  $T$  that are less than  $t$  is empty, then  $\sup \emptyset = \inf T$ .

If  $T$  has a minimum  $t$ , then  $\rho(t) = t$ .

**Definition 2.** Using these two operators for  $t \in T$ , the point can be classified in the following manner,  $t$  is right-scattered if  $\sigma(t) > t$ ,  $t$  is right-dense if  $\sigma(t) = t$ ,  $t$  is left-scattered if  $\rho(t) < t$ ,  $t$  is left-dense if  $\rho(t) = t$ ,  $t$  is isolated if it is left- and right-scattered:  $\rho(t) < t < \sigma(t)$  and  $t$  is dense if it is both left- and right-dense  $\rho(t) = t = \sigma(t)$  for  $t \in T$ .

**Definition 3.** The graininess function,  $\mu : T \rightarrow [0, \infty)$ , is defined to be

$$\mu(t) = \sigma(t) - t$$

The graininess function essentially describes the step size between two consecutive points in  $T$ . Oftentimes the differences in results obtained from discrete and continuous calculus stem from the different value of the graininess function evaluated at a given point  $t$ .

**Definition 4.** The derivative in time scale calculus, called the delta derivative, determines the rates of forward change over a time scale. For a function  $f : T \rightarrow R$  and  $t \in T^k$ , the

delta derivative of  $f$  at  $t$ ,  $f^\Delta(t)$ , is defined to be the number, when it exists, where for any given  $\epsilon > 0$ , there exists a neighborhood  $U$  of  $t$  such that

$$|[f(\sigma(t)) - f(s)] - f^\Delta(t)[\sigma(t) - s]| \leq \epsilon|\sigma(t) - s| \tag{2.2}$$

is true for all  $s \in U$ . Here,  $\mathbb{T}^k = \mathbb{T} \setminus \{m\}$  when  $\mathbb{T}$  has a left-scattered maximum  $m$ ; otherwise,  $\mathbb{T}^k = \mathbb{T}[1]$ . Since the delta derivative definition involves the forward jump operator; if the time scale has a left scattered maximum  $m$ , then one cannot jump past this point. Therefore, this point is removed from the set of points used to determine the delta derivative. However, if the time scale does not contain such a left-scattered maximum, then  $\mathbb{T}^k$  is equivalent to the time scale.

Take  $\mathbb{T} = \mathbb{R}$ , then  $\sigma(t) = t$ ,  $\mu(t) = 0$ ,  $f^\Delta = f'$  is the derivative used in standard calculus. If  $\mathbb{T} = \mathbb{Z}$ ,  $\sigma(t) = t + 1$ ,  $\mu(t) = 1$ ,  $f^\Delta = \Delta f$  is the forward difference operator used in difference equations.

**Theorem 2** (Properties of the Delta Derivative).  $f : \mathbb{T} \rightarrow \mathbb{R}$  be a function and  $t \in \mathbb{T}^k$  as defined above. For such a function, the following properties hold:

- (1) If  $f$  is delta differentiable at  $t$ , then  $f$  is continuous at  $t$ .
- (2) If  $t$  is right-scattered and  $f$  is continuous at  $t$ , then the delta derivative of  $f$ ,  $f^\Delta$ , is defined as follows

$$f^\Delta(t) = \frac{f(\sigma(t)) - f(t)}{\mu(t)}$$

- (3) If  $t$  is right-dense, then the delta derivative at  $t$  is as follows (if and only if the limit exists as a finite number)

$$f^\Delta(t) = \lim_{s \rightarrow t} \frac{f(t) - f(s)}{t - s}$$

- (4) If  $f$  is delta differentiable at  $t$ , then

$$f(\sigma(t)) = f(t) + \mu(t) f^\Delta(t)$$

- (5) If  $\mathbb{T} = \mathbb{R}$ , then the delta derivative is  $f'(t)$  from continuous calculus.
- (6) If  $\mathbb{T} = \mathbb{Z}$ , then the delta derivative is the forward difference,  $\Delta f$ , from discrete calculus.

**Definition 5.** If  $F^\Delta(t) = f(t)$  for all  $t \in \mathbb{T}^k$ , then  $F(t)$  is said to be anti-derivative of  $f(t)$  and  $f(t)$  is said to be delta integrable provided that  $f(t)$  is rd-continuous. The cauchy integral of  $f(t)$  is defined by  $\int_r^s f(t)\Delta(t) = F(s) - F(r)$ .

**Theorem 3.** Let  $f, g$  be rd-continuous,  $a, b, c \in \mathbb{T}$  and  $\alpha, \beta \in \mathbb{R}$ , then

- (1)  $\int_a^b (\alpha f(t) + \beta g(t)) \Delta t = \alpha \int_a^b f(t) \Delta t + \beta \int_a^b g(t) \Delta t$
- (2)  $\int_a^b f(t) \Delta t = - \int_b^a f(t) \Delta t$
- (3)  $\int_a^c f(t) \Delta t = \int_a^b f(t) \Delta t + \int_b^c f(t) \Delta t$
- (4)  $\int_a^b f(t) g^\Delta(t) \Delta t = f(b)g(b) - f(a)g(a) - \int_a^b f^\Delta(t)g(\sigma(t)) \Delta t$

**Theorem 4.** If  $f$  is  $\Delta$ -integrable on  $[a, b]$ , then so is  $|f|$ , and

$$\left| \int_a^b f(t) \Delta t \right| \leq \int_a^b |f(t)| \Delta t$$

Let  $\mathbb{T}_1, \mathbb{T}_2$  be two time scales. Let  $\sigma_i, \rho_i$  and  $\Delta_i$  be the forward jump operator, the backward jump operator and the delta differentiation, respectively on  $\mathbb{T}_i$ , for  $i = 1, 2$ . Let  $a, b \in \mathbb{T}_1, c, d \in \mathbb{T}_2$ , with  $a < b, c < d$ .  $[a, b]$  and  $[c, d]$  are the half-closed bounded intervals in  $\mathbb{T}_1$  and  $\mathbb{T}_2$  respectively, and a "rectangle" in  $\mathbb{T}_1 \times \mathbb{T}_2$  by

$$\mathbb{R} = [a, b] \times [c, d] = \{(t_1, t_2) : t_1 \in [a, b], t_2 \in [c, d]\}$$

Let  $f$  be a real valued function on  $\mathbb{T}_1 \times \mathbb{T}_2$ . This function  $f$  is said to be rd-continuous in  $t_2$  if  $a_1 \in \mathbb{T}_1$ , then function  $f$  is real valued function on  $\mathbb{T}_1 \times \mathbb{T}_2$ , this function  $f$  is said to be rd-continuous in  $t_2$  if  $a_1 \in \mathbb{T}_1$ , then  $f(a_1, t_2)$  is rd-continuous on  $\mathbb{T}_2$ .  $CC_{rd}$  denotes the set of functions  $f(a_1, t_2)$  on  $\mathbb{T}_1 \times \mathbb{T}_2$ , having the properties:

- (1)  $f$  is rd-continuous in  $t_1$  and  $t_2$ .
- (2) If  $(x_1, x_2) \in \mathbb{T}_1 \times \mathbb{T}_2$  with  $x_1$  right dense and  $x_2$  right dense, then  $f$  is continuous at  $(x_1, x_2)$ .

**Definition 6.** Let  $g_k, h_k : \mathbb{T}^2 \rightarrow \mathbb{R}, k \in \mathbb{N}_0$  be defined by  $g_0(t, s) = h_0(t, s) = 1$  for all  $s, t \in \mathbb{T}$  and then recursively by

$$g_{k+1}(t, s) = \int_s^t g_k(\sigma(\tau), s) \Delta\tau \forall s, t \in \mathbb{T} \tag{2.3}$$

$$h_{k+1}(t, s) = \int_s^t h_k(\sigma(\tau), s) \Delta\tau \forall s, t \in \mathbb{T} \tag{2.4}$$

### 3. MAIN RESULTS

**Lemma 1** (Weighted Montgomery Identity on Time Scales). Let  $g : [a, b] \rightarrow [0, \infty), G : [c, d] \rightarrow [0, \infty)$  be rd-continuous and positive and  $h : [a, b] \rightarrow \mathbb{R}, H : [c, d] \rightarrow \mathbb{R}$  be invertible and differentiable, such that  $g(t_1) = h^{\Delta_1}(t_1)$  on  $[a, b]$  and  $G(t_2) = H^{\Delta_2}(t_2)$ . Let  $a, b, s_1, t_1 \in \mathbb{T}_1, c, d, s_2, t_2 \in \mathbb{T}_2$  with  $a < b, c < d$ ,

$$A = h^{-1} \left( \left(1 - \frac{\alpha}{2}\right) h(a) + \frac{\alpha}{2} h(b) \right), B = h^{-1} \left( \left(1 - \frac{\alpha}{2}\right) h(b) + \frac{\alpha}{2} h(a) \right),$$

$$C = H^{-1} \left( \left(1 - \frac{\beta}{2}\right) G(c) + \frac{\beta}{2} H(d) \right), D = H^{-1} \left( \left(1 - \frac{\beta}{2}\right) G(d) + \frac{\beta}{2} H(c) \right)$$

and

$f : [a, b]_{\mathbb{T}_1} \times [c, d]_{\mathbb{T}_2} \rightarrow \mathbb{R}$  is  $\Delta_1 \Delta_2$  differentiable. Then for all  $s_1 \in [A, B], s_2 \in [C, D]$ ,

$0 \leq \alpha, \beta \leq 1$ , we have

$$\int_a^b \int_c^d W(t_1, t_2, s_1, s_2) \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \Delta_1 s_1 \Delta_2 s_2$$

$$= \left( \frac{\beta}{2} H(d) - \frac{\beta}{2} H(c) \right) \left\{ h(b) - \left( \left(1 - \frac{\alpha}{2}\right) h(b) + \frac{\alpha}{2} h(a) \right) \right\} (f(b, c) - f(a, d) + f(b, d))$$

$$+ \left( \frac{\alpha}{2} h(a) + \frac{\alpha}{2} h(b) \right) \left\{ (1 - \beta) H(c) + (\beta - 1) H(d) \right\} (f(a, t_2) + f(b, t_2))$$

$$+ \left( \frac{\beta}{2} H(c) + \frac{\beta}{2} H(d) \right) \left\{ (1 - \alpha) h(b) + (\alpha - 1) h(a) \right\} (f(t_1, c) - f(t_1, d))$$

$$+ ((1 - \alpha) h(b) + (\alpha - 1) h(a)) ((1 - \beta) H(d) + (\beta - 1) H(c)) f(t_1, t_2)$$

$$+ \left[ H(c) - \left( \left(1 - \frac{\beta}{2}\right) H(c) + \frac{\beta}{2} H(d) \right) \right] \int_a^b h'(s_1) f(\sigma(s_1), c) \Delta_1 s_1$$

$$- \left[ H(t_2) - \left( \left(1 - \frac{\beta}{2}\right) H(c) + \frac{\beta}{2} H(d) \right) \right] \int_a^b h'(s_1) f(\sigma(s_1), t_2) \Delta_1 s_1$$

$$- \int_c^d H'(s_2) ((1 - \alpha) h(a) + (\alpha - 1) h(b)) f(t_1, \sigma(s_2)) \Delta_2 s_2$$

$$- \int_c^d H'(s_2) \left\{ \left( h(a) - \left( \left(1 - \frac{\alpha}{2}\right) h(a) + \frac{\alpha}{2} h(b) \right) \right) \right\} f(a, \sigma(s_2)) \Delta_2 s_2$$

$$- \int_c^d H'(s_2) \left\{ \left( h(b) - \left( \left(1 - \frac{\alpha}{2}\right) h(b) + \frac{\alpha}{2} h(a) \right) \right) \right\} f(b, \sigma(s_2)) \Delta_2 s_2$$

$$+ \int_c^d H'(s_2) \left\{ \int_a^b h'(s_1) f(\sigma(s_1), \sigma(s_2)) \Delta_1 s_1 \right\} \Delta_2(t_2)$$

where

$$W_1(t_1, t_2, s_1, s_2) = \begin{cases} h(s_1) - \left( \left(1 - \frac{\alpha}{2}\right) h(a) + \frac{\alpha}{2} h(b) \right), & s_1 \in [a, t_1) \\ h(s_1) - \left( \frac{\alpha}{2} h(a) + \left(1 - \frac{\alpha}{2}\right) h(b) \right), & s_1 \in [t_1, b] \\ H(s_2) - \left( \left(1 - \frac{\beta}{2}\right) H(c) + \frac{\beta}{2} H(d) \right), & s_2 \in [c, t_2) \\ H(s_2) - \left( \frac{\beta}{2} H(c) + \left(1 - \frac{\beta}{2}\right) H(d) \right), & s_2 \in [t_2, d] \end{cases}$$

*Proof.* Let's start with

$$\int_a^b \int_c^d W(t_1, t_2, s_1, s_2) \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \Delta_1 s_1 \Delta_2 s_2$$

$$\begin{aligned}
 &= \int_a^{t_1} \int_c^{t_2} \left\{ h(s_1) - \left( \left( 1 - \frac{\alpha}{2} \right) h(a) + \frac{\alpha}{2} h(b) \right) \right\} \left\{ H(s_2) - \left( \left( 1 - \frac{\beta}{2} \right) H(c) + \frac{\beta}{2} H(d) \right) \right\} \\
 &\quad \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \Delta_1 s_1 \Delta_2 s_2 + \int_a^{t_1} \int_{t_2}^d \left\{ h(s_1) - \left( \left( 1 - \frac{\alpha}{2} \right) h(a) + \frac{\alpha}{2} h(b) \right) \right\} \\
 &\quad \left\{ H(s_2) - \left( \left( 1 - \frac{\beta}{2} \right) H(d) + \frac{\beta}{2} H(c) \right) \right\} \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \Delta_1 s_1 \Delta_2 s_2 \\
 &+ \int_{t_1}^b \int_c^{t_2} \left\{ h(s_1) - \left( \left( 1 - \frac{\alpha}{2} \right) h(b) + \frac{\alpha}{2} h(a) \right) \right\} \left\{ H(s_2) - \left( \left( 1 - \frac{\beta}{2} \right) H(c) + \frac{\beta}{2} H(d) \right) \right\} \\
 &\quad \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \Delta_1 s_1 \Delta_2 s_2 + \int_{t_1}^b \int_{t_2}^d \left\{ h(s_1) - \left( \left( 1 - \frac{\alpha}{2} \right) h(b) + \frac{\alpha}{2} h(a) \right) \right\} \left\{ H(s_2) - \right. \\
 &\quad \left. \left( \left( 1 - \frac{\beta}{2} \right) H(d) + \frac{\beta}{2} H(c) \right) \right\} \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \Delta_1 s_1 \Delta_2 s_2.
 \end{aligned}$$

Now

$$\begin{aligned}
 &= \int_a^{t_1} \left\{ h(s_1) - \left( \left( 1 - \frac{\alpha}{2} \right) h(a) + \frac{\alpha}{2} h(b) \right) \right\} \left[ \left\{ H(t_2) - \left( \left( 1 - \frac{\beta}{2} \right) H(c) + \frac{\beta}{2} H(d) \right) \right\} \right. \\
 &\quad \left. \frac{\partial f(s_1, t_2)}{\Delta_1 s_1} - \left\{ H(c) - \left( \left( 1 - \frac{\beta}{2} \right) H(c) + \frac{\beta}{2} H(d) \right) \right\} \frac{\partial f(s_1, c)}{\Delta_1 s_1} - \int_c^{t_2} H'(s_2) \frac{\partial f(s_1, \sigma(s_2))}{\Delta_1 s_1} \right] \Delta_2 s_2 \\
 &+ \int_a^{t_1} \left\{ h(s_1) - \left( \left( 1 - \frac{\alpha}{2} \right) h(a) + \frac{\alpha}{2} h(b) \right) \right\} \left[ \left\{ H(d) - \left( \left( 1 - \frac{\beta}{2} \right) H(d) + \frac{\beta}{2} H(c) \right) \right\} \right. \\
 &\quad \left. \frac{\partial f(s_1, d)}{\Delta_1 s_1} - \left\{ H(t_2) - \left( \left( 1 - \frac{\beta}{2} \right) H(d) + \frac{\beta}{2} H(c) \right) \right\} \frac{\partial f(s_1, t_2)}{\Delta_1 s_1} - \int_{t_2}^d H'(s_2) \frac{\partial f(s_1, \sigma(s_2))}{\Delta_1 s_1} \right] \Delta_2 s_2 \\
 &+ \int_{t_1}^b \left\{ h(s_1) - \left( \left( 1 - \frac{\alpha}{2} \right) h(b) + \frac{\alpha}{2} h(a) \right) \right\} \left[ \left\{ H(t_2) - \left( \left( 1 - \frac{\beta}{2} \right) H(c) + \frac{\beta}{2} H(d) \right) \right\} \right. \\
 &\quad \left. \frac{\partial f(s_1, t_2)}{\Delta_1 s_1} - \left\{ H(c) - \left( \left( 1 - \frac{\beta}{2} \right) H(c) + \frac{\beta}{2} H(d) \right) \right\} \frac{\partial f(s_1, c)}{\Delta_1 s_1} - \int_c^{t_2} H'(s_2) \frac{\partial f(s_1, \sigma(s_2))}{\Delta_1 s_1} \right] \Delta_2 s_2 \\
 &+ \int_{t_1}^b \left\{ h(s_1) - \left( \left( 1 - \frac{\alpha}{2} \right) h(b) + \frac{\alpha}{2} h(a) \right) \right\} \left[ \left\{ H(d) - \left( \left( 1 - \frac{\beta}{2} \right) H(d) + \frac{\beta}{2} H(c) \right) \right\} \right. \\
 &\quad \left. \frac{\partial f(s_1, t_2)}{\Delta_1 s_1} - \left\{ H(t_2) - \left( \left( 1 - \frac{\beta}{2} \right) H(d) + \frac{\beta}{2} H(c) \right) \right\} \frac{\partial f(s_1, c)}{\Delta_1 s_1} - \int_c^{t_2} H'(s_2) \frac{\partial f(s_1, \sigma(s_2))}{\Delta_1 s_1} \right] \Delta_2 s_2
 \end{aligned}$$

and

$$\begin{aligned}
 &= \left[ H(t_2) - \left( \left( 1 - \frac{\beta}{2} \right) H(c) + \frac{\beta}{2} H(d) \right) \right] \left\{ \left[ h(t_1) - \left( 1 - \frac{\alpha}{2} \right) h(a) - \frac{\alpha}{2} h(b) \right] f(t_1, t_2) \right. \\
 &\quad \left. - \left[ h(a) - \left( 1 - \frac{\alpha}{2} \right) h(a) - \frac{\alpha}{2} h(b) \right] f(a, t_2) - \int_a^{t_1} h'(s_1) f(\sigma(s_1), t_2) \Delta_1 s_1 \right\}
 \end{aligned}$$



$$\begin{aligned}
 & - \left[ H(c) - \left( \left( 1 - \frac{\beta}{2} \right) H(c) + \frac{\beta}{2} H(d) \right) \right] \left\{ \left[ h(t_1) - \left( 1 - \frac{\alpha}{2} \right) h(a) - \frac{\alpha}{2} h(b) \right] f(t_1, c) \right. \\
 & - \left. \left[ h(a) - \left( 1 - \frac{\alpha}{2} \right) h(a) - \frac{\alpha}{2} h(b) \right] f(a, c) - \int_a^{t_1} h'(s_1) f(\sigma(s_1), c) \Delta_1 s_1 \right\} \\
 & - \int_c^{t_2} H'(s_2) \left\{ \left[ h(t_1) - \left( 1 - \frac{\alpha}{2} \right) h(a) - \frac{\alpha}{2} h(b) \right] f(t_1, \sigma(s_2)) - \left[ h(a) - \left( 1 - \frac{\alpha}{2} \right) \right. \right. \\
 & \quad \left. \left. h(a) - \frac{\alpha}{2} h(b) \right] f(a, \sigma(s_2)) - \int_a^{t_1} h'(s_1) f(\sigma(s_1), \sigma(s_2)) \Delta_1 s_1 \right\} \Delta_2(s_2) \\
 \\
 & \left[ H(d) - \left( 1 - \frac{\beta}{2} \right) H(d) + \frac{\beta}{2} H(c) \right] \left\{ \left[ h(t_1) - \left( \left( 1 - \frac{\alpha}{2} \right) h(a) + \frac{\alpha}{2} h(b) \right) \right] f(t_1, d) \right. \\
 & - \left. \left[ h(a) - \left( \left( 1 - \frac{\alpha}{2} \right) h(a) + \frac{\alpha}{2} h(b) \right) \right] f(a, d) - \int_a^{t_1} h'(s_1) f(\sigma(s_1), d) \Delta_1 s_1 \right\} \\
 & - \left[ H(t_2) - \left( \left( 1 - \frac{\beta}{2} \right) H(d) + \frac{\beta}{2} H(c) \right) \right] \left\{ \left[ h(s_1) - \left( 1 - \frac{\alpha}{2} \right) h(a) - \frac{\alpha}{2} h(b) \right] f(t_1, t_2) \right. \\
 & - \left. \left[ h(a) - \left( 1 - \frac{\alpha}{2} \right) h(a) - \frac{\alpha}{2} h(b) \right] f(a, t_2) - \int_a^{t_1} h'(s_1) f(\sigma(s_1), d) \Delta_1 s_1 \right\} \\
 & - \int_{t_2}^d H'(s_2) \left\{ \left[ h(t_1) - \left( 1 - \frac{\alpha}{2} \right) h(a) - \frac{\alpha}{2} h(b) \right] f(t_1, \sigma(s_2)) - \left[ h(a) - \left( 1 - \frac{\alpha}{2} \right) h(a) \right. \right. \\
 & \quad \left. \left. - \frac{\alpha}{2} h(b) \right] f(a, \sigma(s_2)) - \int_a^{t_1} h'(s_1) f(\sigma(s_1), \sigma(s_2)) \Delta_1 s_1 \right\} \Delta_2(s_2) + \\
 \\
 & \left[ H(t_2) - \left( \left( 1 - \frac{\beta}{2} \right) H(c) + \frac{\beta}{2} H(d) \right) \right] \left\{ \left[ h(b) - \left( 1 - \frac{\alpha}{2} \right) h(b) - \frac{\alpha}{2} h(a) \right] f(b, t_2) \right. \\
 & - \left. \left[ h(t_1) - \left( 1 - \frac{\alpha}{2} \right) h(b) - \frac{\alpha}{2} h(a) \right] f(t_1, t_2) - \int_{t_1}^b h'(s_1) f(\sigma(s_1), t_2) \Delta_1 s_1 \right\} \\
 & - \left[ H(c) - \left( \left( 1 - \frac{\beta}{2} \right) H(c) + \frac{\beta}{2} H(d) \right) \right] \left\{ \left[ h(t_1) - \left( 1 - \frac{\alpha}{2} \right) h(b) - \frac{\alpha}{2} h(a) \right] f(b, c) \right. \\
 & - \left. \left[ h(t_1) - \left( 1 - \frac{\alpha}{2} \right) h(b) - \frac{\alpha}{2} h(a) \right] f(t_1, c) - \int_{t_1}^b h'(s_1) f(\sigma(s_1), c) \Delta_1 s_1 \right\} \\
 & - \int_c^y h'(t) \left\{ \left( h(b) - \left( \left( 1 - \frac{\alpha}{2} \right) h(b) + \frac{\alpha}{2} h(a) \right) \right) f(b, \sigma(t)) - \left[ h(t_1) - \left( 1 - \frac{\alpha}{2} \right) h(b) \right. \right. \\
 & \quad \left. \left. + \frac{\alpha}{2} h(a) \right] f(t_1, \sigma(s_2)) - \int_{t_1}^b h'(s_1) f(\sigma(s_1), \sigma(s_2)) \Delta_1 s_1 \right\} \Delta_2(s_2) +
 \end{aligned}$$

$$\begin{aligned}
 & \left[ H(d) - \left( \left( 1 - \frac{\beta}{2} \right) H(d) + \frac{\beta}{2} H(c) \right) \right] \left\{ \left[ h(b) - \left( 1 - \frac{\alpha}{2} \right) h(b) - \frac{\alpha}{2} h(a) \right] f(b, d) \right. \\
 & \quad \left. - \left[ h(t_1) - \left( 1 - \frac{\alpha}{2} \right) h(b) - \frac{\alpha}{2} h(a) \right] f(t_1, d) - \int_{t_1}^b h'(s_1) f(\sigma(s_1), \sigma(S_2)) \Delta_1 s_1 \right\} \\
 & - \left[ h(y) - \left( \left( 1 - \frac{\beta}{2} \right) H(d) + \frac{\beta}{2} H(c) \right) \right] \left\{ \left[ h(b) - \left( 1 - \frac{\alpha}{2} \right) h(b) - \frac{\alpha}{2} h(a) \right] f(b, t_2) \right. \\
 & \quad \left. - \left[ h(t_1) - \left( 1 - \frac{\alpha}{2} \right) h(b) + \frac{\alpha}{2} h(a) \right] f(t_1, t_2) - \int_{t_1}^b H'(s_2) f(\sigma(s_1), t_2) \Delta_1 s \right\} \\
 & - \int_{t_2}^d H'(s_2) \left\{ \left( h(t_1) - \left( \left( 1 - \frac{\alpha}{2} \right) h(a) + \frac{\alpha}{2} h(b) \right) \right) f(t_1, \sigma(s_2)) - \left[ h(a) - \left( 1 - \frac{\alpha}{2} \right) h(a) \right. \right. \\
 & \quad \left. \left. + \frac{\alpha}{2} h(b) \right] f(a, \sigma(t)) - \int_a^{t_1} h'(s_1) f(\sigma(s_1), \sigma(s_2)) \Delta_1 s_1 \right\} \Delta_2(s_2) \\
 & = \left( \frac{\beta}{2} H(d) - \frac{\beta}{2} H(c) \right) \left( h(b) - \left( \left( 1 - \frac{\alpha}{2} \right) h(b) + \frac{\alpha}{2} h(a) \right) \right) (f(b, c) - f(a, d) + f(b, d)) \\
 & \quad + \left( \frac{\alpha}{2} h(a) + \frac{\alpha}{2} h(b) \right) ((1 - \beta) H(c) + (\beta - 1) H(d)) (f(a, t_2) + f(b, t_2)) \\
 & \quad + \left( \frac{\beta}{2} H(c) + \frac{\beta}{2} H(d) \right) ((1 - \alpha) h(b) + (\alpha - 1) h(a)) (f(t_1, c) - f(t_1, d)) \\
 & \quad + ((1 - \alpha) h(b) + (\alpha - 1) h(a)) ((1 - \beta) H(d) + (\beta - 1) H(c)) f(t_1, t_2) \\
 & \quad + \left[ H(c) - \left( \left( 1 - \frac{\beta}{2} \right) H(c) + \frac{\beta}{2} H(d) \right) \right] \int_a^b h'(s_1) f(\sigma(s_1), c) \Delta_1 s_1 \\
 & \quad - \left[ h(t_2) - \left( \left( 1 - \frac{\beta}{2} \right) H(c) + \frac{\beta}{2} H(d) \right) \right] \int_a^b h'(s_1) f(\sigma(s_1), t_2) \Delta_1 s_1 \\
 & \quad - \int_c^d H'(s_2) ((1 - \alpha) h(a) + (\alpha - 1) h(b)) f(t_1, \sigma(s_2)) \Delta_2 s_2 \\
 & \quad + \int_c^d H'(s_2) \left\{ \left( h(a) - \left( \left( 1 - \frac{\alpha}{2} \right) h(a) + \frac{\alpha}{2} h(b) \right) \right) \right\} f(a, \sigma(s_2)) \Delta_2 s_2 \\
 & \quad + \int_c^d H'(s_2) \left\{ \int_a^b h'(s_1) f(\sigma(s_1), \sigma(s_2)) \Delta_1 s_1 \right\} \Delta_2(s_2)
 \end{aligned}$$

□

**Remark 1.** When  $\mathbb{T} = \mathbb{R}$

$$\begin{aligned}
 & \int_a^b \int_c^d W(t_1, t_2, s_1, s_2) \frac{d^2 f(s_1, s_2)}{d_1 s_1 d_2 s_2} d_1 s_1 d_2 s_2 \\
 & = \left( \frac{\beta}{2} H(d) - \frac{\beta}{2} H(c) \right) \left\{ h(b) - \left( \left( 1 - \frac{\alpha}{2} \right) h(b) + \frac{\alpha}{2} h(a) \right) \right\} (f(b, c) - f(a, d) + f(b, d))
 \end{aligned}$$

$$\begin{aligned}
 & + \left(\frac{\alpha}{2}h(a) + \frac{\alpha}{2}h(b)\right) \{(1 - \beta)H(c) + (\beta - 1)H(d)\} (f(a, t_2) + f(b, t_2)) \\
 & + \left(\frac{\beta}{2}H(c) + \frac{\beta}{2}H(d)\right) \{(1 - \alpha)h(b) + (\alpha - 1)h(a)\} (f(t_1, c) - f(t_1, d)) \\
 & + ((1 - \alpha)h(b) + (\alpha - 1)h(a)) ((1 - \beta)H(d) + (\beta - 1)H(c)) f(t_1, t_2) \\
 & + \left[H(c) - \left(\left(1 - \frac{\beta}{2}\right)H(c) + \frac{\beta}{2}H(d)\right)\right] \int_a^b h'(s_1)f(s_1, c)d_1s_1 \\
 & - \left[H(t_2) - \left(\left(1 - \frac{\beta}{2}\right)H(c) + \frac{\beta}{2}H(d)\right)\right] \int_a^b h'(s_1)f(s_1, t_2)d_1s_1 \\
 & - \int_c^d H'(s_2) ((1 - \alpha)h(a) + (\alpha - 1)h(b)) f(t_1, s_2)d_2s_2 \\
 & + \int_c^d H'(s_2) \left\{ \left(h(a) - \left(\left(1 - \frac{\alpha}{2}\right)h(a) + \frac{\alpha}{2}h(b)\right)\right) \right\} f(a, s_2)d_2s_2 \\
 & + \int_c^d H'(s_2) \left\{ \int_a^b h'(s_1)f(s_1, s_2)d_1s_1 \right\} d_2(t_2)
 \end{aligned}$$

**Remark 2.**  $\mathbb{T} = \mathbb{Z}$

$$\begin{aligned}
 & \sum_{s_1=a}^{b-1} \sum_{s_2=c}^{d-1} W(t_1, t_2, s_1, s_2) \\
 & = \left(\frac{\beta}{2}H(d) - \frac{\beta}{2}H(c)\right) \left\{ h(b) - \left(\left(1 - \frac{\alpha}{2}\right)h(b) + \frac{\alpha}{2}h(a)\right) \right\} (f(b, c) - f(a, d) + f(b, d)) \\
 & + \left(\frac{\alpha}{2}h(a) + \frac{\alpha}{2}h(b)\right) \{(1 - \beta)H(c) + (\beta - 1)H(d)\} (f(a, t_2) + f(b, t_2)) \\
 & + \left(\frac{\beta}{2}H(c) + \frac{\beta}{2}H(d)\right) \{(1 - \alpha)h(b) + (\alpha - 1)h(a)\} (f(t_1, c) - f(t_1, d)) \\
 & + ((1 - \alpha)h(b) + (\alpha - 1)h(a)) ((1 - \beta)H(d) + (\beta - 1)H(c)) f(t_1, t_2) \\
 & + \left[H(c) - \left(\left(1 - \frac{\beta}{2}\right)H(c) + \frac{\beta}{2}H(d)\right)\right] \sum_{s_1=a}^{b-1} h'(s_1)f(t_1 + 1, c) \\
 & - \left[H(t_2) - \left(\left(1 - \frac{\beta}{2}\right)H(c) + \frac{\beta}{2}H(d)\right)\right] \sum_{s_1=a}^{b-1} h'(s_1)f(t_1 + 1, t_2) \\
 & - \sum_{s_2=c}^{d-1} H'(s_2) ((1 - \alpha)h(a) + (\alpha - 1)h(b)) f(t_1, t_2 + 1) \\
 & + \sum_{s_2=c}^{d-1} H'(s_2) \left\{ \left(h(a) - \left(\left(1 - \frac{\alpha}{2}\right)h(a) + \frac{\alpha}{2}h(b)\right)\right) \right\} f(a, t_2 + 1) \\
 & + \sum_{s_1=a}^{b-1} \sum_{s_2=c}^{d-1} H'(s_2) \left\{ \int_a^b h'(s_1)f(t_1 + 1, t_2 + 1) \right\}
 \end{aligned}$$

**Remark 3.** By taking  $h(s_1) = s_1, H(s_2) = s_2$ , we obtain

$$\begin{aligned} & \int_a^b \int_c^d W(t_1, t_2, s_1, s_2) \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \Delta_1 s_1 \Delta_2 s_2 = (1-\alpha)(1-\beta)(b-a)(d-c)f(t_1, t_2) \\ & + (b-a)(d-c) \left\{ (1-\beta) \frac{\alpha}{2} [f(a, t_2) + f(b, t_2)] + \frac{\beta}{2} (1-\alpha) [f(t_1, c) + f(t_1, d)] \right\} \\ & + \frac{\alpha\beta}{4} (b-a)(d-c) \{f(b, c) + f(a, d) + f(b, d)\} - \frac{\beta}{2} (d-c) \int_a^b \{f(\sigma(s_1), c) \\ & + f(\sigma(s_1), d)\} \Delta_1 s_1 - \frac{\alpha}{2} (b-a) \int_c^d \{f(a, \sigma(t)) + f(b, \sigma(t))\} \Delta_2 t_2 \\ & + \int_a^b \int_c^d f(\sigma(s_1), \sigma(s_2)) \Delta_1 s_1 \Delta_2 s_2 \end{aligned}$$

**Theorem 5.** Under the conditions of Lemma, if  $f^{\Delta_1 \Delta_2} \in L_2((a, b)_{\mathbb{T}_1} \times (c, d)_{\mathbb{T}_2})$ , with  $h(s_1) = s_1, H(s_2) = s_2$ , then we have

$$\begin{aligned} & (1-\alpha)(1-\beta)(b-a)(d-c)f(t_1, t_2) + (b-a)(d-c) \left[ (1-\beta) \frac{\alpha}{2} [f(a, t_2) + f(b, t_2)] \right. \\ & \left. + \frac{\beta}{2} (1-\alpha) \{f(t_1, c) + f(t_1, d)\} \right] + \frac{\alpha\beta}{2} (b-a)(d-c) \{f(b, c) + f(a, d) + f(b, d)\} \\ & - \frac{\beta}{2} (d-c) \int_a^b \{f(\sigma(s_1), c) + f(\sigma(s_1), d)\} \Delta_1 s_1 - \frac{\alpha}{2} (b-a) \int_c^d \{f(a, \sigma(t)) \\ & + f(b, \sigma(t))\} \Delta_2 t_2 + \int_a^b \int_c^d f(\sigma(s_1), \sigma(s_2)) \Delta_1 s_1 \Delta_2 s_2 - \frac{[f(b, d) - f(a, d) - f(b, c) + f(a, c)]}{(b-a)^2(d-c)^2} \\ & \times \left[ h_2\left(t_1, a + \alpha \frac{b-a}{2}\right) - h_2\left(a, a + \alpha \frac{b-a}{2}\right) + h_2\left(b, b - \alpha \frac{b-a}{2}\right) - h_2\left(t_1, b - \alpha \frac{b-a}{2}\right) \right] \\ & \times \left[ h_2\left(t_2, c + \alpha \frac{d-c}{2}\right) - h_2\left(c, c + \alpha \frac{d-c}{2}\right) + h_2\left(d, d - \alpha \frac{d-c}{2}\right) - h_2\left(t_2, d - \alpha \frac{d-c}{2}\right) \right] \\ & \leq \left[ \left\{ \frac{b^3 - a^3}{3} - 2\left(a + \alpha \frac{b-a}{2}\right) \left(h_2\left(t_1, a + \alpha \frac{b-a}{2}\right)\right) - h_2\left(a, a + \alpha \frac{b-a}{2}\right) - \left(a + \alpha \frac{b-a}{2}\right)^2 (t_1 - a) \right\} \right. \\ & \left. - 2\left(b - \alpha \frac{b-a}{2}\right) \left( \left(h_2\left(b, b - \alpha \frac{b-a}{2}\right) - h_2\left(t_1, b - \alpha \frac{b-a}{2}\right)\right) - \left(b - \alpha \frac{b-a}{2}\right)^2 (b - t_1) \right) \right] \\ & \times \left[ \left\{ \frac{d^3 - c^3}{3} - 2\left(c + \alpha \frac{d-c}{2}\right) \left(h_2\left(t_2, c + \alpha \frac{d-c}{2}\right)\right) - h_2\left(c, c + \alpha \frac{d-c}{2}\right) - \left(c + \alpha \frac{d-c}{2}\right)^2 (t_2 - c) \right\} \right. \\ & \left. - 2\left(d - \alpha \frac{d-c}{2}\right) \left( \left(h_2\left(d, d - \alpha \frac{d-c}{2}\right) - h_2\left(t_2, d - \alpha \frac{d-c}{2}\right)\right) - \left(d - \alpha \frac{d-c}{2}\right)^2 (d - t_2) \right) \right] \\ & - \frac{1}{(b-a)(d-c)} \left[ \left( \left(h_2\left(t_1, a + \alpha \frac{b-a}{2}\right)\right) - h_2\left(a, a + \alpha \frac{b-a}{2}\right) + \left( \left(h_2\left(b, b - \alpha \frac{b-a}{2}\right)\right) \right) \right. \right. \end{aligned}$$

$$-h_2\left(t_1, b - \alpha \frac{b-a}{2}\right)\right)\right] \left[ \left( \left( h_2\left(t_2, c + \alpha \frac{d-c}{2}\right) \right) - h_2\left(c, c + \alpha \frac{d-c}{2}\right) \right) \right. \\ \left. + \left( \left( h_2\left(d, d - \alpha \frac{d-c}{2}\right) \right) - h_2\left(t_2, d - \alpha \frac{d-c}{2}\right) \right) \right] \sqrt{T(f\Delta_1\Delta_2)}.$$

where

$$T(f) = \int_a^b \int_c^d f^2(s_1, s_2) \Delta_2 s_2 \Delta_1 s_1 - \frac{1}{(b-a)(d-c)} \left( \int_a^b \int_c^d f(s_1, s_2) \Delta_2 s_2 \Delta_1 s_1 \right)^2$$

*Proof.* From the definition of  $W(t_1, t_2, s_1, s_2)$ , and taking  $h(s_1) = s_1, H(s_2) = s_2$ , we obtain

$$\int_a^b \int_c^d W(t_1, t_2, s_1, s_2) \Delta_2 s_2 \Delta_1 s_1 = \int_a^b W_1(t_1, s_1) \Delta_1 s_1 \int_c^d W_2(t_2, s_2) \Delta_2 s_2 \\ \left[ \int_a^b \left\{ s_1 - \left( a + \alpha \frac{b-a}{2} \right) \right\} \Delta_1 s_1 + \int_{t_1}^b \left\{ s_1 - \left( b - \alpha \frac{b-a}{2} \right) \right\} \Delta_1 s_1 \right] \\ \times \left[ \int_c^{t_2} \left\{ s_2 - \left( c + \alpha \frac{d-c}{2} \right) \right\} \Delta_2 s_2 + \int_{t_2}^d \left\{ s_2 - \left( d - \alpha \frac{d-c}{2} \right) \right\} \Delta_2 s_2 \right] \\ = \left[ h_2\left(t_1, a + \alpha \frac{b-a}{2}\right) - h_2\left(a, a + \alpha \frac{b-a}{2}\right) + h_2\left(b, b - \alpha \frac{b-a}{2}\right) - h_2\left(t_1, b - \alpha \frac{b-a}{2}\right) \right] \\ \times \left[ h_2\left(t_2, c + \alpha \frac{d-c}{2}\right) - h_2\left(c, c + \alpha \frac{d-c}{2}\right) + h_2\left(d, d - \alpha \frac{d-c}{2}\right) - h_2\left(t_2, d - \alpha \frac{d-c}{2}\right) \right]$$

and

$$\int_a^b \int_c^d W^2(t_1, t_2, s_1, s_2) \Delta_2 s_2 \Delta_1 s_1 = \int_a^b W_1^2(t_1, s_1) \Delta_1 s_1 \int_c^d W_2^2(t_2, s_2) \Delta_2 s_2 \\ = \left\{ \int_a^{t_1} \left[ s_1 - \left( a + \alpha \frac{b-a}{2} \right) \right]^2 \Delta_1 s_1 + \int_{t_1}^b \left[ s_1 - \left( b - \alpha \frac{b-a}{2} \right) \right]^2 \Delta_1 s_1 \right\} \Delta_1 s_1 \\ \times \left\{ \int_c^{t_2} \left[ s_2 - \left( c + \alpha \frac{d-c}{2} \right) \right]^2 \Delta_2 s_2 + \int_{t_2}^d \left[ s_2 - \left( d - \alpha \frac{d-c}{2} \right) \right]^2 \Delta_2 s_2 \right\} \Delta_2 s_2 \\ = \left[ \int_a^{t_1} \left\{ s_1^2 - 2 \left( a + \alpha \frac{b-a}{2} \right) \left( s_1 - \left( a + \alpha \frac{b-a}{2} \right) - \left( a + \alpha \frac{b-a}{2} \right)^2 \right) \right\} \Delta_1 s_1 \right. \\ \left. + \int_{t_1}^b \left\{ s_1^2 - 2 \left( b - \alpha \frac{b-a}{2} \right) \left( s_1 - \left( b - \alpha \frac{b-a}{2} \right) - \left( b - \alpha \frac{b-a}{2} \right)^2 \right) \right\} \Delta_1 s_1 \right] \\ \times \left[ \int_c^{t_2} \left\{ s_2^2 - 2 \left( c + \alpha \frac{d-c}{2} \right) \left( s_2 - \left( c + \alpha \frac{d-c}{2} \right) - \left( c + \alpha \frac{d-c}{2} \right)^2 \right) \right\} \Delta_2 s_2 \right. \\ \left. + \int_{t_2}^d \left\{ s_2^2 - 2 \left( d - \alpha \frac{d-c}{2} \right) \left( s_2 - \left( d - \alpha \frac{d-c}{2} \right) - \left( d - \alpha \frac{d-c}{2} \right)^2 \right) \right\} \Delta_2 s_2 \right]$$

$$\begin{aligned}
 &\leq \left[ \int_a^{t_1} \left\{ \frac{s_1^2 + s_1\sigma(s_1) + (\sigma(s_1))^2}{3} - 2 \left( a + \alpha \frac{b-a}{2} \right) \left( s_1 - \left( a + \alpha \frac{b-a}{2} \right) - \left( a + \alpha \frac{b-a}{2} \right)^2 \right) \Delta_1 s_1 \right\} \right] \\
 &\left[ \int_{t_1}^b \left\{ \frac{s_1^2 + s_1\sigma(s_1) + (\sigma(s_1))^2}{3} - 2 \left( b - \alpha \frac{b-a}{2} \right) \left( s_1 - \left( b - \alpha \frac{b-a}{2} \right) - \left( b - \alpha \frac{b-a}{2} \right)^2 \right) \Delta_1 s_1 \right\} \right] \\
 &\left[ \int_c^{t_2} \left\{ \frac{s_2^2 + s_2\sigma(s_2) + (\sigma(s_2))^2}{3} - 2 \left( c + \alpha \frac{d-c}{2} \right) \left( s_2 - \left( c + \alpha \frac{d-c}{2} \right) - \left( c + \alpha \frac{d-c}{2} \right)^2 \right) \Delta_2 s_2 \right\} \right] \\
 &\left[ \int_{t_2}^d \left\{ \frac{s_2^2 + s_2\sigma(s_2) + (\sigma(s_2))^2}{3} - 2 \left( d - \alpha \frac{d-c}{2} \right) \left( s_2 - \left( d - \alpha \frac{d-c}{2} \right) - \left( d - \alpha \frac{d-c}{2} \right)^2 \right) \Delta_2 s_2 \right\} \right] \\
 &= \frac{t_1^3 - a^3}{3} - 2 \left( a + \alpha \frac{b-a}{2} \right) \left[ h_2 \left( t_1, a + \alpha \frac{b-a}{2} \right) - h_2 \left( a, a + \alpha \frac{b-a}{2} \right) \right] - \left( a + \alpha \frac{b-a}{2} \right)^2 (t_1 - a) \\
 &+ \frac{b^3 - t_1^3}{3} - 2 \left( b - \alpha \frac{b-a}{2} \right) \left[ h_2 \left( b, b - \alpha \frac{b-a}{2} \right) - h_2 \left( t_1, b - \alpha \frac{b-a}{2} \right) \right] - \left( b - \alpha \frac{b-a}{2} \right)^2 (b - t_1) \\
 &+ \frac{t_2^3 - c^3}{3} - 2 \left( c + \alpha \frac{d-c}{2} \right) \left[ h_2 \left( t_2, c + \alpha \frac{d-c}{2} \right) - h_2 \left( c, c + \alpha \frac{d-c}{2} \right) \right] - \left( c + \alpha \frac{d-c}{2} \right)^2 (t_2 - c) \\
 &+ \frac{d^3 - t_2^3}{3} - 2 \left( d - \alpha \frac{d-c}{2} \right) \left[ h_2 \left( d, d - \alpha \frac{d-c}{2} \right) - h_2 \left( t_2, d - \alpha \frac{d-c}{2} \right) \right] - \left( d - \alpha \frac{d-c}{2} \right)^2 (d - t_2)
 \end{aligned}$$

Furthermore, we have

$$\begin{aligned}
 &\int_a^b \int_c^d \left[ W(t_1, t_2, s_1, s_2) - \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d W(t_1, t_2, s_1, s_2) \Delta_2 s_2 \Delta_1 s_1 \right] \Delta_2 s_2 \Delta_1 s_1 \\
 &\quad \times \left[ \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} - \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \right] \Delta_2 s_2 \Delta_1 s_1 \\
 &\int_a^b \int_c^d W(t_1, t_2, s_1, s_2) \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \Delta_2 s_2 \Delta_1 s_1 - \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d W(t_1, t_2, s_1, s_2) \Delta_2 s_2 \Delta_1 s_1 \\
 &\quad \times \int_a^b \int_c^d \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \Delta_2 s_2 \Delta_1 s_1 \\
 &= \int_a^b \int_c^d W(t_1, t_2, s_1, s_2) \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \Delta_2 s_2 \Delta_1 s_1 - \frac{[f(b, d) - f(a, d) - f(b, c) + f(a, c)]}{(b-a)(d-c)} \\
 &\quad \times \int_a^b \int_c^d W(t_1, t_2, s_1, s_2) \Delta_2 s_2 \Delta_1 s_1
 \end{aligned}$$

On the other hand

$$\begin{aligned}
 &\int_a^b \int_c^d \left[ W(t_1, t_2, s_1, s_2) - \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d W(t_1, t_2, s_1, s_2) \Delta_2 s_2 \Delta_1 s_1 \right] \Delta_2 s_2 \Delta_1 s_1 \\
 &\quad \times \left[ \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} - \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \right] \Delta_2 s_2 \Delta_1 s_1
 \end{aligned}$$

$$\begin{aligned} &\leq \left\| W(t_1, t_2, \dots) - \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d W(t_1, t_2, s_1, s_2) \Delta_2 s_2 \Delta_1 s_1 \right\|_2 \\ &\quad \times \left\| \frac{\partial^2 f(\cdot, \cdot)}{\Delta_1 s_1 \Delta_2 s_2} - \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \right\|_2 \\ &= \left[ \int_a^b \int_c^d W^2(t_1, t_2, s_1, s_2) \Delta_2 s_2 \Delta_1 s_1 - \frac{1}{(b-a)(d-c)} \left( \int_a^b \int_c^d W(t_1, t_2, s_1, s_2) \Delta_2 s_2 \Delta_1 s_1 \right)^2 \right]^{\frac{1}{2}} \\ &\quad \times \left[ \int_a^b \int_c^d \left( \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \right)^2 - \frac{1}{(b-a)(d-c)} \left( \int_a^b \int_c^d \frac{\partial^2 f(s_1, s_2)}{\Delta_1 s_1 \Delta_2 s_2} \right)^2 \right]^{\frac{1}{2}} \\ &\quad \times \left[ \int_a^b \int_c^d W^2(t_1, t_2, s_1, s_2) \Delta_2 s_2 \Delta_1 s_1 - \frac{1}{(b-a)(d-c)} \left( \int_a^b \int_c^d W(t_1, t_2, s_1, s_2) \Delta_2 s_2 \Delta_1 s_1 \right)^2 \right]^{\frac{1}{2}} \\ &\quad \times \sqrt{T(f^{\Delta_1 \Delta_2})}. \end{aligned}$$

□

#### 4. ACKNOWLEDGEMENT

We are thankful to Prof. Dr. A. D. Raza, Director General Abdus Salam School of Mathematical Sciences (ASSMS), GC University, Lahore-Pakistan for providing us the opportunity to avail the facilities from ASSMS to get this article complete

#### REFERENCES

- [1] R. Agarwal, M. Bohner and A. Peterson, Inequalities on time scales: a survey, *Math. Inequal. Appl.* 4 (2001), no. 4, 53517557.
- [2] M. Bohner, M. Fan and J. M. Zhang, Periodicity of scalar dynamic equations and applications to population models, *J. Math. Anal. Appl.* 330 (2007), 1179.
- [3] M. Bohner and T. Matthews, Ostrowski inequalities on time scales, *JIPAM. J. Inequal. Pure Appl. Math.* 9 (2008), no. 1, Article 6, 8 pp.
- [4] M. Bohner and A. Peterson, *Dynamic equations on time scales*, Birkhauser Boston, Boston, MA, 2001.
- [5] M. Bohner and A. Peterson, *Advances in dynamic equations on time scales*, Birkhauser Boston, Boston, MA, 2003.
- [6] M. Bohner and T. Matthews, Ostrowski inequalities on time scales, *JIPAM. J. Inequal. Pure Appl. Math.* 9 (2008), no. 1, Article 6, 8 pp.
- [7] S. S. Dragomir, The discrete version of Ostrowskis inequality in normed linear spaces, *JIPAM. J. Inequal. Pure Appl. Math.* 3 (2002), no. 1, Article 2, 16 pp.
- [8] C. Dinu, Ostrowski type inequalities on time scales, *An. Univ. Craiova Ser. Mat. Inform.* 34 (2007), 431758.

- [9] B. Karpuz and U. M. Ozkan, Generalized Ostrowski inequality on time scales, JIPAM. J. Inequal. Pure Appl. Math. 9 (2008), no. 4, Article 112, 7 pp.
- [10] W. J. Liu and Q. A. Ngo, An Ostrowski-Grüss type inequality on time scales, Comput. Math. Appl. 58 (2009), no. 6, 1207171210.
- [11] W. J. Liu and Q. A. Ngo, A generalization of Ostrowski inequality on time scales for  $k$  points, Appl. Math. Comput. 203 (2008), no. 2, 75417760.
- [12] W. J. Liu, Q. A. Ngo and W. B. Chen, A new generalization of Ostrowski type inequality on time scales, An. Stiint. Univ. Ovidius Constanta Ser. Mat. 17 (2009), no. 2, 10117114.
- [13] -D. S. Mitrinovic, J. E. Pecaric and A. M. Fink, Inequalities involving functions and their integrals and derivatives, Kluwer Acad. Publ., Dordrecht, 1991.
- [14] A. Ostrowski, Ueber die Absolutabweichung einer differentierbaren Funktion von ihren Integralmittelpunkt, Comment. Math. Helv. 10 (1938) 22617227.
- [15] U. M. Ozkan, M. Z. Sarikaya and H. Yildirim, Extensions of certain integral inequalities on time scales, Appl. Math. Lett. 21 (2008), no. 10, 993171000.
- [16] M. Z. Sarikaya, New weighted Ostrowski and Chebyshev type inequalities on time scales, Comput. Math. Appl. 60 (2010), no. 5, 1510171514.
- [17] M. Z. Sarikaya, N. Aktan and H. Yildirim, On weighted Chebyshev-Grüss type inequalities on time scales, J. Math. Inequal. 2 (2008), no. 2, 18517195.
- [18] A. Tuna, B. I. Yasar and S. Kutukcu, A note on integral inequalities involving the product  $\int_{\zeta_1}^{\zeta_2} f(x)g(x)dx$  of two functions on time scales, J. Appl. Funct. Anal. 3 (2008), no. 3, 34117346.
- [19] S. F. Wang, Q. L. Xue, W. J. Liu, Further generalization of Ostrowski-Grüss type inequalities, Adv. Appl. Math. Anal. 3 (2008), no. 1, 171720.

*E-mail address:* wchattah@hotmail.com

*E-mail address:* uetzone@hotmail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ENGINEERING AND TECHNOLOGY,, LAHORE, PAKISTAN

*E-mail address, corresponding Author:* malik.muddassar@gmail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ENGINEERING AND TECHNOLOGY,, TAXILA, PAKISTAN



# The Henstock-Stieltjes integral for fuzzy-number-valued functions on a infinite interval<sup>†</sup>

Ke-feng Duan\*

*College of Mathematics and Statistics, Longdong University, Qingyang Gansu, 745000, P.R. China*

**Abstract:** In this paper, the Henstock-Stieltjes integral for fuzzy-number-valued functions on a infinite interval which is an extension of the usual fuzzy Riemann-Stieltjes integral on infinite interval is firstly defined and discussed. Several necessary and sufficient conditions of the integrability for fuzzy-number-valued functions are given by means of the Henstock-Stieltjes integral of real-valued functions on infinite interval and Henstock integral of fuzzy-number-valued functions on infinite interval.

**Keywords:** Fuzzy numbers; Fuzzy Henstock integral; Stieltjes integral

**AMS subject classifications.** 26E50; 28E10.

## 1 Introduction

Recently, in order to complete the theory of fuzzy integrals and to meet the solving need of the fuzzy differential equations [1-3], fuzzy integrals of fuzzy-number-valued functions have been studied by many authors from different points of views, including Nanda [4], Wu et al. [5] and other authors [6-9]. As an extension for Riemann integral and Lebesgue integral, the Stieltjes integral plays an important role in probability theory, stochastic processes, physics, econometrics, biometrics and numerical analysis[10-13] in the Mathematics analysis. In fact, the establishment of the Stieltjes integral was related to the moment of inertia in physics [14]. Until 1909, Riesz presented a general expression for the linear functional of the space of the continuous functions in a finite interval by Stieltjes integral [15]. After Riesz' work, people find that the Stieltjes integral is a powerful tool in several branches of mathematics. In the fuzzy analysis, in 1968, Zadeh defined the probability measure of a fuzzy event by using the Lebesgue-Stieltjes integral of the membership function [16]. It is well known that the notion of the Stieltjes integral for fuzzy-number-valued functions was originally proposed by Nanda [4] in 1989. In 1998, Wu [17] discussed and defined the concept of fuzzy Riemann-Stieltjes integral by means of the representation theorem of fuzzy-number-valued functions, whose membership function could be obtained by solving a nonlinear programming problem, but it is difficult to calculate and extend to the higher-dimensional space. In 2006, Ren et al. introduced the concept of two kinds of fuzzy Riemann-Stieltjes integral for fuzzy-number-valued functions [18,19] and showed that a continuous fuzzy-number-valued function was fuzzy Riemann-Stieltjes integrable with respect to a real-valued increasing function. To overcome the limitations of the existing studies and to characterize continuous linear functionals on the space of Henstock integrable fuzzy-number-valued functions, in 2014, the concept of the Henstock-Stieltjes integral for fuzzy-number-valued functions is defined and discussed, and some useful results for this integral are shown [20].

The expectations of fuzzy random variables were investigated by M. L. Puri and D. A. Ralescu in 1986 [21]. It well known that the notion of a fuzzy random variable as a fuzzy-number-valued function and the expectation  $E(X)$  of a fuzzy random variable  $X$  was defined by a fuzzy integral  $E(X) = \int X$  or set-valued integral of  $X_\lambda$  [21]. In 2007, the concept of the fuzzy Henstock integral on infinite interval is proposed and discussed in order to solve the expectation  $E(X)$  of a fuzzy random variable  $X$  which distribution function has some kinds of discontinuity or non-integrability [7]. In this paper, the Henstock-Stieltjes integral for fuzzy-number-valued functions on infinite interval which is an extension of the usual fuzzy Riemann-Stieltjes integral on infinite interval is firstly defined and discussed. Several necessary and sufficient conditions of the integrability for fuzzy-number-valued functions are given by means of the

<sup>†</sup>The work is supported by the Natural Scientific Fund of Qingyang City (zj2014-10)

\* Tel.:+8618293439829, E-mail: kfduanldu@163.com

Henstock-Stieltjes integral of real-valued functions on infinite interval and Henstock integral of fuzzy-number-valued functions on infinite interval.

## 2 Preliminaries

Fuzzy set  $\tilde{u} \in E^1$  is called a fuzzy number if  $\tilde{u}$  is a normal, convex fuzzy set, upper semi-continuous and  $\text{supp } u = \{x \in \mathbb{R} \mid u(x) > 0\}$  is compact. Here  $\bar{A}$  denotes the closure of  $A$ . We use  $E^1$  to denote the fuzzy number space [22].

Let  $\tilde{u}, \tilde{v} \in E^1, k \in \mathbb{R}$ , the addition and scalar multiplication are defined by

$$[\tilde{u} + \tilde{v}]_\lambda = [\tilde{u}]_\lambda + [\tilde{v}]_\lambda, \quad [k\tilde{u}]_\lambda = k[\tilde{u}]_\lambda,$$

respectively, where  $[\tilde{u}]_\lambda = \{x : u(x) \geq \lambda\} = [u_\lambda^-, u_\lambda^+]$ , for any  $\lambda \in [0, 1]$ .

We use the Hausdorff distance between fuzzy numbers given by  $D : E^1 \times E^1 \rightarrow [0, +\infty)$  as follows [22]:

$$D(\tilde{u}, \tilde{v}) = \sup_{\lambda \in [0,1]} d([\tilde{u}]_\lambda, [\tilde{v}]_\lambda) = \sup_{\lambda \in [0,1]} \max\{|u_\lambda^- - v_\lambda^-|, |u_\lambda^+ - v_\lambda^+|\},$$

where  $d$  is the Hausdorff metric.  $D(\tilde{u}, \tilde{v})$  is called the distance between  $\tilde{u}$  and  $\tilde{v}$ .

**Lemma 2.1** [22]. If  $\tilde{u} \in E^1$ , then

- (1)  $[\tilde{u}]_\lambda$  is non-empty bounded closed interval for all  $\lambda \in [0, 1]$ ;
- (2)  $[\tilde{u}]_{\lambda_1} \supset [\tilde{u}]_{\lambda_2}$  for any  $0 \leq \lambda_1 \leq \lambda_2 \leq 1$ ;
- (3) for any  $\{\lambda_n\}$  converging increasingly to  $\lambda \in (0, 1]$ ,

$$\bigcap_{n=1}^{\infty} [\tilde{u}]_{\lambda_n} = [\tilde{u}]_\lambda.$$

Conversely, if for all  $\lambda \in [0, 1]$ , there exists  $A_\lambda \subset \mathbb{R}$  satisfying (1)  $\sim$  (3), then there exists a unique  $\tilde{u} \in E^1$  such that  $[\tilde{u}]_\lambda = A_\lambda, \lambda \in (0, 1]$ , and  $[\tilde{u}]_0 = \bigcup_{\lambda \in (0,1]} \overline{[\tilde{u}]_\lambda} \subset A_0$ .

**Definition 2.1** [7, 20, 23].  $\bar{R}$  denote the generalized real line, for  $\tilde{f}$  defined on  $[a, +\infty)$ , we define  $\tilde{f}(+\infty) = \tilde{0}$ , and  $\tilde{0} \cdot (+\infty) = \tilde{0}$ .

Let  $\delta : [a, +\infty) \rightarrow R^+$  be a positive real function. A division  $P = \{[x_{i-1}, x_i]; \xi_i\}$  is said to be  $\delta$ -fine, if the following conditions are satisfied:

- (1)  $a = x_0 < x_1 < \dots < x_{n-1} = b < x_n = +\infty$ ;
- (2)  $\xi_i \in [x_{i-1}, x_i] \subset O(\xi_i), i = 1, 2, \dots, n$ ;

where  $O(\xi_i) = (\xi_i - \delta(\xi_i), \xi_i + \delta(\xi_i))$  for  $i = 1, 2, \dots, n - 1$ , and  $O(\xi_n) = [b, +\infty)$ .

For brevity, we write  $T = \{[u, v]; \xi\}$ , where  $[u, v]$  denotes a typical interval in  $T$  and  $\xi$  is the associated point of  $[u, v]$ .

**Definition 2.2.** Let  $\alpha : [a, +\infty) \rightarrow \mathbb{R}$  be an increasing function. A function  $f : [a, +\infty) \rightarrow \mathbb{R}$  is Henstock-Stieltjes integrable with respect to  $\alpha$  on  $[a, +\infty)$  if there exists a real number  $I$  such that for every  $\varepsilon > 0$ , there is a function  $\delta(x) > 0$  on  $[a, +\infty)$  such that for any  $\delta$ -fine division  $T = \{[x_{i-1}, x_i]; \xi_i\}_{i=1}^n$ , we have

$$\left| \sum_{i=1}^n f(\xi_i)[\alpha(x_i) - \alpha(x_{i-1})] - I \right| < \varepsilon.$$

As usual, we write  $(HS) \int_a^{+\infty} f(x)d\alpha = I$  and  $(f, \alpha) \in HS[a, +\infty)$ .

Recall, also, that a function  $\tilde{f} : [a, b] \rightarrow E^1$  is said to be bounded if there exists  $M \in \mathbb{R}$  such that  $\|\tilde{f}(x)\| = D(\tilde{f}(x), \tilde{0}) \leq M$  for any  $x \in [a, b]$ . Notice that here  $\|\tilde{f}(x_0)\|$  does not stand for the norm of  $E^1$ .

## 3 The fuzzy Henstock-Stieltjes integral on infinite interval and its properties

In this section we shall give the definition of the Henstock-Stieltjes integral for fuzzy-number-valued functions on a infinite interval.

**Definition 3.1.** Let  $\alpha : [a, +\infty) \rightarrow \mathbb{R}$  be an increasing function. A fuzzy-number-valued function  $\tilde{f}(x)$  is said to be fuzzy Henstock-Stieltjes integrable with respect to  $\alpha$  on  $[a, +\infty)$  if there exists a fuzzy

number  $\tilde{H} \in E^1$  such that for every  $\varepsilon > 0$ , there is a function  $\delta(x) > 0$  on  $[a, +\infty]$  such that for any  $\delta$ -fine division  $T = \{[x_{i-1}, x_i]; \xi_i\}_{i=1}^n$ , we have

$$D\left(\sum_{i=1}^n \tilde{f}(\xi_i)[\alpha(x_i) - \alpha(x_{i-1})], \tilde{H}\right) < \varepsilon.$$

We write  $(FHS) \int_a^{+\infty} \tilde{f}(x)d\alpha = \tilde{H}$  and  $(\tilde{f}, \alpha) \in FHS[a, +\infty]$ .

The definition of  $\tilde{f} \in FHS(-\infty, a]$  is similar. Naturally, we define  $\tilde{f} \in FHS(-\infty, +\infty)$  iff  $\tilde{f} \in FHS(-\infty, a]$  and  $\tilde{f} \in FHS[a, +\infty)$ , and furthermore

$$(FHS) \int_{-\infty}^{+\infty} \tilde{f}(x)d\alpha = (FHS) \int_{-\infty}^a \tilde{f}(x)d\alpha + (FHS) \int_a^{+\infty} \tilde{f}(x)d\alpha.$$

**Remark 3.1.** It is clear, if  $\tilde{f}(x)$  is a real-valued function then Definition 3.1 implies the definition of  $(HS)$  integral on infinite interval introduced by [20]; if  $\alpha(x) = x$ , then Definition 3.1 implies the definition of  $(FH)$  integral introduced by Gong et al. [7].

**Remark 3.2.** From the definition of the fuzzy Henstock-Stieltjes integral and the fact that  $(E^1, D)$  is a complete metric space, we can easily obtain the following conclusions.

**Theorem 3.1.** Let  $\alpha : [a, +\infty] \rightarrow \mathbb{R}$  be an increasing function. A fuzzy-number-valued function  $\tilde{f}$  is fuzzy Henstock-Stieltjes integrable with respect to  $\alpha$  on  $[a, +\infty]$  if and only if for every  $\varepsilon > 0$ , there is a function  $\delta(x) > 0$  on  $[a, +\infty]$  such that for any  $\delta$ -fine division  $T = \{[u, v]; \xi\}$  and  $T' = \{[u', v']; \xi'\}$ , we have

$$D\left(\sum_T \tilde{f}(\xi)[\alpha(v) - \alpha(u)], \sum_{T'} \tilde{f}(\xi')[\alpha(v') - \alpha(u')]\right) < \varepsilon.$$

**Theorem 3.2.** Let  $\alpha : [a, +\infty] \rightarrow \mathbb{R}$  be an increasing function and let  $\tilde{f} : [a, +\infty] \rightarrow E^1$ . Then the following statements are equivalent:

- (1)  $(\tilde{f}, \alpha) \in FHS[a, +\infty]$  and  $(FHS) \int_a^{+\infty} \tilde{f}(x)d\alpha = \tilde{A}$ ;
- (2) for any  $\lambda \in [0, 1]$ ,  $f_\lambda^-$  and  $f_\lambda^+$  are Henstock-Stieltjes integrable with respect to  $\alpha$  on  $[a, +\infty]$  for any  $\lambda \in [0, 1]$  uniformly ( $\delta(x)$  is independent of  $\lambda \in [0, 1]$ ), and

$$[(FHS) \int_a^{+\infty} \tilde{f}(x)d\alpha]_\lambda = [(HS) \int_a^{+\infty} f_\lambda^-(x)d\alpha, (HS) \int_0^{+\infty} f_\lambda^+(x)d\alpha].$$

- (3) For any  $b > a$ ,  $\tilde{f} \in FH[a, b]$ ,  $\lim_{b \rightarrow +\infty} \int_a^b \tilde{f}(x)d\alpha$  as a fuzzy number exists and

$$\lim_{b \rightarrow +\infty} \int_a^b \tilde{f}(x)d\alpha = \int_a^{+\infty} \tilde{f}(x)d\alpha.$$

**Proof.** First, we prove that (1) is equivalent to (2).

(1) implies (2): If  $\int_a^{+\infty} \tilde{f}(x)d\alpha = \tilde{A}$ , then given  $\varepsilon > 0$ , there exists a positive-valued function  $\delta(x)$  on  $[a, +\infty]$  such that for any  $\delta$ -fine division of  $[a, +\infty] : T = \{[x_{i-1}, x_i]; \xi_i\}$ , we have

$$D\left(\sum_i \tilde{f}(\xi_i)(\alpha(x_i) - \alpha(x_{i-1})), \tilde{A}\right) < \varepsilon,$$

i.e.

$$\sup_{\lambda \in [0, 1]} \max\{|\left[\sum_i \tilde{f}(\xi_i)(\alpha(x_i) - \alpha(x_{i-1}))\right]_\lambda^- - A_\lambda^-|, |\left[\sum_i \tilde{f}(\xi_i)(\alpha(x_i) - \alpha(x_{i-1}))\right]_\lambda^+ - A_\lambda^+|\} < \varepsilon,$$

so for any  $\lambda \in [0, 1]$  and any  $\delta$ -fine division  $T = \{[x_{i-1}, x_i]; \xi_i\}$ , we have

$$|\left[\sum_i \tilde{f}(\xi_i)(\alpha(x_i) - \alpha(x_{i-1}))\right]_\lambda^- - A_\lambda^-| = \left|\sum_i f_\lambda^-(\xi_i)(\alpha(x_i) - \alpha(x_{i-1})) - A_\lambda^-\right| < \varepsilon,$$

$$|[\sum_i \tilde{f}(\xi_i)(\alpha(x_i) - \alpha(x_{i-1}))]_{\lambda}^+ - A_{\lambda}^+| = |\sum_i f_{\lambda}^+(\xi_i)(\alpha(x_i) - \alpha(x_{i-1})) - A_{\lambda}^+| < \varepsilon.$$

Thus,  $f_{\lambda}^-$  and  $f_{\lambda}^+$  are Henstock integrable uniformly with respect to  $\lambda \in [0, 1]$  on  $[a, +\infty)$ , and

$$A_{\lambda} = [\int_a^{+\infty} f_{\lambda}^-(x)d\alpha, \int_a^{+\infty} f_{\lambda}^+(x)d\alpha].$$

Conversely, since  $f_{\lambda}^-$  and  $f_{\lambda}^+$  are Henstock integrable uniformly with respect to  $\lambda \in [0, 1]$  on  $[a, +\infty)$ , then given  $\varepsilon > 0$ , there exists a positive-valued function  $\delta(x)$  on  $[a, +\infty]$  such that for any  $\delta$ -fine division  $T = \{[x_{i-1}, x_i]; \xi_i\}$ , and for any  $\lambda \in [0, 1]$ , we have

$$|\sum_i f_{\lambda}^-(\xi_i)(\alpha(x_i) - \alpha(x_{i-1})) - A_{\lambda}^-| < \varepsilon, \tag{0.1}$$

$$|\sum_i f_{\lambda}^+(\xi_i)(\alpha(x_i) - \alpha(x_{i-1})) - A_{\lambda}^+| < \varepsilon. \tag{0.2}$$

We can prove that the class of closed intervals  $\{[A_{\lambda}^-, A_{\lambda}^+] : \lambda \in [0, 1]\}$  determines a fuzzy number. In fact,  $[A_{\lambda}^-, A_{\lambda}^+]$  satisfies the conditions of lemma 2.1.

- (1) Since  $f_{\lambda}^-(x) \leq f_{\lambda}^+(x), \lambda \in [0, 1]$ , then  $A_{\lambda}^- \leq A_{\lambda}^+$ , i.e.  $[A_{\lambda}^-, A_{\lambda}^+]$  is a closed interval,  $\lambda \in [0, 1]$ ,
- (2) For any  $0 \leq \lambda_1 \leq \lambda_2 \leq 1$ ,

$$f_{\lambda_1}^-(x) \leq f_{\lambda_2}^-(x) \leq f_{\lambda_2}^+(x) \leq f_{\lambda_1}^+(x).$$

This implies

$$\int_a^{+\infty} f_{\lambda_1}^-(x)d\alpha \leq \int_a^{+\infty} f_{\lambda_2}^-(x)d\alpha \leq \int_a^{+\infty} f_{\lambda_2}^+(x)d\alpha \leq \int_a^{+\infty} f_{\lambda_1}^+(x)d\alpha.$$

That is,  $[A_{\lambda_1}^-, A_{\lambda_1}^+] \supset [A_{\lambda_2}^-, A_{\lambda_2}^+]$ .

- (3) For any  $\{\lambda_n\}$  increasingly converging to  $\lambda \in (0, 1]$ ,

$$\bigcap_{n=1}^{\infty} [\tilde{f}(x)]_{\lambda_n} = [\tilde{f}(x)]_{\lambda},$$

i.e.

$$\bigcap_{n=1}^{\infty} [f_{\lambda_n}^-(x), f_{\lambda_n}^+(x)] = [f_{\lambda}^-(x), f_{\lambda}^+(x)].$$

That is

$$\lim_{n \rightarrow \infty} f_{\lambda_n}^-(x) = f_{\lambda}^-(x), \lim_{n \rightarrow \infty} f_{\lambda_n}^+(x) = f_{\lambda}^+(x).$$

Note that

$$f_0^-(x) \leq f_{\lambda_n}^-(x) \leq f_1^-(x), f_1^+(x) \leq f_{\lambda_n}^+(x) \leq f_0^+(x).$$

This implies

$$0 \leq f_{\lambda_n}^-(x) - f_0^-(x) \leq f_1^-(x) - f_0^-(x), 0 \leq f_{\lambda_n}^+(x) - f_1^+(x) \leq f_0^+(x) - f_1^+(x).$$

By the non-negativeness and Henstock integrability of  $f_1^- - f_0^-, f_0^+ - f_1^+$ , we know that  $f_1^- - f_0^-, f_0^+ - f_1^+$  are Lebesgue integrable (refer to [9]). Hence  $f_{\lambda_n}^-(x) - f_0^-(x), f_{\lambda_n}^+(x) - f_1^+(x)$  are Lebesgue integrable, and

$$\lim_{n \rightarrow \infty} \int_a^{+\infty} (f_{\lambda_n}^-(x) - f_0^-(x))d\alpha = \int_a^{+\infty} (f_{\lambda}^-(x) - f_0^-(x))d\alpha,$$

$$\lim_{n \rightarrow \infty} \int_a^{+\infty} (f_{\lambda_n}^+(x) - f_1^+(x))d\alpha = \int_a^{+\infty} (f_{\lambda}^+(x) - f_1^+(x))d\alpha.$$

That is

$$\lim_{n \rightarrow \infty} \int_a^{+\infty} f_{\lambda_n}^-(x)d\alpha = \int_a^{+\infty} f_{\lambda}^-(x)d\alpha, \lim_{n \rightarrow \infty} \int_a^{+\infty} f_{\lambda_n}^+(x)d\alpha = \int_a^{+\infty} f_{\lambda}^+(x)d\alpha.$$

Thus,  $\bigcap_{n=1}^{\infty} [A_{\lambda_n}^-, A_{\lambda_n}^+] = [A_{\lambda}^-, A_{\lambda}^+]$ .

Combining the inequality (1) and (2) we obtain

$$D\left(\sum_i (\alpha(x_i) - \alpha(x_{i-1}))\tilde{f}(\xi_i), \tilde{A}\right) < \varepsilon,$$

i.e.

$$\tilde{f} \in FH[a, +\infty), \int_a^{+\infty} \tilde{f}(x)d\alpha = \tilde{A}.$$

We'll prove that (i) is equivalent to (iii) as follows.

(1) *implies* (3): Let  $\varepsilon > 0$ . Suppose  $\tilde{f} \in FH[a, +\infty)$ . There exists a positive-valued function  $\delta$  on  $[a, +\infty)$  such that

$$D\left(\sum_i \tilde{f}(\xi_i)(\alpha(x_i) - \alpha(x_{i-1})), \int_a^{+\infty} \tilde{f}\right) < \varepsilon.$$

for any  $\delta$ -fine division of  $[a, +\infty) : T = \{[x_{i-1}, x_i]; \xi_i\}_{i=1}^n$ . On the other hand, by the Cuachy Rule about  $\tilde{f} \in FH[a, b]$  (refer to Th 2.3 of [24]), then  $\tilde{f} \in FH[a, b]$  for any  $b > a$ . There is a positive-valued function  $\delta_1$  on  $[a, b]$  such that for any  $\delta_1$ -fine division of  $[a, b] : T = \{[x_{i-1}, x_i]; \xi_i\}_{i=1}^n$ , we have

$$D\left(\sum_i \tilde{f}(\xi_i)(\alpha(x_i) - \alpha(x_{i-1})), \int_a^b \tilde{f}\right) < \varepsilon.$$

We may assume that  $\delta_1 \leq \delta$  for any  $\xi \in [a, b]$ . Then

$$\begin{aligned} & D\left(\int_a^{+\infty} \tilde{f}, \int_a^b \tilde{f}\right) \\ & \leq D\left(\sum_i \tilde{f}(\xi_i)(\alpha(x_i) - \alpha(x_{i-1})), \int_a^{+\infty} \tilde{f}\right) + D\left(\sum_i \tilde{f}(\xi_i)(\alpha(x_i) - \alpha(x_{i-1})), \int_a^b \tilde{f}\right) + D(\tilde{f}(+\infty)\mu([b, +\infty)) \\ & < \varepsilon + \varepsilon = 2\varepsilon. \end{aligned}$$

Hence

$$\lim_{b \rightarrow +\infty} \int_a^b \tilde{f}(x)d\alpha = \int_a^{+\infty} \tilde{f}(x)d\alpha.$$

(3) *implies* (1): Let  $\varepsilon > 0$ . Choose a sequence  $a = b_0 < b_1 < b_2 < \dots, b_k \uparrow +\infty$ . Since  $\tilde{f} \in FH[b_{k-1}, b_k], k = 1, 2, 3, \dots$ , there exist  $\delta_k$  such that

$$D\left(\sum_{[b_{k-1}, b_k]} \tilde{f}(\xi)(v - u), \int_{b_{k-1}}^{b_k} \tilde{f}\right) < \varepsilon/2^{k+2}.$$

for any  $\delta_k$ -fine division on  $[b_{k-1}, b_k], k = 1, 2, 3, \dots$ . Suppose  $\lim_{b \rightarrow +\infty} \int_a^b \tilde{f}(x)d\alpha = \tilde{A}$ . Choose  $N$  such that

$b > b_N$  which implies  $D(\int_a^b \tilde{f}(x)d\alpha, \tilde{A}) < \varepsilon/2$ .

Define

$$\delta(\xi) = \begin{cases} \delta_1(\xi), & \xi \in [b_0, b_1), \\ \delta_k(\xi), & \xi \in (b_{k-1}, b_k), k = 1, 2, 3, \dots, \\ \min(\delta_k(b_k), \delta_{k+1}(b_k)) & \xi = b_k, k = 1, 2, 3, \dots \end{cases}$$

For any  $\delta$ -fine division  $P = \{[x_{i-1}, x_i]; \xi_i\}$  satisfies  $i = 1, 2, \dots, n$ ,  $O(\xi_n) = [b, +\infty)$  and  $b > b_N$ , we have

$$\begin{aligned} & D\left(\sum_i \tilde{f}(\xi_i)(\alpha(x_i) - \alpha(x_{i-1})), \tilde{A}\right) \\ & \leq D\left(\int_a^b \tilde{f}, \tilde{A}\right) + D\left(\int_a^b \tilde{f}, \sum_i \tilde{f}(\xi_i)(\alpha(x_i) - \alpha(x_{i-1}))\right) \\ & < \varepsilon/2 + D\left(\int_a^b \tilde{f}, \sum_{i=1}^{n-1} \tilde{f}(\xi_i)(\alpha(x_i) - \alpha(x_{i-1})) + \tilde{f}(+\infty)\mu([b, +\infty))\right) \\ & \leq \varepsilon/2 + \sum_{k=1}^{+\infty} \varepsilon/2^{k+2} = 2\varepsilon \end{aligned}$$

Hence,  $\tilde{f} \in FH[a, +\infty)$  and

$$\lim_{b \rightarrow +\infty} \int_a^b \tilde{f}(x) d\alpha = \int_a^{+\infty} \tilde{f}(x) dx.$$

The proof is complete.

**Theorem 3.3.** Let  $\alpha : [a, +\infty) \rightarrow \mathbb{R}$  be an increasing function such that  $\alpha \in C^1[a, +\infty)$  and  $\tilde{f} : [a, +\infty) \rightarrow E^1$  be a bounded fuzzy-number-valued function. Then  $\tilde{f}$  is fuzzy Henstock-Stieltjes integrable with respect to  $\alpha$  on  $[a, +\infty)$  if and only if  $\tilde{f}\alpha'$  is fuzzy Henstock integrable on  $[a, +\infty)$ . Furthermore, we have

$$(FHS) \int_a^{+\infty} \tilde{f}(x) d\alpha = (FH) \int_a^{+\infty} \tilde{f}(x)\alpha'(x) dx,$$

where  $(FH)$  integral denotes the fuzzy Henstock integral introduced by Wu et al. [5].

**Proof.** Since  $\tilde{f} : [a, +\infty) \rightarrow E^1$  is bounded on  $[a, +\infty)$ ,  $\sup_{x \in [a, +\infty)} D(\tilde{f}(x), \tilde{0})$  exists. Continuity of  $\alpha'$  on  $[a, b]$  implies uniform continuity on  $[a, b]$  for any  $b > a$ . Hence, for each  $\epsilon > 0$ , there exists  $\eta > 0$  such that

$$|\alpha'(x) - \alpha'(y)| < \frac{\epsilon}{3 \sup_{x \in [a, +\infty)} D(\tilde{f}(x), \tilde{0}) \cdot (b - a)}$$

for any  $x, y \in [a, b]$  satisfying  $|x - y| < \eta$ . Choose a positive-valued function  $\delta_1(x)$  on  $[a, b]$  with  $\delta_1(x) < \eta$  for all  $x \in [a, b]$ . Let  $T = \{[x_{i-1}, x_i]; \xi_i\}_{i=1}^n$  be a  $\delta_1$ -fine division on  $[a, b]$ , then by Lagrange mean value theorem, there exists  $\bar{x}_i \in [x_{i-1}, x_i]$  such that

$$\alpha(x_i) - \alpha(x_{i-1}) = \alpha'(\bar{x}_i)(\alpha(x_i) - \alpha(x_{i-1})), \quad (1 \leq i \leq n).$$

Since  $|\bar{x}_i - x_i| \leq \delta_1(x_i) < \eta$  for  $1 \leq i \leq n$ , we have

$$|\alpha'(\bar{x}_i) - \alpha'(x_i)| < \frac{\epsilon}{3 \sup_{x \in [a, +\infty)} D(\tilde{f}(x), \tilde{0}) \cdot (b - a)}$$

for  $1 \leq i \leq n$ . Hence, for any  $\delta_1$ -fine division  $T = \{[x_{i-1}, x_i]; \xi_i\}_{i=1}^n$  on  $[a, b]$ , we have

$$\begin{aligned} & D\left(\sum_{i=1}^n \tilde{f}(\xi_i)[\alpha(x_i) - \alpha(x_{i-1})], \sum_{i=1}^n \tilde{f}(\xi_i)\alpha'(\xi_i)(\alpha(x_i) - \alpha(x_{i-1}))\right) \\ & = D\left(\sum_{i=1}^n \tilde{f}(\xi_i)\alpha'(\bar{x}_i)(\alpha(x_i) - \alpha(x_{i-1})), \sum_{i=1}^n \tilde{f}(\xi_i)\alpha'(\xi_i)(\alpha(x_i) - \alpha(x_{i-1}))\right) \\ & \leq \sum_{i=1}^n D(\tilde{f}(\xi_i)\alpha'(\bar{x}_i)(\alpha(x_i) - \alpha(x_{i-1})), \tilde{f}(\xi_i)\alpha'(\xi_i)(\alpha(x_i) - \alpha(x_{i-1}))) \end{aligned}$$

$$\begin{aligned}
 &= \sum_{i=1}^n \sup_{\lambda \in [0,1]} \max\{|f_{\lambda}^{-}(\xi_i)[\alpha'(\bar{x}_i) - \alpha'(\xi_i)|, |f_{\lambda}^{+}(\xi_i)[\alpha'(\bar{x}_i) - \alpha'(\xi_i)]|\}(\alpha(x_i) - \alpha(x_{i-1})) \\
 &\leq \sum_{i=1}^n |\alpha'(\bar{x}_i) - \alpha'(\xi_i)|(\alpha(x_i) - \alpha(x_{i-1})) \sup_{\lambda \in [0,1]} \max\{|f_{\lambda}^{-}(\xi_i)|, |f_{\lambda}^{+}(\xi_i)|\} \\
 &\leq (b-a) \cdot \frac{\epsilon}{3 \sup_{x \in [a,+\infty)} D(\tilde{f}(x), \tilde{0}) \cdot (b-a)} \cdot \sup_{x \in [a,+\infty)} D(\tilde{f}(x), \tilde{0}) \\
 &< \frac{\epsilon}{3}. \tag{*}
 \end{aligned}$$

On the other hand, since  $\tilde{f}$  is fuzzy Henstock-Stieltjes integrable with respect to  $\alpha$  on  $[a, +\infty)$ , by Theorem 3.1, there is a function  $\delta_2(x) > 0$  such that for any  $\delta_2$ -fine division  $T = \{[u, v]; \xi\}$  and  $T' = \{[u', v']; \xi'\}$  on  $[a, b]$ , we have

$$D\left(\sum_T \tilde{f}(\xi)[\alpha(v) - \alpha(u)], \sum_{T'} \tilde{f}(\xi')[\alpha(v') - \alpha(u')]\right) < \frac{\epsilon}{3}.$$

Define  $\delta(x)$  on  $[a, b]$  by  $\delta(x) = \min\{\delta_1(x), \delta_2(x)\}$ . Then for any  $\delta$ -fine division  $T = \{[u, v]; \xi\}$  and  $T' = \{[u', v']; \xi'\}$  on  $[a, b]$ , we have

$$\begin{aligned}
 &D\left(\sum_T \tilde{f}(\xi)\alpha'(\xi)(v-u), \sum_{T'} \tilde{f}(\xi')\alpha'(\xi')(v'-u')\right) \\
 &\leq D\left(\sum_T \tilde{f}(\xi)\alpha'(\xi)(v-u), \sum_T \tilde{f}(\xi)[\alpha(v) - \alpha(u)]\right) \\
 &+ D\left(\sum_T \tilde{f}(\xi)[\alpha(v) - \alpha(u)], \sum_{T'} \tilde{f}(\xi')[\alpha(v') - \alpha(u')]\right) \\
 &+ D\left(\sum_{T'} \tilde{f}(\xi')[\alpha(v') - \alpha(u')], \sum_{T'} \tilde{f}(\xi')\alpha'(\xi')(v'-u')\right) \\
 &< \epsilon.
 \end{aligned}$$

Hence,  $\tilde{f}\alpha'$  is Henstock integrable on  $[a, b]$  for any  $[a, b]$  by Theorem 2.3 of [5], and by above formula (\*), we know that

$$(FHS) \int_a^b \tilde{f}(x)d\alpha = (FH) \int_a^b \tilde{f}(x)\alpha'(x)dx.$$

Applied Theorem 3.1,  $\tilde{f}\alpha'$  is Henstock integrable on  $[a, +\infty)$ .

Conversely, if  $\tilde{f}\alpha'$  is Henstock integrable on  $[a, +\infty)$ , then by Theorem 2.3 of [5], for each  $\epsilon > 0$ , there is a function  $\delta_3(x) > 0$  such that for any  $\delta_3$ -fine division  $T = \{[u, v]; \xi\}$  and  $T' = \{[u', v']; \xi'\}$ , we have

$$D\left(\sum_T \tilde{f}(\xi)\alpha'(\xi)(v-u), \sum_{T'} \tilde{f}(\xi')\alpha'(\xi')(v'-u')\right) < \frac{\epsilon}{3}.$$

Define  $\delta(x)$  on  $[a, +\infty)$  by  $\delta(x) = \min\{\delta_1(x), \delta_3(x)\}$ . Then for any  $\delta$ -fine division  $T = \{[u, v]; \xi\}$  and  $T' = \{[u', v']; \xi'\}$ , we have

$$\begin{aligned}
 &D\left(\sum_T \tilde{f}(\xi)[\alpha(v) - \alpha(u)], \sum_{T'} \tilde{f}(\xi')[\alpha(v') - \alpha(u')]\right) \\
 &\leq D\left(\sum_T \tilde{f}(\xi)[\alpha(v) - \alpha(u)], \sum_T \tilde{f}(\xi)\alpha'(\xi)(v-u)\right) \\
 &+ D\left(\sum_T \tilde{f}(\xi)\alpha'(\xi)(v-u), \sum_{T'} \tilde{f}(\xi')\alpha'(\xi')(v'-u')\right) \\
 &+ D\left(\sum_{T'} \tilde{f}(\xi')\alpha'(\xi')(v'-u'), \sum_{T'} \tilde{f}(\xi')[\alpha(v') - \alpha(u')]\right) \\
 &< \epsilon.
 \end{aligned}$$

Hence,  $\tilde{f}$  is fuzzy Henstock-Stieltjes integrable with respect to  $\alpha$  on  $[a, +\infty)$ .

In the following part, we will prove the equation  $(FHS) \int_a^{+\infty} \tilde{f} d\alpha = (FH) \int_a^{+\infty} \tilde{f} \alpha' dx$ . For any division  $T : a = x_0 < x_1 < x_2 < \dots < x_n = b$ , according to the Lagrange mean value theorem, we have

$$\alpha(x_i) - \alpha(x_{i-1}) = \alpha'(\bar{x}_i)(\alpha(x_i) - \alpha(x_{i-1})), \quad (x_{i-1} < \bar{x}_i < x_i).$$

This implies

$$\sum_{i=1}^n \tilde{f}(\bar{x}_i)[\alpha(x_i) - \alpha(x_{i-1})] = \sum_{i=1}^n \tilde{f}(\bar{x}_i)\alpha'(\bar{x}_i)(\alpha(x_i) - \alpha(x_{i-1})).$$

That is

$$(FHS) \int_a^{+\infty} \tilde{f}(x) d\alpha = (FH) \int_a^{+\infty} \tilde{f}(x)\alpha'(x) dx.$$

The proof is complete.

**Theorem 3.4.** Let  $\alpha : [a, +\infty) \rightarrow \mathbb{R}$  be an increasing function such that  $\alpha \in C^1[a, +\infty)$ ,  $|\alpha'(x)| \leq M$  and  $\tilde{f} = \tilde{0}$  a.e. on  $[a, +\infty)$  (i.e.  $\tilde{f}(x) = \tilde{0}$  on  $[a, +\infty)$  except a Lebesgue zero measure set). Then  $(\tilde{f}, \alpha) \in FHS[a, +\infty)$  and  $(FHS) \int_a^{+\infty} \tilde{f}(x) d\alpha = \tilde{0}$ .

**Proof.** Since  $\alpha(x) \in C^1[a, +\infty)$ , and  $|\alpha'(x)| \leq M$  for all  $x \in [a, +\infty)$ . By Lagrange mean value theorem, there exists  $\xi \in [x_{i-1}, x_i]$  such that

$$\alpha(x_i) - \alpha(x_{i-1}) = \alpha'(\xi)(\alpha(x_i) - \alpha(x_{i-1})) \leq M(\alpha(x_i) - \alpha(x_{i-1})).$$

Let  $S = \{x | \tilde{f}(x) \neq \tilde{0}\}$  and for each positive integer  $n$ , set  $S = \bigcup S_n \subset [a, +\infty)$ , where  $S_n = \{x | n - 1 < D(\tilde{f}(x), \tilde{0}) \leq n\}$ ,  $n = 1, 2, 3, \dots$ . For every  $\epsilon > 0$  and a positive integer  $n$ , choose an open set  $G_n$  such that  $S_n \subset G_n$  and  $\mu(G_n) < \frac{\epsilon}{nM2^n}$ . Define  $\delta(x)$  on  $[a, +\infty)$  by

$$\delta(x) = \begin{cases} 1, & x \in [a, +\infty) \setminus S, \\ \delta(x), & \text{such that } (x - \delta(x), x + \delta(x)) \subset G_n, x \in S_n, n = 1, 2, \dots \end{cases}$$

For any  $\delta$ -fine division  $T = \{[x_{i-1}, x_i]; \xi_i\}$ , we have

$$\begin{aligned} D\left(\int_a^{+\infty} \tilde{f}(x) d\alpha, \tilde{0}\right) &= D\left(\sum_{\xi_i \in S} \tilde{f}(\xi_i)[\alpha(x_i) - \alpha(x_{i-1})] + \sum_{\xi_i \in [a, +\infty) \setminus S} \tilde{f}(\xi_i)[\alpha(x_i) - \alpha(x_{i-1})], \tilde{0}\right) \\ &= D\left(\sum_{\xi_i \in S} \tilde{f}(\xi_i)[\alpha(x_i) - \alpha(x_{i-1})], \tilde{0}\right) \leq \sum_{\xi_i \in S} D(\tilde{f}(\xi_i)[\alpha(x_i) - \alpha(x_{i-1})], \tilde{0}) \\ &\leq M \sum_{i=1}^{\infty} \sum_{\xi_i \in S_i} D(\tilde{f}(\xi_i), \tilde{0})(\alpha(x_i) - \alpha(x_{i-1})) < M \sum_{i=1}^{\infty} i \cdot \frac{\epsilon}{iM2^i} \\ &= \epsilon. \end{aligned}$$

The proof is complete.

**Remark 3.3.** Let  $\alpha : [a, +\infty) \rightarrow \mathbb{R}$  be an increasing function and  $\alpha \in C^1[a, +\infty)$ , and  $|\alpha'(x)| \leq M$ . If  $\tilde{f}(x) = \tilde{g}(x)$  a.e. on  $[a, +\infty)$  and  $(\tilde{f}, \alpha) \in FHS[a, +\infty)$ , then  $(\tilde{g}, \alpha) \in FHS[a, +\infty)$  and

$$(FHS) \int_a^{+\infty} \tilde{f}(x) d\alpha = (FHS) \int_a^{+\infty} \tilde{g}(x) d\alpha.$$

Using Theorem 3.4, naturally, we have the following conclusion.

**Theorem 3.5.** Let  $\alpha : [a, +\infty) \rightarrow \mathbb{R}$  be an increasing function. If  $\tilde{f}(x) = \tilde{0}$  a.e.s. on  $[a, +\infty)$  (i.e.  $\tilde{f}(x) = \tilde{0}$  on  $[a, +\infty)$  except a  $\alpha$ -Lebesgue-Stieltjes zero measure set), then  $(\tilde{f}, \alpha) \in FHS[a, +\infty)$  and  $(FHS) \int_a^{+\infty} \tilde{f}(x) d\alpha = \tilde{0}$ .



## 4 Conclusion

The aim of this paper is attempt to extend the theory of the fuzzy Henstock-Stieltjes integral on a infinite interval, we firstly define and discuss the Henstock-Stieltjes integral for fuzzy-number-valued functions on a infinite interval. On the other hand, the integrability of the fuzzy Henstock-Stieltjes integral on a infinite interval are also shown and discussed. In the future, we shall consider the continuity and the differentiability of the primitive for the fuzzy Henstock-Stieltjes integral on a infinite interval, the quadrature rules for the fuzzy Henstock-Stieltjes integral on a infinite interval, the convergence theorems for sequences of the fuzzy Henstock-Stieltjes integrable functions on a infinite interval, and so on.

## References

- [1] B. Bede, Note on “numerical solutions of fuzzy differential equations by predictor-corrector method” , *Information Sciences* 178 (2008) 1917-1922.
- [2] Z.T. Gong, Y.B. Shao, Global existence and uniqueness of solutions for fuzzy differential equations under dissipative-type conditions, *Computers and Mathematics with Applications* 56 (2008) 2716-2723.
- [3] O. Kaleva, Fuzzy differential equations, *Fuzzy Sets and Systems* 24 (1987) 301-317.
- [4] S. Nanda, On fuzzy integrals, *Fuzzy Sets and Systems* 32 (1989) 95-101.
- [5] C.X. Wu, Z.T. Gong, On Henstock integral of fuzzy-number-valued functions(I), *Fuzzy Sets and Systems* 120 (2001) 523-532.
- [6] B. Bede, S.G. Gal, Quadrature rules for integrals of fuzzy-number-valued functions, *Fuzzy Sets and Systems* 145 (2004) 359-380.
- [7] Z.T. Gong, L. Wang, The numerical calculus of expectations of fuzzy random variables, *Fuzzy Sets and Systems* 158 (2007) 722-738.
- [8] K. Kwak, W. Pedrycz, Face Recognition: A study in information fusion using fuzzy integral, *Pattren Recognition Letters* 26 (2005) 719-733.
- [9] H.C. Wu, The improper fuzzy Riemann integral and its numerical integration, *Information Sciences* 111 (1998) 109-137.
- [10] P. Billingsley, *Probability Measures*, John Wiley and Sons, Inc., New York, 1968.
- [11] S.S. Dragomir, The unified treatment of trapezoid, Simpson, and Ostrowski type inequality for monotonic mappings and applications, *Mathematical and Computer Modelling* 31 (2000) 61-70.
- [12] L. Egghe, Construction of concentration measures for general Lorenz curves using Riemann-Stieltjes integral, *Mathematical and Computer Modelling* 35 (2002) 1149-1163.
- [13] I. Štajner-Papuga, T. Grbić, M. Daňková, Pseudo-Riemann-Stieltjes integral, *Information Sciences* 179 (2009) 2923-2933.
- [14] F.A. Medvedev, *Development of the concept of integral*, Nauka, Moscow, 1974 (in Russian).
- [15] F. Riesz, B. Sz.-Nagy, *Functional Analysis*, Ungar, New York, 1955.
- [16] L.A. Zadeh, Probability measures of fuzzy events, *Journal of Mathematical Analysis and Applications* 23 (1968) 421-427.
- [17] H.C. Wu, The fuzzy Riemann-Stieltjes integral. *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems* 6 (1998) 51-67.
- [18] X.K. Ren, C.X. Wu, Z.G. Zhu, A new kind of fuzzy Riemann-Stieltjes integral, in: X.Z. Wang, D. Yeung, X.L. Wang (Eds.), *Proc. 5th Int. Conf. on Machine Learning and Cybernetics, ICMLC 2006*, Dalian, China, 2006, pp. 1885-1888.
- [19] X.K. Ren, *The non-additive measure and the fuzzy Riemann-Stieltjes integral*, Ph.D dissertation, Harbin Institute of Technology, 2008.
- [20] Z.T. Gong, L.L. Wang, The Henstock-Stieltjes Integral for Fuzzy-number-valued Functions, *Information Sciences* 188 (2012) 276-297

- [21] M.L. Puri, D.A. Ralescu, Fuzzy random variables, *Journal of Mathematics Analysis and Applications* 114 (1986) 409-422.
- [22] C.V. Negoita, D. Ralescu, *Applications of Fuzzy Sets to Systems Analysis*, Wiley, New York, 1975.
- [23] P.Y. Lee, *Lanzhou Lectures on Henstock Integration*, World Scientific, Singapore, 1989.
- [24] R. Henstock, *Theory of Integration*, Butterworth, London, 1963.

# New weighted $q$ -Čebyšev-Grüss type inequalities for double integrals

Zhen Liu<sup>1</sup> and Wengui Yang<sup>2\*</sup>

<sup>1</sup>*Department of Mathematics, Kashi Teacher's College, Kashi, Xinjiang 844000, China*

<sup>2</sup>*Department of Public Education, Sanmenxia Polytechnic, Sanmenxia, Henan 472000, China*

**Abstract:** In this paper, we establish the weighted double  $q$ -integrals Montgomery identity for functions of two independent variables, then obtain weighted  $q$ -Čebyšev-Grüss type inequalities for double integrals. Furthermore, weighted  $q$ -Ostrowski type inequalities for double integrals are also given.

**Keywords:** Čebyšev-Grüss type inequalities; Ostrowski type inequalities; Montgomery identity; double  $q$ -integrals

**2010 Mathematics Subject Classification:** 34A08; 26D10; 26D15.

## 1 Introduction and preliminaries

1882, P.L. Čebyšev [7] prove that, if  $f', g' \in L_\infty[a, b]$ , then

$$|T(f, g)| \leq \frac{1}{12}(b-a)^2 \|f'\|_\infty \|g'\|_\infty, \tag{1.1}$$

where for two functions  $f, g : [a, b] \rightarrow \mathbb{R}$ , the functional

$$T(f, g) = \frac{1}{b-a} \int_a^b f(x)g(x)dx - \left( \frac{1}{b-a} \int_a^b f(x)dx \right) \left( \frac{1}{b-a} \int_a^b g(x)dx \right), \tag{1.2}$$

and  $\|\cdot\|_\infty$  denotes the norm in  $L_\infty[a, b]$  defined as  $\|f\|_\infty = \operatorname{ess\,sup}_{t \in [a, b]} |f(t)|$ .

In 1935, G. Grüss [13] showed that

$$|T(f, g)| \leq \frac{1}{4}(M-m)(N-n), \tag{1.3}$$

provided  $m, M, n$  and  $N$  are real numbers satisfying the conditions,

$$m \leq f(x) \leq M, \quad n \leq g(x) \leq N, \tag{1.4}$$

for all  $x \in [a, b]$ , where  $T(f, g)$  is as defined by (1.2).

In 1938, Ostrowski [19] proved the following integral inequality:

Let  $f : I \rightarrow \mathbb{R}$ , where  $I \subset \mathbb{R}$  is an interval, be a mapping that is differentiable in the interior of  $I$  ( $\operatorname{Int}I$ ), and let  $a, b \in \operatorname{Int}I$ ,  $a < b$ . If  $|f'(t)| \leq M, \forall t \in (a, b)$ , then,

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t)dt \right| \leq \left[ \frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a)M, \tag{1.5}$$

for all  $x \in [a, b]$ .

During the past few years, many researchers have given considerable attention to the above results and various generalizations, extensions and variants of these inequalities (1.1), (1.3) and (1.5) have appeared in the literature, see [1, 2, 3, 6, 8, 9, 11, 12, 16, 17, 18, 20, 21, 22] and the references cited therein. Find new

\*Corresponding author.

Email: [lz790821ks@126.com](mailto:lz790821ks@126.com) (Z. Liu) and [wgyang0617@yahoo.com](mailto:wgyang0617@yahoo.com) (W. Yang)

inequalities in the multidimensional cases still an interesting problem. In [4, 10], the authors proved the double integrals Montgomery identity:

$$f(x, y) = \frac{1}{b-a} \int_a^b f(t, y) dt + \frac{1}{d-c} \int_c^d f(x, s) ds - \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(t, s) dt ds + \int_a^b \int_c^d P(x, t) Q(y, s) \frac{\partial^2 f(t, s)}{\partial t \partial s} dt ds, \quad (1.6)$$

where  $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$  is differentiable, the derivative  $\frac{\partial^2 f(t, s)}{\partial t \partial s}$  is integrable on  $[a, b] \times [c, d]$ , and the Peano kernels  $P(x, t)$  and  $Q(y, s)$  are defined by

$$P(x, t) = \begin{cases} \frac{t-a}{b-a}, & a \leq t \leq x, \\ \frac{t-b}{b-a}, & x \leq t \leq b \end{cases} \quad \text{and} \quad Q(y, s) = \begin{cases} \frac{s-c}{d-c}, & c \leq s \leq y, \\ \frac{s-d}{d-c}, & y \leq s \leq d. \end{cases}$$

Furthermore, Guezane-Lakoud and Aissaoui [14] established new extension of the weighted Montgomery identity (1.6) for functions of two independent variables, then obtained new Čebyšev type inequalities.

For the sake of convenience, some definitions and propositions are cited on  $q$ -integral as follows. Some details see [5, 15].

In what follows,  $q$  is a real number satisfying  $0 < q < 1$ .

**Definition 1.1** ([5]). For an arbitrary function  $f(x)$ , the  $q$ -differential is defined by  $(d_q f)(x) = f(qx) - f(x)$ . In particular,  $d_q x = (q - 1)x$ .  $q$ -derivative is defined by

$$(D_q f)(x) = \frac{d_q f(x)}{d_q x} = \frac{f(qx) - f(x)}{(q - 1)x}, \quad (D_q f)(0) = \lim_{x \rightarrow 0} (D_q f)(x).$$

Clearly, if  $f(x)$  is differentiable, then  $\lim_{q \rightarrow 1^-} (D_q f)(x) = \frac{df(x)}{dx}$ . And  $q$ -derivatives of higher order by

$$(D_q^0 f)(x) = f(x) \quad \text{and} \quad (D_q^n f)(x) = D_q(D_q^{n-1} f)(x), \quad n \in \mathbb{N}.$$

**Definition 1.2** ([5]). Suppose  $0 < a < b$ . The definite  $q$ -integral is defined as

$$(I_q f)(x) = \int_0^x f(t) d_q t = x(1 - q) \sum_{n=0}^{\infty} f(xq^n) q^n, \quad x \in [0, b]. \quad (1.7)$$

and

$$\int_a^b f(x) d_q x = \int_0^b f(x) d_q x - \int_0^a f(x) d_q x. \quad (1.8)$$

Similarly as done for derivatives, an operator  $I_q^n$  can be defined, namely,

$$(I_q^0 f)(x) = f(x) \quad \text{and} \quad (I_q^n f)(x) = I_q(I_q^{n-1} f)(x), \quad n \in \mathbb{N}.$$

The definite  $q$ -integral defined above is too general for our purpose of studying inequalities. For example, if  $f(x) \geq 0$ , it is not necessarily true that  $\int_a^b f(x) d_q x \geq 0$ .

From now on, we will use a special type of the definite  $q$ -integral, which we will call the restricted definite  $q$ -integral. Throughout all the paper, we will use the following notations:

$$c_j = bq^j, \quad \text{for } j \in \{0, 1, \dots, n\}, \quad a = c_n = bq^n.$$

**Definition 1.3** ([5]). Let  $0 < q < 1$ ,  $b > 0$ , and  $n \in \mathbb{Z}^+$ . The restricted  $q$ -integral is defined as  $\int_{bq^n}^b f(x) d_q x$ .

The following formula readily follows from (1.7) and (1.8):

$$\int_a^b f(x)d_q x = \int_{bq^n}^b f(x)d_q x = (1-q)b \sum_{j=0}^{n-1} q^j f(bq^j) = (1-q) \sum_{j=0}^{n-1} c_j f(c_j).$$

Note that the restricted integral  $\int_a^b f(x)d_q x$  is just a finite sum, so no questions about convergency arise. It is easy to check that

$$\int_a^b D_q f d_q x = f(b) - f(a).$$

Obviously, if  $f(x) \geq g(x)$  on  $[a, b]$ , then  $\int_a^b f(x)d_q x \geq \int_a^b g(x)d_q x$ . If  $0 < k < n$ , then

$$\int_a^b f(x)d_q x = \int_a^{c_k} f(x)d_q x + \int_{c_k}^b f(x)d_q x.$$

The following is the formula for the  $q$ -integration by parts:

$$\int_a^b f(x)(D_q g)(x)d_q x = [f(x)g(x)]_a^b - \int_a^b g(qx)(D_q f)(x)d_q x.$$

Cauciman [15] gave  $q$ -integral Grüss's inequality as follows: Assume that (1.4) holds, then

$$\left| \frac{1}{b-a} \int_a^b f(x)g(x)d_q x - \left( \frac{1}{b-a} \int_a^b f(x)d_q x \right) \left( \frac{1}{b-a} \int_a^b g(x)d_q x \right) \right| \leq \frac{1}{4}(M-m)(N-n).$$

Assume that  $w : [a, b] \rightarrow [0, \infty)$  satisfying  $\int_a^b w(x)d_q x = 1$ . Set  $W(t) = \int_a^t w(x)d_q x$  for  $t \in [a, b]$ ,  $W(t) = 0$  for  $t < a$ , and  $W(t) = 1$  for  $t > b$ . We give weighted  $q$ -integral Peano kernel  $P_w(x, t)$  defined by

$$P_w(x, t) = \begin{cases} W(t), & a \leq t \leq x, \\ W(t) - 1, & x \leq t \leq b. \end{cases}$$

Then the following weighted  $q$ -integral Montgomery identity holds: (see [23])

$$f(x) = \int_a^b w(t)f(qt)d_q t + \int_a^b P_w(x, t)(D_q f)(t)d_q t.$$

In 2011, Yang [23] obtained the following inequalities:

$$|T(w, f, g)| \leq \|D_q f\| \|D_q g\| \int_a^b w(x)H^2(qx)d_q x,$$

and

$$|T(w, f, g)| \leq \frac{1}{2} \int_a^b w(x)[|g(qx)| \|D_q f\| + |f(qx)| \|D_q g\|]H(qx)d_q x,$$

where  $\|\cdot\|$  as  $\|h\| = \sup_{t \in [a, b]} |h(t)|$  for  $h \in C[a, b]$ ,

$$T(w, f, g) = \int_a^b w(x)f(qx)g(qx)d_q x - \left( \int_a^b w(x)f(qx)d_q x \right) \left( \int_a^b w(x)g(qx)d_q x \right),$$

and

$$H(x) = \int_a^b |P_w(x, t)|d_q t$$

for all  $x \in [a, b]$ .

Motivated by the results mentioned above, by using weighted  $q$ -integral Montgomery identity for functions of two independent variables, we establish some new weighted  $q$ -Čebyšev type inequalities for double integrals. Furthermore, weighted  $q$ -Ostrowski type inequalities for double integrals are also given.

## 2 Weighted $q$ -Čebyšev type inequalities for double integrals

Assume that  $w : [a, b] \rightarrow \mathbb{R}_0 = [0, \infty)$  and  $u : [c, d] \rightarrow \mathbb{R}_0$  satisfying  $\int_a^b w(x)d_{q_1}x = \int_c^d u(y)d_{q_2}y = 1$ , where  $0 < q_1, q_2 < 1$ . Set  $W(t) = \int_a^t w(x)d_{q_1}x$  for  $t \in [a, b]$  and  $U(s) = \int_a^s u(y)d_{q_2}y$  for  $s \in [c, d]$ , so we have  $W(a) = U(c) = 0$  and  $W(b) = U(d) = 1$ . We give the following weighted  $q$ -integral Peano kernels  $P_w(x, t)$  and  $Q_u(y, s)$  defined by

$$P_w(x, t) = \begin{cases} W(t), & a \leq t \leq x, \\ W(t) - 1, & x \leq t \leq b \end{cases} \quad \text{and} \quad Q_u(y, s) = \begin{cases} U(s), & c \leq s \leq y, \\ U(s) - 1, & y \leq s \leq d. \end{cases} \quad (2.1)$$

We use the following notations to simplify details of the presentation. Let  $\frac{\partial f(t,s)}{d_{q_1}t}$  and  $\frac{\partial f(t,s)}{d_{q_2}s}$  be partial  $q$ -derivative on  $t$  and  $s$ , respectively. For some suitable functions  $w : [a, b] \rightarrow \mathbb{R}_0$ ,  $u : [c, d] \rightarrow \mathbb{R}_0$  and  $f, g : \Omega = [a, b] \times [c, d] \rightarrow \mathbb{R}$ , we set

$$\begin{aligned} T(w, u, f, g) &= \int_a^b \int_c^d w(x)u(y)f(q_1x, q_2y)g(q_1x, q_2y)d_{q_1}xd_{q_2}y \\ &\quad - \int_a^b \int_c^d w(x)u(y)g(q_1x, q_2y) \left( \int_a^b w(t)f(q_1t, q_2y)d_{q_1}t \right) d_{q_1}xd_{q_2}y \\ &\quad - \int_a^b \int_c^d w(x)u(y)g(q_1x, q_2y) \left( \int_c^d u(s)f(q_1x, q_2s)d_{q_2}s \right) d_{q_1}xd_{q_2}y \\ &\quad + \left( \int_a^b \int_c^d w(x)u(y)f(q_1x, q_2y)d_{q_1}xd_{q_2}y \right) \left( \int_a^b \int_c^d w(x)u(y)g(q_1x, q_2y)d_{q_1}xd_{q_2}y \right) \end{aligned}$$

and define  $\| \cdot \|$  as  $\|h\| = \sup_{(t,s) \in \Omega} |h(t, s)|$  for  $h \in C(\Omega, \mathbb{R})$ .

**Theorem 2.1.** *Let  $f : \Omega \rightarrow \mathbb{R}$ ,  $w : [a, b] \rightarrow \mathbb{R}_0$  and  $u : [c, d] \rightarrow \mathbb{R}_0$  satisfying  $\int_a^b w(x)d_{q_1}x = \int_c^d u(y)d_{q_2}y = 1$ , then*

$$\begin{aligned} f(x, y) &= \int_a^b w(t)f(q_1t, y)d_{q_1}t + \int_c^d u(s)f(x, q_2s)d_{q_2}s - \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s \\ &\quad + \int_a^b \int_c^d P_w(x, t)Q_u(y, s) \frac{\partial f(t,s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s, \quad (2.2) \end{aligned}$$

for  $(x, y) \in \Omega$ , where the weighted  $q$ -integral Peano kernels  $P_w(x, t)$  and  $Q_u(y, s)$  are defined by (2.1).

*Proof.* According to the weighted  $q$ -integral Peano kernels  $P_w(x, t)$  and  $Q_u(y, s)$  and the proof of Theorem 1 in [23], we obtain

$$\begin{aligned} &\int_a^b \int_c^d P_w(x, t)Q_u(y, s) \frac{\partial f(t,s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s = \int_a^b P_w(x, t) \left( \int_c^d Q_u(y, s) \frac{\partial f(t,s)}{d_{q_1}td_{q_2}s} d_{q_2}s \right) d_{q_1}t \\ &= \int_a^b P_w(x, t) \left( \frac{\partial f(t,y)}{d_{q_1}t} - \int_c^d u(s) \frac{\partial f(t, q_2s)}{d_{q_1}t} d_{q_2}s \right) d_{q_1}t \\ &= \int_a^b P_w(x, t) \frac{\partial f(t,y)}{d_{q_1}t} d_{q_1}t - \int_c^d u(s) \left( \int_a^b P_w(x, t) \frac{\partial f(t, q_2s)}{d_{q_1}t} d_{q_1}t \right) d_{q_2}s \\ &= \left( f(x, y) - \int_a^b w(t)f(q_1t, y)d_{q_1}t \right) - \int_c^d u(s) \left( f(x, q_2s) - \int_a^b w(t)f(q_1t, q_2s)d_{q_1}t \right) d_{q_2}s \\ &= f(x, y) - \int_a^b w(t)f(q_1t, y)d_{q_1}t - \int_c^d u(s)f(x, q_2s)d_{q_2}s + \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s. \end{aligned}$$

Thus we have

$$f(x, y) = \int_a^b w(t)f(q_1t, y)d_{q_1}t + \int_c^d u(s)f(t, q_2s)d_{q_2}s - \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s + \int_a^b \int_c^d P_w(x, t)Q_u(y, s) \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s}d_{q_1}td_{q_2}s,$$

and this completes the proof.  $\square$

**Theorem 2.2.** Let  $f, g : \Omega \rightarrow \mathbb{R}$ ,  $w : [a, b] \rightarrow \mathbb{R}_0$  and  $u : [c, d] \rightarrow \mathbb{R}_0$  satisfying  $\int_a^b w(x)d_{q_1}x = \int_c^d u(y)d_{q_2}y = 1$ , then

$$|T(w, u, f, g)| \leq \left\| \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} \right\| \left\| \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} \right\| \int_a^b \int_c^d w(x)u(y)H^2(q_1x, q_2y)d_{q_1}xd_{q_2}y, \tag{2.3}$$

where

$$H(x, y) = \int_a^b \int_c^d |P_w(x, t)Q_u(y, s)|d_{q_1}td_{q_2}s.$$

*Proof.* Since the functions  $f$  and  $g$  satisfy the hypothesis of Theorem 2.1, the following identities hold:

$$f(x, y) = \int_a^b w(t)f(q_1t, y)d_{q_1}t + \int_c^d u(s)f(t, q_2s)d_{q_2}s - \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s + \int_a^b \int_c^d P_w(x, t)Q_u(y, s) \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s}d_{q_1}td_{q_2}s, \tag{2.4}$$

and

$$g(x, y) = \int_a^b w(t)g(q_1t, y)d_{q_1}t + \int_c^d u(s)g(x, q_2s)d_{q_2}s - \int_a^b \int_c^d w(t)u(s)g(q_1t, q_2s)d_{q_1}td_{q_2}s + \int_a^b \int_c^d P_w(x, t)Q_u(y, s) \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s}d_{q_1}td_{q_2}s. \tag{2.5}$$

Due to the above two inequalities (2.4) and (2.5), we have

$$f(q_1x, q_2y) = \int_a^b w(t)f(q_1t, q_2y)d_{q_1}t + \int_c^d u(s)f(q_1x, q_2s)d_{q_2}s - \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s + \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s}d_{q_1}td_{q_2}s, \tag{2.6}$$

and

$$g(q_1x, q_2y) = \int_a^b w(t)g(q_1t, q_2y)d_{q_1}t + \int_c^d u(s)g(q_1x, q_2s)d_{q_2}s - \int_a^b \int_c^d w(t)u(s)g(q_1t, q_2s)d_{q_1}td_{q_2}s + \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s}d_{q_1}td_{q_2}s. \tag{2.7}$$

Multiplying (2.6) by (2.7), we obtain

$$\begin{aligned} & \left( f(q_1x, q_2y) - \int_a^b w(t)f(q_1t, q_2y)d_{q_1}t - \int_c^d u(s)f(q_1x, q_2s)d_{q_2}s + \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s \right) \\ & \times \left( g(q_1x, q_2y) - \int_a^b w(t)g(q_1t, q_2y)d_{q_1}t - \int_c^d u(s)g(q_1x, q_2s)d_{q_2}s + \int_a^b \int_c^d w(t)u(s)g(q_1t, q_2s)d_{q_1}td_{q_2}s \right) \\ & = \left( \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s}d_{q_1}td_{q_2}s \right) \left( \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s}d_{q_1}td_{q_2}s \right). \end{aligned}$$

Consequently,

$$\begin{aligned}
 & f(q_1x, q_2y)g(q_1x, q_2y) - f(q_1x, q_2y) \int_a^b w(t)g(q_1t, q_2y)d_{q_1}t - f(q_1x, q_2y) \int_c^d u(s)g(q_1x, q_2s)d_{q_2}s \\
 & \quad + f(q_1x, q_2y) \int_a^b \int_c^d w(t)u(s)g(q_1t, q_2s)d_{q_1}td_{q_2}s - g(q_1x, q_2y) \int_a^b w(t)f(q_1t, q_2y)d_{q_1}t \\
 & \quad + \int_a^b w(t)f(q_1t, q_2y)d_{q_1}t \int_a^b w(t)g(q_1t, q_2y)d_{q_1}t + \int_a^b w(t)f(q_1t, q_2y)d_{q_1}t \int_c^d u(s)g(q_1x, q_2s)d_{q_2}s \\
 & \quad - \int_a^b w(t)f(q_1t, q_2y)d_{q_1}t \int_a^b \int_c^d w(t)u(s)g(q_1t, q_2s)d_{q_1}td_{q_2}s - g(q_1x, q_2y) \int_c^d u(s)f(q_1x, q_2s)d_{q_2}s \\
 & \quad + \int_c^d u(s)f(q_1x, q_2s)d_{q_2}s \int_a^b w(t)g(q_1t, q_2y)d_{q_1}t + \int_c^d u(s)f(q_1x, q_2s)d_{q_2}s \int_c^d u(s)g(q_1x, q_2s)d_{q_2}s \\
 & - \int_c^d u(s)f(q_1x, q_2s)d_{q_2}s \int_a^b \int_c^d w(t)u(s)g(q_1t, q_2s)d_{q_1}td_{q_2}s + g(q_1x, q_2y) \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s \\
 & \quad - \int_a^b w(t)g(q_1t, q_2y)d_{q_1}t \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s - \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s \\
 & \quad \times \int_c^d u(s)g(q_1x, q_2s)d_{q_2}s + \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s \int_a^b \int_c^d w(t)u(s)g(q_1t, q_2s)d_{q_1}td_{q_2}s \\
 & = \left( \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s \right) \left( \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s \right) \quad (2.8)
 \end{aligned}$$

Multiplying both sides of (2.8) by  $w(x)u(y)$ , then  $q$ -integrating the resultant identity over  $\Omega$ , we get

$$\begin{aligned}
 T(w, u, f, g) &= \int_a^b \int_c^d w(x)u(y) \left[ \left( \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s \right) \right. \\
 & \quad \left. \times \left( \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s \right) \right] d_{q_1}xd_{q_2}y.
 \end{aligned}$$

Finally, using the properties of modulus we observe that

$$\begin{aligned}
 |T(w, u, f, g)| &\leq \left\| \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} \right\| \left\| \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} \right\| \int_a^b \int_c^d w(x)u(y) \left[ \left( \int_a^b \int_c^d |P_w(q_1x, t)Q_u(q_2y, s)|d_{q_1}td_{q_2}s \right) \right. \\
 & \quad \left. \times \left( \int_a^b \int_c^d |P_w(q_1x, t)Q_u(q_2y, s)|d_{q_1}td_{q_2}s \right) \right] d_{q_1}xd_{q_2}y \\
 &= \left\| \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} \right\| \left\| \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} \right\| \int_a^b \int_c^d w(x)u(y)H^2(q_1x, q_2y)d_{q_1}xd_{q_2}y.
 \end{aligned}$$

This completes the proof of Theorem 2.2. □

**Theorem 2.3.** Let  $f, g : \Omega \rightarrow \mathbb{R}$ ,  $w : [a, b] \rightarrow \mathbb{R}_0$  and  $u : [c, d] \rightarrow \mathbb{R}_0$  satisfying  $\int_a^b w(x)d_{q_1}x = \int_c^d u(y)d_{q_2}y = 1$ , then

$$\begin{aligned}
 |T(w, u, f, g)| &\leq \frac{1}{2} \int_a^b \int_c^d w(x)u(y) \\
 & \quad \times \left( |g(q_1x, q_2y)| \left\| \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} \right\| + |f(q_1x, q_2y)| \left\| \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} \right\| \right) H(q_1x, q_2y)d_{q_1}xd_{q_2}y, \quad (2.9)
 \end{aligned}$$

where  $H(x, y)$  is defined in Theorem 2.2.



*Proof.* Multiplying both sides of (2.6) and (2.7) by  $w(x)u(y)g(q_1x, q_2y)$  and  $w(x)u(y)f(q_1x, q_2y)$ , adding the resulting identities and rewriting, we have

$$\begin{aligned}
 & w(x)u(y)f(q_1x, q_2y)g(q_1x, q_2y) \\
 &= \frac{1}{2} \left( w(x)u(y)g(q_1x, q_2y) \int_a^b w(t)f(q_1t, q_2y)d_{q_1}t + w(x)u(y)g(q_1x, q_2y) \int_c^d u(s)f(q_1x, q_2s)d_{q_2}s \right. \\
 &\quad - w(x)u(y)g(q_1x, q_2y) \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s + w(x)u(y)f(q_1x, q_2y) \int_a^b w(t)g(q_1t, q_2y)d_{q_1}t \\
 &\quad \left. + w(x)u(y)f(q_1x, q_2y) \int_c^d u(s)g(q_1x, q_2s)d_{q_2}s - w(x)u(y)f(q_1x, q_2y) \int_a^b \int_c^d w(t)u(s)g(q_1t, q_2s)d_{q_1}td_{q_2}s \right) \\
 &\quad + \frac{1}{2} \left( w(x)u(y)g(q_1x, q_2y) \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s \right. \\
 &\quad \left. + w(x)u(y)f(q_1x, q_2y) \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s \right). \quad (2.10)
 \end{aligned}$$

Q-integrating both sides of (2.10) with respect to  $x$  from  $a$  to  $b$  and  $y$  from  $c$  to  $d$  and rewriting we have

$$\begin{aligned}
 T(w, u, f, g) &= \frac{1}{2} \left( \int_a^b \int_c^d w(x)u(y)g(q_1x, q_2y) \left( \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s \right) d_{q_1}xd_{q_2}y \right. \\
 &\quad \left. + \int_a^b \int_c^d w(x)u(y)f(q_1x, q_2y) \left( \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s \right) d_{q_1}xd_{q_2}y \right). \quad (2.11)
 \end{aligned}$$

Finally, from (2.11) and using the properties of modulus we observe that

$$\begin{aligned}
 T(w, u, f, g) &\leq \frac{1}{2} \left( \int_a^b \int_c^d w(x)u(y)|g(q_1x, q_2y)| \left( \int_a^b \int_c^d |P_w(q_1x, t)Q_u(q_2y, s)| \left| \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} \right| d_{q_1}td_{q_2}s \right) d_{q_1}xd_{q_2}y \right. \\
 &\quad \left. + \int_a^b \int_c^d w(x)u(y)|f(q_1x, q_2y)| \left( \int_a^b \int_c^d |P_w(q_1x, t)Q_u(q_2y, s)| \left| \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} \right| d_{q_1}td_{q_2}s \right) d_{q_1}xd_{q_2}y \right) \\
 &\leq \frac{1}{2} \int_a^b \int_c^d w(x)u(y) \left( |g(q_1x, q_2y)| \left\| \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} \right\| + |f(q_1x, q_2y)| \left\| \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} \right\| \right) \\
 &\quad \times \left( \int_a^b \int_c^d |P_w(q_1x, t)Q_u(q_2y, s)| d_{q_1}td_{q_2}s \right) d_{q_1}xd_{q_2}y \\
 &= \frac{1}{2} \int_a^b \int_c^d w(x)u(y) \left( |g(q_1x, q_2y)| \left\| \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} \right\| + |f(q_1x, q_2y)| \left\| \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} \right\| \right) H(q_1x, q_2y) d_{q_1}xd_{q_2}y.
 \end{aligned}$$

This completes the proof of Theorem 2.3. □

**Remark 2.4.** If  $q_1, q_2 \rightarrow 1^-$ , by Definitions 1.2 and 1.2, the partial  $q$ -derivative and double  $q$ -integrals are the usual partial derivative and double integrals, so Theorems 2.2 and 2.3 are reduced to Theorems 3 and 4 in [14].

**Theorem 2.5.** Let  $f, g : \Omega \rightarrow \mathbb{R}$ ,  $w : [a, b] \rightarrow \mathbb{R}_0$  and  $u : [c, d] \rightarrow \mathbb{R}_0$  satisfying  $\int_a^b w(x)d_{q_1}x = \int_c^d u(y)d_{q_2}y = 1$ , then

$$|T(w, u, f, g)| \leq \|g(q_1x, q_2y)\| \left\| \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} \right\| \int_a^b \int_c^d w(x)u(y)H(q_1x, q_2y)d_{q_1}xd_{q_2}y, \quad (2.12)$$

and

$$|T(w, u, f, g)| \leq \|f(q_1x, q_2y)\| \left\| \frac{\partial g(t, s)}{d_{q_1}t d_{q_2}s} \right\| \int_a^b \int_c^d w(x)u(y)H(q_1x, q_2y)d_{q_1}x d_{q_2}y, \quad (2.13)$$

where  $H(x, y)$  is defined in Theorem 2.2.

*Proof.* We prove only (2.12), since the proof of (2.13) is similar. The identity (2.6) shows that

$$\begin{aligned} f(q_1x, q_2y) &= \int_a^b w(t)f(q_1t, q_2y)d_{q_1}t + \int_c^d u(s)f(q_1x, q_2s)d_{q_2}s - \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}t d_{q_2}s \\ &\quad + \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial f(t, s)}{d_{q_1}t d_{q_2}s} d_{q_1}t d_{q_2}s, \end{aligned} \quad (2.14)$$

for  $(x, y) \in \Omega$ .

Now, if we multiply (2.14) by  $w(x)u(y)g(q_1x, q_2y)$  and  $q$ -integrate over  $(x, y) \in \Omega$ , we deduce

$$\begin{aligned} &\int_a^b \int_c^d w(x)u(y)f(q_1x, q_2y)g(q_1x, q_2y)d_{q_1}x d_{q_2}y \\ &= \int_a^b \int_c^d w(x)u(y)g(q_1x, q_2y) \left( \int_a^b w(t)f(q_1t, q_2y)d_{q_1}t \right) d_{q_1}x d_{q_2}y \\ &\quad + \int_a^b \int_c^d w(x)u(y)g(q_1x, q_2y) \left( \int_c^d u(s)f(q_1x, q_2s)d_{q_2}s \right) d_{q_1}x d_{q_2}y \\ &\quad - \int_a^b \int_c^d w(x)u(y)g(q_1x, q_2y)d_{q_1}x d_{q_2}y \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}t d_{q_2}s \\ &\quad + \int_a^b \int_c^d w(x)u(y)g(q_1x, q_2y) \left( \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial f(t, s)}{d_{q_1}t d_{q_2}s} d_{q_1}t d_{q_2}s \right) d_{q_1}x d_{q_2}y, \end{aligned}$$

which provides another representation for the functional  $T(w, u, f, g)$  namely,

$$T(w, u, f, g) = \int_a^b \int_c^d w(x)u(y)g(q_1x, q_2y) \left( \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial f(t, s)}{d_{q_1}t d_{q_2}s} d_{q_1}t d_{q_2}s \right) d_{q_1}x d_{q_2}y, \quad (2.15)$$

From (2.15) and using modules properties, it yields

$$\begin{aligned} |T(w, u, f, g)| &\leq \int_a^b \int_c^d w(x)u(y)|g(q_1x, q_2y)| \left( \int_a^b \int_c^d |P_w(q_1x, t)Q_u(q_2y, s)| \left| \frac{\partial f(t, s)}{d_{q_1}t d_{q_2}s} \right| d_{q_1}t d_{q_2}s \right) d_{q_1}x d_{q_2}y \\ &\leq \|g(q_1x, q_2y)\| \left\| \frac{\partial f(t, s)}{d_{q_1}t d_{q_2}s} \right\| \int_a^b \int_c^d w(x)u(y) \left( \int_a^b \int_c^d |P_w(q_1x, t)Q_u(q_2y, s)| d_{q_1}t d_{q_2}s \right) d_{q_1}x d_{q_2}y \\ &= \|g(q_1x, q_2y)\| \left\| \frac{\partial f(t, s)}{d_{q_1}t d_{q_2}s} \right\| \int_a^b \int_c^d w(x)u(y)H(q_1x, q_2y)d_{q_1}x d_{q_2}y. \end{aligned}$$

This completes the proof of Theorem 2.5. □

### 3 Weighted $q$ -Ostrowski type inequalities for double integrals

For some given functions  $f, g : \Omega \rightarrow \mathbb{R}$  and  $w : [a, b] \rightarrow \mathbb{R}_0$  and  $u : [c, d] \rightarrow \mathbb{R}_0$  satisfying  $\int_a^b w(x)d_{q_1}x = \int_c^d u(y)d_{q_2}y = 1$ ,

$$S(w, u, f, g) = f(q_1x, q_2y)g(q_1x, q_2y) - \frac{1}{2} \left( f(q_1x, q_2y) \int_a^b w(t)g(q_1t, q_2y)d_{q_1}t + \int_c^d u(s)g(q_1x, q_2s)d_{q_2}s \right. \\ \times f(q_1x, q_2y) + g(q_1x, q_2y) \int_a^b w(t)f(q_1t, q_2y)d_{q_1}t + g(q_1x, q_2y) \int_c^d u(s)f(q_1x, q_2s)d_{q_2}s \\ \left. - f(q_1x, q_2y) \int_a^b \int_c^d w(t)u(s)g(q_1t, q_2s)d_{q_1}td_{q_2}s - g(q_1x, q_2y) \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s \right).$$

**Theorem 3.1.** Let  $f, g : \Omega \rightarrow \mathbb{R}$ ,  $w : [a, b] \rightarrow \mathbb{R}_0$  and  $u : [c, d] \rightarrow \mathbb{R}_0$  satisfying  $\int_a^b w(x)d_{q_1}x = \int_c^d u(y)d_{q_2}y = 1$ , then

$$|S(w, u, f, g)| \leq \frac{1}{2} \left( |f(q_1x, q_2y)| \left\| \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} \right\| + |g(q_1x, q_2y)| \left\| \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} \right\| \right) H(q_1x, q_2y), \tag{3.1}$$

and

$$|T(w, u, f, g)| \leq \frac{1}{2} \left( |f(q_1x, q_2y)| \left\| \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} \right\| + |g(q_1x, q_2y)| \left\| \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} \right\| \right) \int_a^b \int_c^d w(x)u(y)H(q_1x, q_2y)d_{q_1}xd_{q_2}y. \tag{3.2}$$

where  $H(x, y)$  is defined in Theorem 2.2.

*Proof.* Multiplying both sides of (2.6) and (2.7) by  $g(q_1x, q_2y)$  and  $f(q_1x, q_2y)$ , adding the resulting identities and rewriting we have

$$f(q_1x, q_2y)g(q_1x, q_2y) = \frac{1}{2} \left( f(q_1x, q_2y) \int_a^b w(t)g(q_1t, q_2y)d_{q_1}t + f(q_1x, q_2y) \int_c^d u(s)g(q_1x, q_2s)d_{q_2}s \right. \\ + g(q_1x, q_2y) \int_a^b w(t)f(q_1t, q_2y)d_{q_1}t + g(q_1x, q_2y) \int_c^d u(s)f(q_1x, q_2s)d_{q_2}s \\ \left. - f(q_1x, q_2y) \int_a^b \int_c^d w(t)u(s)g(q_1t, q_2s)d_{q_1}td_{q_2}s - g(q_1x, q_2y) \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s \right) \\ + \frac{1}{2} \left( f(q_1x, q_2y) \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s \right. \\ \left. + g(q_1x, q_2y) \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s \right), \tag{3.3}$$

which implies

$$S(w, u, f, g) = \frac{1}{2} \left( f(q_1x, q_2y) \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s \right. \\ \left. + g(q_1x, q_2y) \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s \right).$$

We observe

$$|S(w, u, f, g)| \leq \frac{1}{2} \left( |f(q_1x, q_2y)| \left\| \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} \right\| + |g(q_1x, q_2y)| \left\| \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} \right\| \right) H(q_1x, q_2y).$$

Multiplying both sides of (3.3) by  $w(x)u(y)$  and  $q$ -integrate over  $(x, y) \in \Omega$ , we deduce

$$T(w, u, f, g) = \frac{1}{2} \left( \int_a^b \int_c^d w(x)u(y)f(q_1x, q_2y) \left( \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s \right) d_{q_1}xd_{q_2}y \right. \\ \left. + \int_a^b \int_c^d w(x)u(y)g(q_1x, q_2y) \left( \int_a^b \int_c^d P_w(q_1x, t)Q_u(q_2y, s) \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} d_{q_1}td_{q_2}s \right) d_{q_1}xd_{q_2}y \right).$$

We observe

$$|T(w, u, f, g)| \leq \frac{1}{2} \left( |f(q_1x, q_2y)| \left\| \frac{\partial g(t, s)}{d_{q_1}td_{q_2}s} \right\| + |g(q_1x, q_2y)| \left\| \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} \right\| \right) \int_a^b \int_c^d w(x)u(y)H(q_1x, q_2y)d_{q_1}xd_{q_2}y.$$

This completes the proof of Theorem 3.1. □

Let  $g(x, y) = 1$ , we have the following corollary.

**Corollary 3.2.** *Let  $f : \Omega \rightarrow \mathbb{R}$ ,  $w : [a, b] \rightarrow \mathbb{R}_0$  and  $u : [c, d] \rightarrow \mathbb{R}_0$  satisfying  $\int_a^b w(x)d_{q_1}x = \int_c^d u(y)d_{q_2}y = 1$ , then*

$$\left| f(x, y) - \int_a^b w(t)f(q_1t, y)d_{q_1}t - \int_c^d u(s)f(x, q_2s)d_{q_2}s + \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s \right| \leq H(x, y) \left\| \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} \right\|,$$

where  $H(x, y)$  is defined in Theorem 2.2, and especially, let  $w(x) = \frac{1}{b-a}$  and  $u(y) = \frac{1}{d-c}$ , we get

$$\left| f(x, y) - \int_a^b w(t)f(q_1t, y)d_{q_1}t - \int_c^d u(s)f(x, q_2s)d_{q_2}s + \int_a^b \int_c^d w(t)u(s)f(q_1t, q_2s)d_{q_1}td_{q_2}s \right| \leq \frac{1}{4(q_1+1)(q_2+1)} \left( (b-a)^2 + 4 \left( x - \frac{a+b}{2} \right)^2 \right) \left( (d-c)^2 + 4 \left( y - \frac{c+d}{2} \right)^2 \right) \left\| \frac{\partial f(t, s)}{d_{q_1}td_{q_2}s} \right\|. \quad (3.4)$$

**Remark 3.3.** If  $q_1, q_2 \rightarrow 1^-$ , the inequality (3.4) are reduced to the main result in [4].

## References

- [1] F. AHMAD, N. S. BARNETT, S. S. DRAGOMIR, *New weighted Ostrowski and Čebyšev type inequalities*, *Nonlinear Analysis*, **71**, 12 (2009), 1408-1412.
- [2] F. AHMAD, P. CERONE, S. S. DRAGOMIR AND N. A. MIR, *On some bounds of Ostrowski and Čebyšev type*, *Journal of Mathematical Inequalities*, **4**, 1 (2010), 53-65.
- [3] M. ALOMARI, M. DARUS, S. S. DRAGOMIR, AND P. CERONE, *Ostrowski type inequalities for functions whose derivatives are  $s$ -convex in the second sense*, *Applied Mathematics Letters*, **23**, 9 (2010), 1071-1076.
- [4] N. S. BARNETT, S. S. DRAGOMIR, *An Ostrowski type inequality for double integrals and applications for cubature formulae*, *Soochow Journal of Mathematics*, **27**, 1 (2001), 1-10.
- [5] H. CAUCIMAN, *Integral inequalities in  $q$ -calculus*, *Computers & Mathematics with Applications*, **47**, 2-3 (2004), 281-300.
- [6] K. BOUKERRIOUA, A. GUEZANE-LAKOUD, *On generalization of Čebyšev type inequalities*, *Journal of Inequalities in Pure and Applied Mathematics*, **8**, 2 (2007), Art. 15.

- [7] P. L. ČEBYŠEV, *Sur les expressions approximatives des intégrales définies par les autres prises entre les mêmes limites*, Proceedings of the Mathematical Society of Kharkov, **2** (1882), 93-98.
- [8] P. CERONE AND S. S. DRAGOMIR, *Some new Ostrowski-type bounds for the Čebyšev functional and applications*, Journal of Mathematical Inequalities, **8**, 1 (2014), 159-170.
- [9] S. S. DRAGOMIR AND N. S. BARNETT, *An Ostrowski type inequality for mapping whose second derivatives are bounded and applications*, Journal of the Indian Mathematical Society (N.S.), **66**, 1-4 (1999), 237-245.
- [10] S. S. DRAGOMIR, P. CERONE, N. S. BARNETT, J. ROUMELIOTIS, *An inequality of the Ostrowski type for double integrals and applications for cubature formulae*, Tamsui Oxford Journal of Mathematical Sciences, **16**, 1 (2000), 1-16.
- [11] S. S. DRAGOMIR AND A. SOFO, *An inequality for monotonic functions generalizing Ostrowski and related results*, Computers & Mathematics with Applications, **51**, 3-4(2006), 497-506.
- [12] S. S. DRAGOMIR AND S. WANG, *An inequality of Ostrowski-Grüss type and its applications to the estimation of error bounds for some special means and for some numerical quadrature rules*, Computers Mathematics with Applications, **33** (1997), 15-20.
- [13] G. GRÜSS, *Über das maximum des absoluten Betrages von  $\frac{1}{b-a} \int_a^b f(x)g(x)dx - \frac{1}{(b-a)^2} \int_a^b f(x)dx \times \int_a^b g(x)dx$* , Mathematische Zeitschrift, **39**, 1 (1935), 215-226.
- [14] A. GUEZANE-LAKOUD AND F. AISSAOUI, *New Čebyšev type inequalities for double integrals*, Journal of Mathematical Inequalities, **5**, 4 (2011), 453-462.
- [15] V. KAC AND P. CHEUNG, *Quantum Calculus*, Springer Verlag, 2002.
- [16] Z. LIU, *A sharp general Ostrowski type inequality for double integrals*, Tamsui Oxford Journal of Information and Mathematical Sciences, **28**, 2 (2012), 217-226.
- [17] Z. LIU, *A variant of Chebyshev inequality with applications*, Journal of Mathematical Inequalities, **7**, 4 (2013), 551-561.
- [18] M. MATIĆ, J. E. PEČARIĆ AND N. UJEVIC, *On new estimation of the remainder in generalized Taylor's formula*, Mathematical Inequalities & Applications, **3**, 2 (1999), 343-361.
- [19] A. OSTROWSKI, *Über die Absolutabweichung einer differentüerbaren Funktion von ihrem Integralmittelwert*, Commentarii Mathematici Helvetici, **10**, 1 (1937), 226-227.
- [20] B. G. PACHPATTE, *On Čebyšev-Grüss type inequalities via Pečarić's extension of the montgomery identity*, Journal of Inequalities in Pure and Applied Mathematics, **7**, 1 (2006), Art. 11.
- [21] B. G. PACHPATTE, *New inequalities of Čebyšev type for double integrals*, Demonstratio Mathematica, **11**, 1 (2007), 43-50.
- [22] K. L. TSENG, S. R. HWANG, AND S. S. DRAGOMIR, *Generalizations of weighted Ostrowski type inequalities for mappings of bounded variation and their applications*, Computers & Mathematics with Applications, **55**, 8 (2008), 1785-1793.
- [23] W. YANG, *On weighted  $q$ -Čebyšev-Grüss type inequalities*, Computers & Mathematics with Applications, **61**, 5 (2011), 1342-1347.

## QUADRATIC $\rho$ -FUNCTIONAL INEQUALITIES IN NORMED SPACES

IKAN CHOI, SUNGHOON KIM\*, GEORGE A. ANASTASSIOU, AND CHOONKIL PARK\*

ABSTRACT. In this paper, we solve the quadratic  $\rho$ -functional inequalities

$$\begin{aligned} & \|f(x+y) + f(x-y) - 2f(x) - 2f(y)\| \\ & \leq \left\| \rho \left( 2f\left(\frac{x+y}{2}\right) + 2f\left(\frac{x-y}{2}\right) - f(x) - f(y) \right) \right\|, \end{aligned} \tag{0.1}$$

where  $\rho$  is a number with  $|\rho| < 1$  and

$$\begin{aligned} & \left\| 2f\left(\frac{x+y}{2}\right) + 2f\left(\frac{x-y}{2}\right) - f(x) - f(y) \right\| \\ & \leq \|\rho(f(x+y) + f(x-y) - 2f(x) - 2f(y))\|, \end{aligned} \tag{0.2}$$

where  $\rho$  is a number with  $|\rho| < \frac{1}{2}$ . Using the fixed point method, we prove the Hyers-Ulam stability of the quadratic  $\rho$ -functional inequalities (0.1) and (0.2) in normed spaces.

### 1. INTRODUCTION AND PRELIMINARIES

The stability problem of functional equations originated from a question of Ulam [29] concerning the stability of group homomorphisms.

The functional equation

$$f(x+y) = f(x) + f(y)$$

is called the *Cauchy equation*. In particular, every solution of the Cauchy equation is said to be an *additive mapping*. Hyers [14] gave a first affirmative partial answer to the question of Ulam for Banach spaces. Hyers' Theorem was generalized by Aoki [2] for additive mappings and by Rassias [20] for linear mappings by considering an unbounded Cauchy difference. A generalization of the Rassias theorem was obtained by Găvruta [11] by replacing the unbounded Cauchy difference by a general control function in the spirit of Rassias' approach.

The functional equation

$$f(x+y) + f(x-y) = 2f(x) + 2f(y)$$

is called a *quadratic functional equation*. In particular, every solution of the quadratic functional equation is said to be a *quadratic mapping*. A Hyers-Ulam stability problem for the quadratic functional equation was proved by Skof [23] for mappings  $f : X \rightarrow Y$ , where  $X$  is a normed space and  $Y$  is a Banach space. Cholewa [6] noticed that the theorem of Skof is still true if the relevant domain  $X$  is replaced by an Abelian group. Czerwik [7] proved the Hyers-Ulam stability of the quadratic functional equation. The functional equation  $f\left(\frac{x+y}{2}\right) = \frac{1}{2}f(x) + \frac{1}{2}f(y)$  is called the *Jensen type quadratic functional equation*. The stability problems of several functional equations

---

2010 *Mathematics Subject Classification*. Primary 39B52, 47H10, 39B72.

*Key words and phrases*. Hyers-Ulam stability; quadratic  $\rho$ -functional inequality; fixed point.

\*Corresponding authors.

have been extensively investigated by a number of authors and there are many interesting results concerning this problem (see [1, 3, 16, 17, 21, 22, 25, 26, 27, 28, 30, 31]).

In [12], Gilányi showed that if  $f$  satisfies the functional inequality

$$\|2f(x) + 2f(y) - f(xy^{-1})\| \leq \|f(xy)\| \tag{1.1}$$

then  $f$  satisfies the Jordan-von Neumann functional equation

$$2f(x) + 2f(y) = f(xy) + f(xy^{-1}).$$

See also [10, 24]. Gilányi [13] and Fechner [9] proved the Hyers-Ulam stability of the functional inequality (1.1). Park, Cho and Han [18] proved the Hyers-Ulam stability of additive functional inequalities.

**Lemma 1.1.** (Banach fixed-point theorem) *Let  $(S, d)$  be a complete metric space and let  $T : S \rightarrow S$  be a strictly contractive mapping with Lipschitz constant  $\alpha < 1$ . Then for each given element  $x \in S$ , there exists a positive integer  $n_0$  such that*

- (1)  $d(T^n x, T^{n+1} x) < \infty, \quad \forall n \geq n_0;$
- (2) *the sequence  $\{T^n x\}$  converges to a fixed point  $y^*$  of  $T$ ;*
- (3)  *$y^*$  is the unique fixed point of  $T$  in the set  $Y = \{y \in S \mid d(T^{n_0} x, y) < \infty\}$ ;*
- (4)  $d(y, y^*) \leq \frac{1}{1-\alpha} d(y, Ty)$  for all  $y \in Y$ .

Since we defined the metric  $d$  as generalized metric in order to use this lemma in the proof of the problem we extend the lemma.

**Lemma 1.2.** ([8]) *Let  $(S, d)$  be a complete generalized metric space and let  $J : S \rightarrow S$  be a strictly contractive mapping with Lipschitz constant  $\alpha < 1$ . Then for each given element  $x \in S$ , either*

$$d(J^n x, J^{n+1} x) = \infty$$

*for all nonnegative integers  $n$  or there exists a positive integer  $n_0$  such that*

- (1)  $d(J^n x, J^{n+1} x) < \infty, \quad \forall n \geq n_0;$
- (2) *the sequence  $\{J^n x\}$  converges to a fixed point  $y^*$  of  $J$ ;*
- (3)  *$y^*$  is the unique fixed point of  $J$  in the set  $Y = \{y \in S \mid d(J^{n_0} x, y) < \infty\}$ ;*
- (4)  $d(y, y^*) \leq \frac{1}{1-\alpha} d(y, Jy)$  for all  $y \in Y$ .

In 1996, Isac and Rassias [15] were the first to provide applications of stability theory of functional equations for the proof of new fixed point theorems with applications. By using fixed point methods, the stability problems of several functional equations have been extensively investigated by a number of authors (see [4, 5, 19]).

In Section 2, we solve the quadratic  $\rho$ -functional inequality (0.1) and prove the Hyers-Ulam stability of the quadratic  $\rho$ -functional inequality (0.1) in normed spaces.

In Section 3, we solve the quadratic  $\rho$ -functional inequality (0.2) and prove the Hyers-Ulam stability of the quadratic  $\rho$ -functional inequality (0.2) in normed spaces.

Throughout this paper, assume that  $X$  is a normed space and  $Y$  is a Banach space.

QUADRATIC  $\rho$ -FUNCTIONAL INEQUALITIES

2. HYERS-ULAM STABILITY OF THE QUADRATIC  $\rho$ -FUNCTIONAL INEQUALITY (0.1): A FIXED POINT APPROACH

In this section, assume that  $|\rho| < 1$ .

We solve the quadratic  $\rho$ -functional inequality (0.1) in normed spaces.

**Lemma 2.1.** *A mapping  $f : X \rightarrow Y$  satisfies*

$$\begin{aligned} & \|f(x+y) + f(x-y) - 2f(x) - 2f(y)\| \\ & \leq \left\| \rho \left( 2f\left(\frac{x+y}{2}\right) + 2f\left(\frac{x-y}{2}\right) - f(x) - f(y) \right) \right\| \end{aligned} \tag{2.1}$$

for all  $x, y \in X$  if and only if  $f : X \rightarrow Y$  is quadratic.

*Proof.* Assume that  $f : X \rightarrow Y$  satisfies (2.1).

Letting  $x = y = 0$  in (2.1), we get  $\|2f(0)\| \leq |\rho|\|2f(0)\|$ . So  $f(0) = 0$ .

Letting  $y = x$  in (2.1), we get  $\|f(2x) - 4f(x)\| \leq 0$  and so  $f(2x) = 4f(x)$  for all  $x \in X$ . Thus

$$f\left(\frac{x}{2}\right) = \frac{1}{4}f(x) \tag{2.2}$$

for all  $x \in X$ .

It follows from (2.1) and (2.2) that

$$\begin{aligned} & \|f(x+y) + f(x-y) - 2f(x) - 2f(y)\| \\ & \leq \left\| \rho \left( 2f\left(\frac{x+y}{2}\right) + 2f\left(\frac{x-y}{2}\right) - f(x) - f(y) \right) \right\| \\ & = \frac{|\rho|}{2} \|f(x+y) + f(x-y) - 2f(x) - 2f(y)\| \end{aligned}$$

and so

$$f(x+y) + f(x-y) = 2f(x) + 2f(y)$$

for all  $x, y \in X$ .

The converse is obviously true. □

We prove the Hyers-Ulam stability of the quadratic  $\rho$ -functional inequality (0.1) in Banach spaces.

**Theorem 2.2.** *Let  $\varphi : X^2 \rightarrow [0, \infty)$  be a function such that there exists an  $\alpha < 1$  with*

$$\varphi(a, b) \leq 4\alpha\varphi\left(\frac{a}{2}, \frac{b}{2}\right) \tag{2.3}$$

for all  $a, b \in X$ . Let  $f : X \rightarrow Y$  be a mapping satisfying

$$\begin{aligned} & \|f(x+y) + f(x-y) - 2f(x) - 2f(y)\| \\ & \leq \left\| \rho \left( 2f\left(\frac{x+y}{2}\right) + 2f\left(\frac{x-y}{2}\right) - f(x) - f(y) \right) \right\| + \varphi(x, y) \end{aligned} \tag{2.4}$$

for all  $x, y \in X$ . Then there exists a unique quadratic mapping  $Q : X \rightarrow Y$  such that

$$\|f(x) - Q(x)\| \leq \frac{1}{4 - 4\alpha} \varphi(x, x)$$

for all  $x \in X$ .



*Proof.* Consider the set

$$S := \{h : X \rightarrow Y\}$$

and let  $d$  be the generalized metric on  $S$ :

$$d(g, h) := \inf\{\mu \in \mathbb{R}_+ : \|g - h\| \leq \mu\varphi(x, x), x \in S\}$$

It is easy to show that  $(S, d)$  is complete. Let  $J$  be the linear mapping from  $S$  to  $S$  such that

$$Jg(x) := \frac{1}{4}g(2x) \tag{2.5}$$

Let  $g, h \in S$  be given such that  $d(g, h) = \varepsilon$ . Then from (2.3) and (2.5), we get

$$\|Jg(x) - Jh(x)\| = \left\| \frac{1}{4}g(2x) - \frac{1}{4}h(2x) \right\| \leq \frac{1}{4}\varepsilon\varphi(2x, 2x) \leq \alpha\varepsilon\varphi(x, x)$$

This means  $d(Jg, Jh) \leq \alpha d(g, h)$ .

So the function  $J : S \rightarrow S$  is a contractive mapping such that

$$d(Jg, Jh) \leq \alpha d(g, h)$$

for  $0 \leq \alpha < 1$ .

Letting  $y = x$  in (2.4), we get

$$\|f(2x) - 4f(x)\| \leq \varphi(x, x)$$

and so

$$\|f(x) - Jf(x)\| \leq \frac{1}{4}\varphi(x, x)$$

for all  $x \in X$ . Thus we get  $d(f, Jf) \leq \frac{1}{4}$ .

By Lemma 1.2, there exists a mapping  $Q : X \rightarrow Y$  satisfying the following:

(1)  $Q$  is a fixed point of  $J$ , i.e.,

$$Q(2a) = 4Q(a) \tag{2.6}$$

for all  $a \in X$ . The mapping  $Q$  is a unique fixed point of  $J$  in the set

$$M = \{g \in S : d(f, g) < \infty\}.$$

This implies that  $Q$  is a unique mapping satisfying (2.6) such that there exists a  $\mu \in (0, \infty)$  satisfying

$$\|f(a) - Q(a)\| \leq \mu\varphi(a, a)$$

for all  $a \in X$ ;

(2)  $d(J^l f, Q) \rightarrow 0$  as  $l \rightarrow \infty$ . This implies the equality

$$\lim_{l \rightarrow \infty} \frac{1}{4^l} f(2^l a) = Q(a)$$

for all  $a \in X$ ;

(3)  $d(f, Q) \leq \frac{1}{1-\alpha}d(f, Jf)$ , which implies the inequality

$$d(f, Q) \leq \frac{1}{4-4\alpha}.$$

QUADRATIC  $\rho$ -FUNCTIONAL INEQUALITIES

So

$$\|f(a) - Q(a)\| \leq \frac{1}{4 - 4\alpha} \varphi(a, a)$$

for all  $a \in X$ .

Then

$$\begin{aligned} & \|Q(x + y) + Q(x - y) - 2Q(x) - 2Q(y)\| \\ &= \lim_{l \rightarrow \infty} \left\| \frac{1}{4^l} (f(2^l(x + y)) + f(2^l(x - y)) - 2f(2^l x) - 2f(2^l y)) \right\| \\ &\leq \lim_{l \rightarrow \infty} \left\| \frac{\rho}{4^l} (2f(2^{l-1}(x + y)) + 2f(2^{l-1}(x - y)) - f(2^l x) - f(2^l y)) \right\| \\ &\quad + \lim_{l \rightarrow \infty} \frac{1}{4^l} \varphi(2^l x, 2^l y) \\ &= \left\| \rho \left( 2Q\left(\frac{x + y}{2}\right) + 2Q\left(\frac{x - y}{2}\right) - Q(x) - Q(y) \right) \right\| \end{aligned}$$

for all  $x, y \in X$ . Hence

$$Q(x + y) + Q(x - y) = 2Q(x) + 2Q(y)$$

for all  $x, y$ . So  $Q : X \rightarrow Y$  is quadratic. □

**Remark 2.3.** We could prove the same statement with the same manner in spite of replacing the condition  $\varphi(a, b) \leq 4\alpha\varphi\left(\frac{a}{2}, \frac{b}{2}\right)$  into  $\varphi(a, b) \leq \frac{1}{4}\alpha\varphi(2a, 2b)$  by defining  $J$  such that  $Jg(x) = 4g\left(\frac{x}{2}\right)$  instead of  $Jg(x) = \frac{1}{4}g(2x)$ . It could be also applied to Theorem 3.2.

**Corollary 2.4.** Let  $r \neq 2$  and  $\theta$  be nonnegative real numbers, and let  $f : X \rightarrow Y$  be a mapping such that

$$\begin{aligned} & \|f(x + y) + f(x - y) - 2f(x) - 2f(y)\| \\ &\leq \left\| \rho \left( 2f\left(\frac{x + y}{2}\right) + 2f\left(\frac{x - y}{2}\right) - f(x) - f(y) \right) \right\| + \theta(\|x\|^r + \|y\|^r) \end{aligned}$$

for all  $x, y \in X$ . Then there exists a unique quadratic mapping  $Q : X \rightarrow Y$  such that

$$\|f(x) - Q(x)\| \leq \frac{2\theta}{|4 - 2^r|} \|x\|^r$$

for all  $x \in X$ .

3. HYERS-ULAM STABILITY OF THE QUADRATIC  $\rho$ -FUNCTIONAL INEQUALITY (0.2): A FIXED POINT APPROACH

In this section, assume that  $|\rho| < \frac{1}{2}$ .

We solve the quadratic  $\rho$ -functional inequality (0.2) in normed spaces.

**Lemma 3.1.** A mapping  $f : X \rightarrow Y$  satisfies

$$\begin{aligned} & \left\| 2f\left(\frac{x + y}{2}\right) + 2f\left(\frac{x - y}{2}\right) - f(x) - f(y) \right\| \\ &\leq \|\rho(f(x + y) + f(x - y) - 2f(x) - 2f(y))\| \end{aligned} \tag{3.1}$$

for all  $x, y \in X$  if and only if  $f : X \rightarrow Y$  is quadratic.

*Proof.* Assume that  $f : X \rightarrow Y$  satisfies (3.1).

Letting  $x = y = 0$  in (3.1), we get  $\|2f(0)\| \leq |\rho|\|2f(0)\|$ . So  $f(0) = 0$ .

Letting  $y = 0$  in (3.1), we get

$$\left\| 4f\left(\frac{x}{2}\right) - f(x) \right\| \leq 0 \tag{3.2}$$

and so  $f\left(\frac{x}{2}\right) = \frac{1}{4}f(x)$  for all  $x \in X$ .

It follows from (3.1) and (3.2) that

$$\begin{aligned} & \frac{1}{2} \|f(x+y) + f(x-y) - 2f(x) - 2f(y)\| \\ &= \left\| 2f\left(\frac{x+y}{2}\right) + 2f\left(\frac{x-y}{2}\right) - f(x) - f(y) \right\| \\ &\leq |\rho| \|f(x+y) + f(x-y) - 2f(x) - 2f(y)\| \end{aligned}$$

and so

$$f(x+y) + f(x-y) = 2f(x) + 2f(y)$$

for all  $x, y \in X$ .

The converse is obviously true. □

We prove the Hyers-Ulam stability of the quadratic  $\rho$ -functional inequality (0.2) in Banach spaces.

**Theorem 3.2.** *Let  $\varphi : X^2 \rightarrow [0, \infty)$  be a function such that there exists an  $\alpha < 1$  with*

$$\varphi(a, b) \leq 4\alpha\varphi\left(\frac{a}{2}, \frac{b}{2}\right)$$

for all  $a, b \in X$ . Let  $f : X \rightarrow Y$  be a mapping satisfying  $f(0) = 0$  and

$$\begin{aligned} & \left\| 2f\left(\frac{x+y}{2}\right) + 2f\left(\frac{x-y}{2}\right) - f(x) - f(y) \right\| \\ & \leq |\rho| \|f(x+y) + f(x-y) - 2f(x) - 2f(y)\| + \varphi(x, y) \end{aligned}$$

for all  $x, y \in X$ . Then there exists a unique quadratic mapping  $Q : X \rightarrow Y$  such that

$$\|f(x) - Q(x)\| \leq \frac{1}{4 - 4\alpha} \varphi(x, 0)$$

for all  $x \in X$ .

*Proof.* Consider the set

$$S := \{h : X \rightarrow Y\}$$

and let  $d$  be the generalized metric on  $S$ :

$$d(g, h) := \inf\{\mu \in \mathbb{R}_+ : \|g - h\| \leq \mu\varphi(x, 0), x \in S\}$$

It is easy to show that  $(S, d)$  is complete. Let  $J$  be the linear mapping from  $S$  to  $S$  such that

$$Jg(x) := \frac{1}{4}g(2x)$$

QUADRATIC  $\rho$ -FUNCTIONAL INEQUALITIES

Let  $Q : X \rightarrow Y$  be defined as in the proof of Theorem 2.2. Then

$$\begin{aligned} & \left\| 2Q\left(\frac{x+y}{2}\right) + 2Q\left(\frac{x-y}{2}\right) - Q(x) - Q(y) \right\| \\ &= \lim_{l \rightarrow \infty} \left\| \frac{1}{4^l} (2f(2^{l-1}(x+y)) + 2f(2^{l-1}(x-y)) - f(2^l x) - f(2^l y)) \right\| \\ &\leq \lim_{l \rightarrow \infty} \left\| \frac{\rho}{4^l} (f(2^l(x+y)) + f(2^l(x-y)) - 2f(2^l x) - 2f(2^l y)) \right\| \\ &\quad + \lim_{l \rightarrow \infty} \frac{1}{4^l} \varphi(2^l x, 2^l y) \\ &= \|\rho(Q(x+y) + Q(x-y) - 2Q(x) - 2Q(y))\| \end{aligned}$$

for all  $x, y \in X$ . Hence

$$Q(x+y) + Q(x-y) = 2Q(x) + 2Q(y)$$

for all  $x, y$ . So  $Q : X \rightarrow Y$  is quadratic. □

**Corollary 3.3.** *Let  $r \neq 2$  and  $\theta$  be nonnegative real numbers, and let  $f : X \rightarrow Y$  be a mapping satisfying  $f(0) = 0$  and*

$$\begin{aligned} & \left\| 2f\left(\frac{x+y}{2}\right) + 2f\left(\frac{x-y}{2}\right) - f(x) - f(y) \right\| \\ & \leq \|\rho(f(x+y) + f(x-y) - 2f(x) - 2f(y))\| + \theta(\|x\|^r + \|y\|^r) \end{aligned}$$

for all  $x, y \in X$ . Then there exists a unique quadratic mapping  $Q : X \rightarrow Y$  such that

$$\|f(x) - Q(x)\| \leq \frac{\theta}{|4 - 2^r|} \|x\|^r$$

for all  $x \in X$ .

ACKNOWLEDGMENTS

I. Choi and S. Kim were supported by the Seoul Science High School *R&E/I&D* Program in 2014. C. Park was supported by Basic Science Research Program through the National Research Foundation of Korea funded by the Ministry of Education, Science and Technology (NRF-2012R1A1A2004299).

REFERENCES

- [1] M. Adam, *On the stability of some quadratic functional equation*, J. Nonlinear Sci. Appl. **4** (2011), 50–59.
- [2] T. Aoki, *On the stability of the linear transformation in Banach spaces*, J. Math. Soc. Japan **2** (1950), 64–66.
- [3] L. Cădariu, L. Găvruta and P. Găvruta, *On the stability of an affine functional equation*, J. Nonlinear Sci. Appl. **6** (2013), 60–67.
- [4] L. Cădariu and V. Radu, *On the stability of the Cauchy functional equation: a fixed point approach*, Grazer Math. Ber. **346** (2004), 43–52.
- [5] L. Cădariu and V. Radu, *Fixed point methods for the generalized stability of functional equations in a single variable*, Fixed Point Theory and Applications **2008**, Art. ID 749392 (2008).
- [6] P.W. Cholewa, *Remarks on the stability of functional equations*, Aequationes Math. **27** (1984), 76–86.
- [7] S. Czerwik, *On the stability of the quadratic mapping in normed spaces*, Abh. Math. Sem. Univ. Hamburg **62** (1992), 59–64.
- [8] J. Diaz and B. Margolis, *A fixed point theorem of the alternative for contractions on a generalized complete metric space*, Bull. Amer. Math. Soc. **74** (1968), 305–309.
- [9] W. Fechner, *Stability of a functional inequalities associated with the Jordan-von Neumann functional equation*, Aequationes Math. **71** (2006), 149–161.

I. CHOI, S. KIM, G.A. ANASTASSIOU, AND C. PARK

- [10] W. Fechner, *On some functional inequalities related to the logarithmic mean*, Acta Math. Hungar. **128** (2010), 31–45.
- [11] P. Găvruta, *A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings*, J. Math. Anal. Appl. **184** (1994), 431–43.
- [12] A. Gilányi, *Eine zur Parallelogrammgleichung äquivalente Ungleichung*, Aequationes Math. **62** (2001), 303–309.
- [13] A. Gilányi, *On a problem by K. Nikodem*, Math. Inequal. Appl. **5** (2002), 707–710.
- [14] D. H. Hyers, *On the stability of the linear functional equation*, Proc. Natl. Acad. Sci. U.S.A. **27** (1941), 222–224.
- [15] G. Isac and Th.M. Rassias, *Stability of  $\psi$ -additive mappings: Applications to nonlinear analysis*, Internat. J. Math. Math. Sci. **19** (1996), 219–228.
- [16] S. Jung, *Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis*, Hadronic Press Inc., Palm Harbor, Florida, 2001.
- [17] C. Park, *Homomorphisms between Poisson  $JC^*$ -algebras*, Bull. Braz. Math. Soc. **36** (2005), 79–97.
- [18] C. Park, Y. Cho and M. Han, *Functional inequalities associated with Jordan-von Neumann-type additive functional equations*, J. Inequal. Appl. **2007** (2007), Article ID 41820, 13 pages.
- [19] V. Radu, *The fixed point alternative and the stability of functional equations*, Fixed Point Theory **4** (2003), 91–96.
- [20] Th. M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc. **72** (1978), 297–300.
- [21] Th.M. Rassias (ed.), *Functional Equations and Inequalities*, Kluwer Academic, Dordrecht, 2000.
- [22] S. Schin, D. Ki, J. Chang and M. Kim, *Random stability of quadratic functional equations: a fixed point approach*, J. Nonlinear Sci. Appl. **4** (2011), 37–49.
- [23] F. Skof, *Proprietà locali e approssimazione di operatori*, Rend. Sem. Mat. Fis. Milano **53** (1983), 113–129.
- [24] J. Rätz, *On inequalities associated with the Jordan-von Neumann functional equation*, Aequationes Math. **66** (2003), 191–200.
- [25] S. Shagholi, M. Eshaghi Gordji and M. Bavand Savadkouhi, *Stability of ternary quadratic derivations on ternary Banach algebras*, J. Comput. Anal. Appl. **13** (2011), 1097–1105.
- [26] S. Shagholi, M. Eshaghi Gordji and M. Bavand Savadkouhi, *Nearly ternary cubic homomorphisms in ternary Fréchet algebras*, J. Comput. Anal. Appl. **13** (2011), 1106–1114.
- [27] D. Shin, C. Park and Sh. Farhadabadi, *On the superstability of ternary Jordan  $C^*$ -homomorphisms*, J. Comput. Anal. Appl. **16** (2014), 964–973.
- [28] D. Shin, C. Park and Sh. Farhadabadi, *Stability and superstability of  $J^*$ -homomorphisms and  $J^*$ -derivations for a generalized Cauchy-Jensen equation*, J. Comput. Anal. Appl. **17** (2014), 125–134.
- [29] S. M. Ulam, *A Collection of the Mathematical Problems*, Interscience Publ. New York, 1960.
- [30] C. Zaharia, *On the probabilistic stability of the monomial functional equation*, J. Nonlinear Sci. Appl. **6** (2013), 51–59.
- [31] S. Zolfaghari, *Approximation of mixed type functional equations in  $p$ -Banach spaces*, J. Nonlinear Sci. Appl. **3** (2010), 110–122.

IKAN CHOI, SUNGHOON KIM

MATHEMATICS BRANCH, SEOUL SCIENCE HIGH SCHOOL, SEOUL 110-530, KOREA

*E-mail address:* dlrgrks623@naver.com; askwhy10@naver.com

GEORGE A. ANASTASSIOU

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF MEMPHIS, MEMPHIS, TN 38152, USA

*E-mail address:* ganastss@memphis.edu

CHOONKIL PARK

RESEARCH INSTITUTE FOR NATURAL SCIENCES, HANYANG UNIVERSITY, SEOUL 133-791, KOREA

*E-mail address:* baak@hanyang.ac.kr

# Solution of the Ulam stability problem for quartic $(a, b)$ -functional equations

Abdullah Alotaibi<sup>1</sup>, John Michael Rassias<sup>2</sup> and S.A. Mohiuddine<sup>1</sup>

<sup>1</sup>Department of Mathematics, Faculty of Science, King Abdulaziz University,  
P.O. Box 80203, Jeddah 21589, Saudi Arabia

<sup>2</sup>National and Capodistrian University of Athens, Pedagogical Department, Mathematics and Informatics, 4, Agamemnonos Str., Aghia Paraskevi, Attikis 15342, Greece  
Email: mathker11@hotmail.com<sup>1</sup>; jrassias@primedu.uoa.gr<sup>2</sup>; mohiuddine@gmail.com<sup>1</sup>

**Abstract.** The “oldest quartic” functional equation was introduced and solved by the author of this paper (see: Glas. Mat. Ser. III 34 (54) (1999), no. 2, 243-252) which is of the form:

$$f(x + 2y) + f(x - 2y) = 4[f(x + y) + f(x - y)] - 6f(x) + 24f(y).$$

Interesting results have been achieved by S.A. Mohiuddine et al., since 2009. In this paper, we are introducing new quartic functional equations, and establish fundamental formulas for the general solution of such functional equations and for “Ulam stability” of pertinent quartic functional inequalities.

*Keywords and phrases:* Quartic functional equations and inequalities; Various normed spaces; Ulam stability.

*AMS subject classification (2000):* 39B.

## 1. INTRODUCTION

In 1940 S. M. Ulam [1] gave a talk before the Mathematics Club of the University of Wisconsin in which he discussed a number of unsolved problems. Among these was the following famous “*stability Ulam question*”:

*We are given a group  $G$  and a metric group  $G'$  with metric  $\rho(.,.)$ . Given  $\epsilon > 0$ , does there exist a  $\delta > 0$  such that if  $f : G \rightarrow G'$  satisfies  $\rho(f(xy), f(x)f(y)) < \delta$  for all  $x, y$  in  $G$ , then a homomorphism  $h : G \rightarrow G'$  exists with  $\rho(f(x), h(x)) < \epsilon$  for all  $x \in G$ ?*

By now an affirmative answer has been given in several cases, and some interesting variations of the problem have also been investigated. We shall call such an  $f : G \rightarrow G'$  an *approximate homomorphism*.

In 1941 D. H. Hyers [2] considered the case of approximately additive mappings  $f : E \rightarrow E'$  where  $E$  and  $E'$  are Banach spaces and  $f$  satisfies the following *Hyers' inequality*

$$\|f(x + y) - f(x) - f(y)\| \leq \epsilon$$

for all  $x, y \in E$ . It was shown that the limit

$$L(x) = \lim_{n \rightarrow \infty} 2^{-n} f(2^n x)$$

exists for all  $x \in E$  and that  $L : E \rightarrow E'$  is the unique additive mapping satisfying  $\|f(x) - L(x)\| \leq \epsilon$ .

No continuity conditions are required for this result, but if  $f(tx)$  is continuous in the real variable  $t$  for each fixed  $x$ , then  $L$  is linear, and if  $f$  is continuous at a single point of  $E$  then  $L : E \rightarrow E'$  is also continuous.

In 1982-1994, a generalization of this result was proved by the author J. M. Rassias [3–7], as follows. He introduced the following *weaker condition (or weaker inequality or the generalized Cauchy inequality)*

$$\|f(x + y) - [f(x) + f(y)]\| \leq \theta \|x\|^p \|y\|^q$$

for all  $x, y$  in  $E$ , controlled by (or involving) a product of different powers of norms, where  $\theta \geq 0$  and real  $p, q : r = p + q \neq 1$ , and retained the condition of continuity of  $f(tx)$  in  $t$  for fixed  $x$ . Besides he investigated that it is possible to replace  $\epsilon$  in the above *Hyers' inequality*, by a non-negative real-valued function such that the pertinent series converges and other conditions hold and still obtain stability results. In all the cases investigated in these results, the approach to the existence question was to prove

asymptotic type formulas:  $L(x) = \lim_{n \rightarrow \infty} 2^{-n} f(2^n x)$ ;  $L(x) = \lim_{n \rightarrow \infty} 2^n f(2^{-n} x)$ .

**Theorem** (J. M. Rassias:1982-1994). Let  $X$  be a real normed linear space and let  $Y$  be a real complete normed linear space. Assume in addition that  $f : X \rightarrow Y$  is an approximately additive mapping for which there exist constants  $\theta \geq 0$  and  $p, q \in \mathbb{R}$  such that  $r = p + q \neq 1$  and  $f$  satisfies the “generalized Cauchy inequality”

$$\|f(x + y) - [f(x) + f(y)]\| \leq \theta \|x\|^p \|y\|^q$$

for all  $x, y \in X$ . Then there exists a unique additive mapping  $L : X \rightarrow Y$  satisfying

$$\|f(x) - L(x)\| \leq \frac{\theta}{|2^r - 2|} \|x\|^r$$

for all  $x \in X$ . If in addition  $f : X \rightarrow Y$  is a mapping such that the transformation  $t \rightarrow f(tx)$  is continuous in  $t \in \mathbb{R}$  for each fixed  $x \in X$ , then  $L$  is an  $\mathbb{R}$ -linear mapping.

In 1940, Ulam, S. M. [1] proposed the “Ulam stability problem”: *When does a linear transformation near an “approximately linear” transformation exist?* Since then, many specialists on this “famous Ulam problem”, have investigated interesting functional equations, for instance: D. H. Hyers [2], in 1941; T. Aoki [8], in 1950; T. M. Rassias [9], in 1978; Z. Gajda [10], in 1991; T. M. Rassias and P. Šemrl [11], in 1992; P. Găvruta [12], in 1994; S.-M. Jung [13], in 1998; K. W. Jun and H. M. Kim [14], in 2002; R. P. Agarwal et al. [15], in 2003, and others. Interesting Ulam-Hyers stability results have been established by S. A. Mohiuddine et al. ([16–19]). The “oldest quartic” functional equation was introduced and solved by the author of this paper, [20], which is of the form:

$$f(x + 2y) + f(x - 2y) = 4[f(x + y) + f(x - y)] - 6f(x) + 24f(y).$$

Since then various quartic equations have been proposed and solved by a number of experts in the area of functional equations and inequalities. For more details on these concepts, one can be referred to [21–30]. For further research in various normed spaces, we are introducing new quartic functional equations, and establish fundamental formulas for the general solution of such functional equations and for “Ulam stability” of pertinent quartic functional inequalities.

## 2. ON $(a, b)$ -QUARTIC FUNCTIONAL EQUATIONS

### 1. Stability of General $a$ -Quartic Functional Equation

$$2[f(ax + y) + f(x + ay)] + a(a - 1)^2 f(x - y) = 2(a^2 - 1)^2 [f(x) + f(y)] + a(a + 1)^2 f(x + y) \quad (1.1)$$

where  $a \neq 0, a \neq \pm 1$ .

Replacing  $x = y = 0$  in (1.1) one gets  $4a^2(1 - a^2)f(0) = 0$ , or

$$f(0) = 0. \quad (1.2)$$

Similarly, substituting  $x = x, y = 0$  in (1.1), we obtain

$$f(ax) = a^4 f(x) + (a^2 - 1)^2 f(0) = a^4 f(x) \quad (1.3)$$

Also assuming  $f(2x) = 16f(x)$ , replacing  $x = x, y = x$  in (1.1), and setting  $k = a + 1 \neq 0, \pm 1$ , one obtains

$$4f(kx) + a(a - 1)^2 f(0) = 4(a^2 - 1)^2 f(x) + a(a + 1)^2 f(2x), \quad (1.4)$$

or

$$f(kx) + \frac{1}{4}a(a - 1)^2 f(0) = (a^2 - 1)^2 f(x) + 4a(a + 1)^2 f(x) = k^4 f(x), \text{ or}$$

$$f(kx) = k^4 f(x) - \frac{1}{4}a(a - 1)^2 f(0) = k^4 f(x). \quad (1.5)$$

Without assuming  $f(2x) = 16f(x)$ , replacing  $x = x, y = -x$  in (1.1), and setting  $l = a - 1 \neq 0, \pm 1$ , one obtains

$$2[f(lx) + f(-lx)] + a(a - 1)^2 f(2x) = 2(a^2 - 1)^2 [f(x) + f(-x)] + a(a + 1)^2 f(0) \tag{1.6}$$

Placing  $-x$  on  $x$  in (1.6), and then subtracting the new equation from (1.6), we get  $a(a - 1)^2 [f(2x) - f(-2x)] = 0$ . Letting  $x/2$  on  $x$ , we find that  $f$  is an “*even function*”, such that  $f(-x) = f(x)$ . Thus from (1.6), we obtain

$$4f(lx) + a(a - 1)^2 f(2x) = 4(a^2 - 1)^2 f(x) + a(a + 1)^2 f(0). \tag{1.7}$$

Assuming  $f(2x) = 16f(x)$ , we get

$$f(lx) = l^4 f(x) + \frac{1}{4} a(a + 1)^2 f(0) = l^4 f(x).$$

Without assuming  $f(2x) = 16f(x)$ , subtracting (1.7) from (1.4), we obtain from (1.2) that

$$f(kx) - f(lx) = \frac{1}{2} a(a^2 + 1) f(2x) = \left[ \left(\frac{k}{2}\right)^4 - \left(\frac{l}{2}\right)^4 \right] f(2x).$$

Replacing  $x \rightarrow x/2$ , we get that

$$“f\left(\frac{k}{2}x\right) = \left(\frac{k}{2}\right)^4 f(x) \text{ if and only if } f\left(\frac{l}{2}x\right) = \left(\frac{l}{2}\right)^4 f(x)” \tag{1.8}$$

Employing the “quartic mean”, we have equivalently that

$$“\bar{f}_{\frac{k}{2}}(x) = \frac{f\left(\frac{k}{2}x\right)}{\left(\frac{k}{2}\right)^4} = f(x) \text{ iff } \bar{f}_{\frac{l}{2}}(x) = \frac{f\left(\frac{l}{2}x\right)}{\left(\frac{l}{2}\right)^4} = f(x)”.$$

Let  $X$  be a real normed linear space and let  $Y$  be a real complete normed linear space. Assume  $f : X \rightarrow Y$ , satisfying the following general  $a$ -quartic functional inequality

$$\|2[f(ax + y) + f(x + ay)] + a(a - 1)^2 f(x - y) - 2(a^2 - 1)^2 [f(x) + f(y)] - a(a + 1)^2 f(x + y)\| \leq c \tag{1.9}$$

where  $a \neq 0, a \neq \pm 1$ . Replacing  $x = y = 0$  in (1.9), one gets

$$\|f(0)\| \leq c/4a^2 |a^2 - 1|. \tag{1.10}$$

Substituting  $x = x, y = 0$  in (1.9), and employing the triangle inequality, we obtain:

$$\|f(ax) - a^4 f(x)\| \leq c_2 = \frac{c_1}{2} = \frac{2a^2 + |a^2 - 1|}{4a^2} c.$$

Note that

$$c_1 = \frac{2a^2 + |a^2 - 1|}{2a^2} c = \begin{cases} \frac{3a^2 - 1}{2a^2} c & \text{if } |a| > 1 \\ \frac{a^2 + 1}{2a^2} c & \text{if } |a| < 1; a \neq 0. \end{cases}$$

Therefore

$$\|f(ax) - a^4 f(x)\| \leq c_2 = \frac{c_1}{2} = \frac{3a^2 - 1}{4a^2} c, \text{ if } |a| > 1,$$

and

$$\|f(ax) - a^4 f(x)\| \leq c_3 = \frac{c_1}{2} = \frac{a^2 + 1}{4a^2} c, \text{ if } |a| < 1; a \neq 0.$$

Therefore we get

$$\|f(x) - a^{-4} f(ax)\| \leq a^{-4} c_2 = \frac{3a^2 - 1}{4a^6} c, \text{ if } |a| > 1,$$

and

$$\|f(x) - a^4 f(a^{-1}x)\| \leq c_3 = \frac{a^2 + 1}{4a^2} c, \text{ if } |a| < 1; a \neq 0.$$



Thus we can easily obtain the following general inequality

$$\|f(x) - a^{-4n}f(a^n x)\| \leq \frac{3a^2 - 1}{4a^2(a^4 - 1)}(1 - a^{-4n})c, \text{ if } |a| > 1. \tag{1.11}$$

In fact,

$$\begin{aligned} \|f(x) - a^{-4n}f(a^n x)\| &\leq \|f(x) - a^{-4}f(a^1 x)\| + a^{-4}\|f(ax) - a^{-4}f(a^2 x)\| + \dots \\ &\quad \dots + a^{-4(n-2)}\|f(a^{n-2} x) - a^{-4}f(a^{n-1} x)\| \\ &\quad + a^{-4(n-1)}\|f(a^{n-1} x) - a^{-4}f(a^n x)\| \tag{1.12} \\ &\leq \left(1 + a^{-4} + a^{-8} + \dots + a^{-4(n-2)} + a^{-4(n-1)}\right) \frac{3a^2 - 1}{4a^6}c \\ &\leq \frac{3a^2 - 1}{4a^2(a^4 - 1)}(1 - a^{-4n})c, \text{ if } |a| > 1. \end{aligned}$$

Also,

$$\begin{aligned} \|f(x) - \alpha^{4n}f(\alpha^{-n} x)\| &\leq \|f(x) - \alpha^4 f(\alpha^{-1} x)\| + \alpha^4 \|f(\alpha^{-1} x) - \alpha^4 f(\alpha^{-2} x)\| \\ &\quad + \dots + \alpha^{4(n-1)} \|f(\alpha^{-(n-1)} x) - \alpha^4 f(\alpha^{-n} x)\| \\ &\leq \left(1 + \alpha^4 + \dots + \alpha^{4(n-1)}\right) \frac{\alpha^2 + 1}{4\alpha^2}c \\ &\leq \frac{\alpha^2 + 1}{4\alpha^2} \frac{1}{1 - \alpha^4} (1 - \alpha^{4n})c, \text{ if } |\alpha| < 1. \end{aligned}$$

The “*altemative*” general inequality for  $|\alpha| < 1$ ;  $\alpha \neq 0$  is similarly established.

**Note 1.** (i) Assume  $|\alpha| > 1$ , and denote

$$Q_n : Q_n(x) = \alpha^{-4n} f(\alpha^n x).$$

Claim that sequence  $\{Q_n\}$ ,  $|\alpha| > 1$  is a Cauchy sequence.

In fact, if  $m > n > 0$ , then

$$\begin{aligned} 0 \leq \|Q_n(x) - Q_m(x)\| &= \|\alpha^{-4n} f(\alpha^n x) - \alpha^{-4m} f(\alpha^m x)\| \\ &= |\alpha|^{-4n} \|f(\alpha^n x) - \alpha^{-4(m-n)} f(\alpha^{m-n} \cdot \alpha^n x)\| \\ &\leq |\alpha|^{-4n} \frac{3\alpha^2 - 1}{4\alpha^2(\alpha^4 - 1)} (1 - \alpha^{-4(m-n)})c \\ &= \frac{3\alpha^2 - 1}{4\alpha^2(\alpha^4 - 1)} (|\alpha|^{-4n} - |\alpha|^{-4n} \alpha^{-4(m-n)})c \\ &\rightarrow 0, \text{ as } n \rightarrow \infty \text{ (and } m \rightarrow \infty). \end{aligned}$$

Thus  $\{Q_n\}$ ,  $|\alpha| > 1$ , is Cauchy sequence.

Similarly, if  $|\alpha| < 1$ ,  $\alpha \neq 0$ , one proves that

$$\{Q_n\}, |\alpha| < 1, \alpha \neq 0$$

is Cauchy sequence, as well.

(ii) Claim the quarticness of

$$Q : Q(x) = \lim_{n \rightarrow \infty} Q_n(x) = \lim_{n \rightarrow \infty} \alpha^{-4n} f(\alpha^n x), \alpha > 1.$$

In fact, replacing  $x \rightarrow \alpha^n x$ ,  $y \rightarrow \alpha^n y$  in the  $\alpha$ -quartic functional inequality (1.9) and then multiplying by  $|\alpha|^{-4n}$ , and taking limit  $n \rightarrow \infty$ , we obtain

$$\begin{aligned} 0 &\leq \|2[Q(\alpha x + y) + Q(x + \alpha y)] + \alpha(\alpha - 1)^2 Q(x - y) \\ &\quad - 2(\alpha^2 - 1)^2 [Q(x) + Q(y)] - \alpha(\alpha + 1)^2 Q(x + y)\| \\ &\leq |\alpha|^{-4n} c \rightarrow 0, \text{ } n \rightarrow \infty. \end{aligned}$$

Thus

$$2[Q(\alpha x + y) + Q(x + \alpha y)] + \alpha(\alpha - 1)^2Q(x - y) = 2(\alpha^2 - 1)^2[Q(x) + Q(y)] + \alpha(\alpha + 1)^2Q(x + y)$$

leading to (1.1) and thus the quarticness of  $Q : |\alpha| > 1$ .

Similarly, we prove that  $Q$  is quartic for  $|\alpha| < 1, \alpha \neq 0$ . Thus the existence of  $Q$  is complete.

If

$$Q(x) = \lim_{n \rightarrow \infty} Q_n(x) = \lim_{n \rightarrow \infty} \begin{cases} a^{-4n} f(a^n x) & \text{if } |a| > 1 \\ a^{4n} f(a^{-n} x) & \text{if } |a| < 1; a \neq 0 \end{cases}, \tag{1.13}$$

then

$$\|f(x) - Q(x)\| \leq \frac{2a^2 + |a^2 - 1|}{4a^2} c \cdot \begin{cases} \frac{1}{a^4 - 1} & \text{if } |a| > 1 \\ \frac{1}{1 - a^4} & \text{if } |a| < 1; a \neq 0. \end{cases} \tag{1.14}$$

**Note 2.** Claim the uniqueness of

$$Q : Q(x) = \lim_{n \rightarrow \infty} Q_n(x) = \lim_{n \rightarrow \infty} \alpha^{-4n} f(\alpha^n x), \alpha > 1.$$

In fact, if there is another quartic mapping  $Q'$  satisfying (1.14), then

$$\begin{aligned} 0 &\leq \|Q(x) - Q'(x)\| \\ &\leq \|Q(x) - f(x)\| + \|f(x) - Q'(x)\|, \end{aligned}$$

or

$$\begin{aligned} 0 &\leq \|Q(x) - Q'(x)\| \\ &= |\alpha|^{-4n} \|Q(\alpha^n x) - Q'(\alpha^n x)\| \\ &= \|\alpha^{-4n} Q(\alpha^n x) - \alpha^{-4n} Q'(\alpha^n x)\| \\ &\leq \|\alpha^{-4n} Q(\alpha^n x) - \alpha^{-4n} f(\alpha^n x)\| + \|\alpha^{-4n} f(\alpha^n x) - \alpha^{-4n} Q'(\alpha^n x)\| \\ &= |\alpha|^{-4n} \left\{ \|Q(\alpha^n x) - f(\alpha^n x)\| + \|f(\alpha^n x) - Q'(\alpha^n x)\| \right\} \\ &\leq \frac{2\alpha^2 + |\alpha^2 - 1|}{2\alpha^2} |\alpha|^{-4n} c \rightarrow 0, \quad n \rightarrow \infty \end{aligned}$$

or

$$Q(x) = Q'(x),$$

proving uniqueness of  $Q : |\alpha| > 1$ . Similarly, one proves uniqueness of  $Q : |\alpha| < 1, \alpha \neq 0$ .

**Theorem 1.1.** Let  $X$  be a normed space and  $Y$  be a Banach space. If  $f : X \rightarrow Y$  is a mapping satisfying (1.9), then there exists a unique quartic mapping  $Q : X \rightarrow Y$ , satisfying inequality (1.14).

If  $f(0) = 0$ , then  $\|f(x) - Q(x)\| \leq c/2|1 - a^4|$ , for  $\forall a \neq 0; \pm 1$ .

## 2. Stability of General $(a, b)$ -Quartic Functional Equation

$$2[f(ax + by) + f(bx + ay)] + ab(a - b)^2 f(x - y) = 2(a^2 - b^2)^2 [f(x) + f(y)] + ab(a + b)^2 f(x + y), \tag{2.1}$$

where  $a \neq \pm b, a, b \neq 0, \pm 1, k = a + b \neq 1, l = a - b \neq 1$ , and  $a^4 + b^4 - a^2 b^2 - 1 \neq 0$ .

Replacing  $x = y = 0$  in this equation, one gets

$$\begin{aligned} 4(a^4 + b^4 - a^2 b^2 - 1)f(0) &= 0, \text{ or} \\ f(0) &= 0 \end{aligned} \tag{2.2}$$

Similarly substituting  $x = x, y = 0$  in (2.1), we obtain

$$f(ax) + f(bx) = (a^4 + b^4)f(x) + (a^2 - b^2)^2 f(0) = (a^4 + b^4)f(x). \tag{2.3}$$

From (2.3), we observe that

$$“f(ax) = a^4 f(x) \quad \text{if and only if} \quad f(bx) = b^4 f(x).” \tag{2.4}$$

Let us introduce the following “ $(a, b)$ -quartic functional mean”

$$\overline{f_{(a,b)}}(x) = \frac{f(ax) + f(bx)}{a^4 + b^4}. \tag{2.5}$$

From (2.3) and (2.5), we find the quartic functional mean equation

$$\overline{f_{(a,b)}}(x) = f(x) \tag{2.6}$$

From (2.4)-(2.5)-(2.6), one establishes

$$“\overline{f_{(a,0)}}(x) = \frac{f(ax)}{a^4} = f(x) \quad \text{iff} \quad \overline{f_{(0,b)}}(x) = \frac{f(bx)}{b^4} = f(x).”$$

Also assuming  $f(2x) = 16f(x)$ , replacing  $x = x, y = x$  in (2.1) and setting  $k = a + b \neq 0, \pm 1$ , one obtains

$$4f(k) + ab(a - b)^2 f(0) = 4(a^2 - b^2)^2 f(x) + ab(a + b)^2 f(2x), \tag{2.8}$$

or

$$f(k) + \frac{1}{4}ab(a - b)^2 f(0) = (a^2 - b^2)^2 f(x) + 4ab(a + b)^2 f(x) = k^4 f(x),$$

or

$$f(k) = k^4 f(x) - \frac{1}{4}ab(a - b)^2 f(0) = k^4 f(x). \tag{2.9}$$

Without assuming  $f(2x) = 16f(x)$ , replacing  $x = x, y = -x$  in (2.1) and setting  $l = a - b \neq 0, \pm 1$  with  $a \neq \pm b$ , one obtains

$$2[f(lx) + f(-lx)] + ab(a - b)^2 f(2x) = 2(a^2 - b^2)^2 [f(x) + f(-x)] + ab(a + b)^2 f(0) \tag{2.10}$$

Placing  $-x$  on  $x$  in (2.10), and then subtracting the new equation from (2.10), we get  $ab(a - b)^2 [f(2x) - f(-2x)] = 0$ . Letting  $x/2$  on  $x$ , we find that  $f$  is an “*even function*”, such that  $f(-x) = f(x)$ . Thus from (2.10), we obtain

$$4f(lx) + ab(a - b)^2 f(2x) = 4(a^2 - b^2)^2 f(x) + ab(a + b)^2 f(0). \tag{2.11}$$

Assuming  $f(2x) = 16f(x)$ , we get

$$f(lx) = l^4 f(x) + \frac{1}{4}ab(a + b)^2 f(0) = l^4 f(x).$$

Without assuming  $f(2x) = 16f(x)$ , subtracting (2.11) from (2.8), we obtain from (2.2) that

$$f(kx) - f(lx) = \frac{1}{2}ab(a^2 + b^2)f(2x) = \left[ \left(\frac{k}{2}\right)^4 - \left(\frac{l}{2}\right)^4 \right] f(2x).$$

Replacing  $x \rightarrow x/2$ , we obtain that

$$f\left(\frac{k}{2}x\right) - \left(\frac{k}{2}\right)^4 f(x) = f\left(\frac{l}{2}x\right) - \left(\frac{l}{2}\right)^4 f(x).$$

Therefore, we observe that

$$“f\left(\frac{k}{2}x\right) = \left(\frac{k}{2}\right)^4 f(x) \quad \text{if and only if} \quad f\left(\frac{l}{2}x\right) = \left(\frac{l}{2}\right)^4 f(x).” \tag{2.12}$$

Employing the “quartic mean”, we have equivalently that

$$“\bar{f}_{\frac{k}{2}}(x) \frac{f(\frac{k}{2}x)}{(\frac{k}{2})^4} = f(x) \text{ iff } \bar{f}_{\frac{1}{2}}(x) = \frac{f(\frac{1}{2}x)}{(\frac{1}{2})^4} = f(x)”.$$

Let  $X$  be a real normed linear space and let  $Y$  be a real complete normed linear space. Assume  $f : X \rightarrow Y$ , satisfying the following general  $(a, b)$ -quartic functional inequality

$$\|2[f(ax+by) + f(bx+ay)] + ab(a-b)^2 f(x-y) - 2(a^2 - b^2)^2[f(x) + f(y)] - ab(a+b)^2 f(x+y)\| \leq c, \tag{2.13}$$

where  $a, b \neq 0; a, b \neq \pm 1$ . Replacing  $x = y = 0$  in (2.13), one gets

$$\|f(0)\| \leq c/4|a^4 + b^4 - a^2b^2 - 1|. \tag{2.14}$$

Substituting  $x = x, y = 0$  in (2.13), and employing the triangle inequality and (2.14), we obtain

$$\|f(ax) + f(bx) - (a^4 + b^4)f(x)\| \leq \frac{c}{2} + (a^2 - b^2)^2|f(0)|,$$

or

$$\|f(ax) + f(bx) - (a^4 + b^4)f(x)\| \leq c \frac{(a^2 - b^2)^2 + 2|a^4 + b^4 - a^2b^2 - 1|}{4|a^4 + b^4 - a^2b^2 - 1|}. \tag{2.15}$$

Substituting  $x = x, y = x$  in (2.13), and employing the triangle inequality and (2.14), as well as the following hypothesis

$$\|f(2x) - 16f(x)\| \leq c_1 (\geq 0), \tag{2.16}$$

and denoting  $k = a + b \neq 0, \pm 1$ , we obtain:

$$\|f(kx) - k^4 f(x)\| \leq \frac{1}{4}(c + ab(a-b)^2|f(0)| + ab(a+b)^2c_1),$$

or

$$\|f(kx) - k^4 f(x)\| \leq c_2 = \frac{1}{4} \left( \frac{ab(a-b)^2 + 4|a^4 + b^4 - a^2b^2 - 1|}{4|a^4 + b^4 - a^2b^2 - 1|} c + ab(a+b)^2c_1 \right), \tag{2.17}$$

or

$$\|f(x) - k^4 f(kx)\| \leq k^{-4}c_2 \tag{2.18}$$

if  $|k| > 1$ .

Thus we easily obtain, the following general inequality:

$$\|f(x) - k^{-4n} f(k^n x)\| \leq \frac{1}{k^4 - 1}(1 - k^{-4n})c_2, \text{ if } |k| > 1. \tag{2.19}$$

In fact,

$$\begin{aligned} \|f(x) - k^{-4n} f(k^n x)\| &\leq \|f(x) - k^{-4} f(k^1 x)\| + k^{-4} \|f(k^1 x) - k^{-4} f(k^2 x)\| + \dots \\ &\quad \dots + k^{-4(n-2)} \|f(k^{n-2} x) - k^{-4} f(k^{n-1} x)\| \\ &\quad + k^{-4(n-1)} \|f(k^{n-1} x) - k^4 f(k^n x)\| \tag{2.20} \\ &\leq k^{-4} \left( 1 + k^{-4} + k^{-8} + \dots + k^{-4(n-2)} + k^{-4(n-1)} \right) c_2 \\ &= \frac{k^{-4}}{1 - k^{-4}}(1 - k^{-4n})c_2 = \frac{1}{k^4 - 1}(1 - k^{-4n})c_2, \text{ if } |k| > 1. \end{aligned}$$

The “*alternative*” general inequality is similarly established, as follows

$$\|f(x) - k^{4n} f(k^{-n} x)\| \leq \frac{1}{1 - k^4}(1 - k^{4n})c_2, \text{ if } |k| < 1; k \neq 0. \tag{2.21}$$

In fact,

$$\begin{aligned} \|f(x) - k^{4n}f(k^{-n}x)\| &\leq \|f(x) - k^4f(k^{-1}x)\| + k^4\|f(k^{-1}x) - k^4f(k^{-2}x)\| + \dots \\ &\quad \dots + k^{4(n-2)}\|f(k^{-(n-2)}x) - k^4f(k^{-(n-1)}x)\| \\ &\quad + k^{4(n-1)}\|f(k^{-(n-1)}x) - k^4f(k^{-n}x)\| \\ &\leq (1 + k^4 + k^8 + \dots + k^{4(n-2)} + k^{4(n-1)})c_2 \\ &\leq \frac{1}{1 - k^4}(1 - k^{4n})c_2, \text{ if } |k| < 1; k \neq 0. \end{aligned} \tag{2.22}$$

If we denote

$$Q(x) = \lim_{n \rightarrow \infty} Q_n(x) = \lim_{n \rightarrow \infty} \begin{cases} k^{-4n}f(k^n x) & \text{if } |k| > 1 \\ k^{4n}f(k^{-n}x) & \text{if } |k| < 1; k \neq 0. \end{cases}$$

It follows

$$|f(x) - Q(x)| \leq c_2 \cdot \begin{cases} \frac{1}{k^4 - 1} & \text{if } |k| > 1 \\ \frac{1}{1 - k^4} & \text{if } |k| < 1; k \neq 0. \end{cases} \tag{2.23}$$

**Note 3.** Following Notes 1-2, we establish the existence and uniqueness of the quartic mapping  $Q$ .

If  $f(0) = 0$ , and  $f(2x) = 16f(x)$ , then

$$\|f(x) - Q(x)\| \leq \frac{c}{2|1 - k^4|}, \text{ for } \forall k \neq 0; \pm 1. \tag{2.24}$$

**Theorem 2.1.** Let  $X$  be normed space and  $Y$  a Banach space. If  $f : X \rightarrow Y$  is a mapping satisfying (2.13) then there exists a unique quartic mapping  $Q : X \rightarrow Y$ , satisfying inequality (2.23).

If  $f(0) = 0$ , and  $f(2x) = 16f(x)$ , then

$$\|f(x) - Q(x)\| \leq \frac{c}{2|1 - k^4|}, \text{ for } \forall k = a + b \neq 0; \pm 1.$$

### 3. General Alternative $a$ -Quartic Functional Equation

$$\begin{aligned} &2f(ax + y) + f(x + ay) + f(ax - y) \\ &= \frac{a}{2}[(a^2 + 4a + 1)f(x + y) - (a^2 - 4a + 1)f(x - y)] \\ &\quad + (3a^4 - 4a^2 + 1)f(x) + (a^4 - 4a^2 + 3)f(y), \end{aligned} \tag{3.1}$$

where  $a \neq 0, a \neq \pm 1$ .

Replacing  $x = y = 0$  in (3.1), one gets

$$4a^2(a^2 - 1)f(0) = 0,$$

or

$$f(0) = 0. \tag{3.2}$$

Substituting  $x = x, y = 0$  in (3.1), we obtain

$$f(ax) = a^4f(x) + \frac{1}{3}(a^4 - 4a^2 + 3)f(0) = a^4f(x). \tag{3.3}$$

Let  $X$  be a real normed linear space and let  $Y$  be a real complete normed linear space. Assume  $f : X \rightarrow Y$ , satisfying the following general alternative  $a$ -quartic functional inequality

$$\begin{aligned} &\|2f(ax + y) + f(x + ay) + f(ax - y) \\ &\quad - \frac{a}{2}[(a^2 + 4a + 1)f(x + y) - (a^2 - 4a + 1)f(x - y)]\| \end{aligned}$$

$$-(3a^4 - 4a^2 + 1)f(x) - (a^4 - 4a^2 + 3)f(y) \Big\| \leq c, \tag{3.4}$$

where  $a \neq 0, a \neq \pm 1$ . Replacing  $x = y = 0$  in (3.4), one gets

$$4a^2|a^2 - 1| \|f(0)\| \leq c,$$

or

$$\|f(0)\| \leq \frac{c}{a^2|a^2 - 1|}. \tag{3.5}$$

Substituting  $x = x, y = 0$  in (3.4) and employing (3.5), we obtain

$$\|f(x) - a^{-4}f(ax)\| \leq \frac{3a^2 + |a^2 - 3|}{3a^2} a^{-4}c = a^{-4}c_1. \tag{3.6}$$

Thus

$$\|f(x) - a^{-4n}f(a^n x)\| \leq \frac{1}{a^4 - 1} (1 - a^{-4(n+1)})c_1. \tag{3.7}$$

Similarly, we obtain

$$\|f(x) - a^4f(a^{-1}x)\| \leq \frac{3a^2 + |a^2 - 3|}{3a^2} c = c_1. \tag{3.8}$$

Thus

$$\|f(x) - a^{4n}f(a^{-n}x)\| \leq \frac{1}{1 - a^4} (1 - a^{4(n+1)})c_1. \tag{3.9}$$

If

$$Q(x) = \lim_{n \rightarrow \infty} Q_n(x) = \lim_{n \rightarrow \infty} \begin{cases} a^{-4n}f(a^n x) & \text{if } |a| > 1 \\ a^{4n}f(a^{-n}x) & \text{if } |a| < 1; a \neq 0 \end{cases}, \text{ then}$$

$$\|f(x) - Q(x)\| \leq \frac{3a^2 + |a^2 - 3|}{3a^2} c \cdot \begin{cases} \frac{1}{a^4 - 1} & \text{if } |a| > 1 \\ \frac{1}{1 - a^4} & \text{if } |a| < 1; a \neq 0. \end{cases} \tag{3.10}$$

**Note 4.** Following Notes 1-2, we establish existence and uniqueness of  $Q$ .

If  $f(0) = 0$ , then  $\|f(x) - Q(x)\| \leq \frac{c}{2|1 - a^4|}$ , for  $\forall a \neq 0; \pm 1$ .

**Theorem 3.1.** Let  $X$  be a normed space and  $Y$  a Banach space. If  $f : X \rightarrow Y$  is a mapping satisfying (3.1), then there exists a unique quartic mapping  $Q : X \rightarrow Y$ , satisfying inequality (3.4).

If  $f(0) = 0$ , then  $\|f(x) - Q(x)\| \leq c/2|1 - a^4|$ , for  $\forall a \neq 0; \pm 1$ .

### OPEN RESEARCH PROBLEMS

#### OPEN PROBLEM A.

Employing both the “*Hyers’ direct method*” and the “*fixed point method*”, it is still “*open*”, the investigation of “*generalized Ulam stabilities*” and “*generalized Ulam superstabilities*” of these quartic functional equations in various normed spaces, domains and groups such as in

1. Banach spaces;
2. Banach algebras;  $C^*$ -algebras;
3.  $N$ -multi-Banach spaces; multi-Banach spaces;
4. Multi-normed spaces;
5. Quasi-Banach spaces;
6. Quasi- $\beta$ [beta]-normed spaces;
7. Non-Archimedean normed spaces;
8. Fuzzy normed spaces;

9. Quasi fuzzy normed spaces;
10. Non- Archimedean fuzzy normed spaces;
11. Intuitionistic normed spaces;
12. Random normed spaces; and probabilistic normed spaces;
13. Non-Archimedean  $RN$ [Random Normed]-spaces;
14. Intuitionistic random normed spaces;
15. Intuitionistic fuzzy normed spaces;
16. intuitionistic fuzzy Banach algebras;
17. Intuitionistic Non-Archimedean fuzzy normed spaces;
18. Menger normed spaces;
19. Menger probabilistic normed spaces;
20. Non-Archimedean Menger normed spaces;
21. Intuitionistic Menger normed spaces;
22.  $L$ -non-Archimedean- fuzzy Euclidean normed spaces;
23.  $F$ -spaces; Fréchet spaces;
24. Banach modules;
25. Distributions and Hyperfunctions;  
as well as, on:
26. Restricted domains;
27. Heisenberg groups.

**OPEN PROBLEM B.**

28. Exploiting the elementary “*M. Hosszu’s method*”, due to M. Hosszu (see: “On the Fréchet’s functional equation”, Bull. Inst. Politech. Iasi 10, (1964), 1-2, 27-28), determine the general solution and the Ulam stability of each one of these quartic functional equations. The advantage of this method is that we do not assume any regularity conditions on the unknown function  $f$ . See also the paper of the authors Xu et al.: [31].
29. Employing two “*L. Szekelyhidi’s fundamental results*”, due to L. Szekelyhidi (see: “Convolution type functional equation on topological abelian groups”, World Scientific, Singapore, 1991), determine the general solution and the Ulam stability of each one of these quartic functional equations in certain types of groups, such as, “commutative groups”. See: [31].

**OPEN PROBLEM C.**

30. Exploiting the elementary “*M. Hosszu’s method*”, due to M. Hosszu (see: “On the Fréchet’s functional equation”, Bull. Inst. Politech. Iasi 10, (1964), 1-2, 27-28 ), determine the general solution and the Ulam stability of each one of these pertinent “*Pexider quartic*” functional equations “with or without *involution*” [32]. We do not assume any regularity conditions on the unknown functions. For “*Pexider quartic*” equations [31]:

$$(1) \quad f(ax + y) + f(x + ay) = g(x + y) + \bar{g}(x - y) + h(x) + \bar{h}(y),$$

$$(2) \quad f(ax + by) + f(bx + ay) = g(x + y) + \bar{g}(x - y) + h(x) + \bar{h}(y),$$

with fixed integers  $a, b \neq 0, \pm 1$  and unknown functions  $f, \bar{f}, g, \bar{g}, h, \bar{h}$ . See also the papers of the author, et al.: ( [31, 33]).

31. Employing two “*L. Szekelyhidi’s fundamental results*”, due to L. Szekelyhidi (see: “Convolution type functional equation on topological abelian groups”, World Scientific, Singapore, 1991), determine the general solution and the Ulam stability of each one of these pertinent “*Pexider quartic*” functional equations with or without “*involution*” in certain types of groups, such as, “commutative groups”. See: [31, 33].
32. Investigate Ulam-Hyers stabilities of pertinent “*quartic derivations*” from a Banach algebra into its Banach modules. See: ( [34, 35]).

33. Establish the solution and stabilities of “*conditionally*” quartic functional equations, for instance:  $\|x\| = \|y\|$ .

34. Prove stabilities of “*orthogonally*” quartic functional equations, “*in the sense of J. Rätz*” [36]: “ $x \perp y \iff \langle x, y \rangle = 0$ ”.

35. Work on stabilities of “*quartic-like*” functional equations, such as:

$$\begin{aligned}
 1. \quad & 2[f_1(ax + y) + f_2(x + ay + \sigma(y))] + a(a - 1)^2g_1(x - y) \\
 & = 2(a^2 - 1)^2[h_1(x) + h_2(y)] + a(a + 1)^2g_2(x + y), \\
 & \sigma = \sigma(y) \text{ is “}i\text{nvolution”}: \sigma(x + y) = \sigma(x) + \sigma(y); \sigma(\sigma(x)) = x.
 \end{aligned}$$

$$\begin{aligned}
 2. \quad & 2f(ax + y + \rho) + f(x + ay + \tau) + f(ax - y) \\
 & = \frac{a}{2}[(a^2 + 4a + 1)f(x + y) - (a^2 - 4a + 1)f(x - y)] \\
 & \quad + (3a^4 - 4a^2 + 1)f(x) + (a^4 - 4a^2 + 3)f(y).
 \end{aligned}$$

**Acknowledgement.** The authors gratefully acknowledge the financial support from King Abdulaziz University, Jeddah, Saudi Arabia.

## References

- [1] S.M. Ulam, A Collection of Mathematical Problems, Interscience Tracts in Pure and Applied Mathematics, No.8, Interscience. Publ., New York , 1960; “Problems in Modern Mathematics”, Ch. VI, Science Ed., Wiley, 1940.
- [2] D.H. Hyers, On the stability of the linear functional equation, Proc. Nat. Acad. Sci., USA, 27 (1941) 222-224.
- [3] J.M. Rassias, On the stability of the Euler-Lagrange functional equation, C. R. Acad. Bulgare Sci. 45 (1992) 17-20.
- [4] J.M. Rassias, On the stability of the Euler-Lagrange functional equation, Chinese J. Math. 20 (1992) 185-190.
- [5] J.M. Rassias, Solution of a problem of Ulam, J. Approx. Theory 57 (1989) 268-273.
- [6] J.M. Rassias, On a new approximation of approximately linear mappings by linear mappings, Discuss, Math. 7 (1985) 193-196.
- [7] J.M. Rassias, On approximation of approximately linear mappings by linear mappings, J. Funct. Anal. 46 (1982) 126-130.
- [8] T. Aoki, On the stability of the linear transformation in Banach spaces, J. Math. Soc. Japan, 2 (1950) 64-66.
- [9] T.M. Rassias, On the stability of the linear mapping in Banach spaces, Proc. Am. Math. Soc. 72 (1978) 297-300.
- [10] Z. Gajda, On stability of additive mappings, Int. J. Math. Math. Sci. 14 (1991) 431-434.
- [11] T.M. Rassias, P. Šemrl, On the behaviour of mappings which do not satisfy Hyers-Ulam stability, Proc. Am. Math. Soc. 114 (1992) 989-993.
- [12] P. Găvruta, A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings, J. Math. Anal. Appl. 184 (1994) 431-436.
- [13] S.-M. Jung, On the Hyers-Ulam stability of the functional equations that have the quadratic property, J. Math. Anal. Appl. 222 (1998) 126-137.
- [14] K.W. Jun, H.M. Kim, The generalized Hyers-Ulam-Rassias stability of a cubic functional equation, J. Math. Anal. Appl. 274 (2002) 867-878.
- [15] R.P. Agarwal, B. Xu, W. Zhang, Stability of functional equations in single variable, J. Math. Anal. Appl. 288 (2003) 852-869
- [16] S.A. Mohiuddine, H. Sevli, Stability of Pexiderized quadratic functional equation in intuitionistic fuzzy normed space, J. Comp. Appl. Math., 235 (2011) 2137-2146.
- [17] S.A. Mohiuddine, Stability of Jensen functional equation in intuitionistic fuzzy normed space, Chaos, Solitons Fract., 42 (2009) 2989-2996.
- [18] M. Mursaleen, S.A. Mohiuddine, On stability of a cubic functional equation in intuitionistic fuzzy normed spaces, Chaos, Solitons Fract., 42 (2009) 2997-3005.
- [19] S.A. Mohiuddine, A. Alotaibi, Fuzzy stability of of a cubic functional equation via fixed point technique, Adv. Difference Equ., 2012 , 2012:48.
- [20] J.M. Rassias, Solution of the Ulam stability problem for quartic mappings, Glas. Mat. Ser. III 34(54) (1999) 243-252.



- [21] S.-M. Jung, Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis, Hadronic Press, Palm Harbor (2001)
- [22] M. Mursaleen, K.J. Ansari, Stability results in intuitionistic fuzzy normed spaces for a cubic functional equation, *Appl. Math. Inf. Sci.* 7(5) (2013) 1685-1692.
- [23] A. Najati, C. Park, On the stability of an  $n$ -dimensional functional equation originating from quadratic forms, *Taiwan. J. Math.* 12 (2008) 1609-1624.
- [24] T.M. Rassias, On the stability of functional equations and a problem of Ulam, *Acta Appl. Math.* 62 (2000) 123-130.
- [25] C. Park, Y. Cho, M. Han, Functional inequalities associated with Jordan-von Neumann-type additive functional equations. *J. Inequal. Appl.* 2007, Article ID 41820 (2007).
- [26] C. Park, J.R. Lee, D.Y. Shin, Functional equations and inequalities in matrix paranormed spaces, *J. Inequal. Appl.* 2013, 2013:547
- [27] Z. Wang, T.M. Rassias, Intuitionistic fuzzy stability of functional equations associated with inner product spaces, *Abstr. Appl. Anal.* Volume 2011, Article ID 456182, 19 pages.
- [28] I.-S. Chang, Higher ring derivation and intuitionistic fuzzy stability, *Abstr. Appl. Anal.* Volume 2012, Article ID 503671, 16 pages.
- [29] J. Roh, I.-S. Chang, On the intuitionistic fuzzy stability of ring homomorphism and ring derivation, *Abstr. Appl. Anal.* Volume 2013, Article ID 192845, 8 pages.
- [30] M. Mirmostafae, M.S. Moslehian, Fuzzy versions of Hyers-Ulam-Rassias theorem, *Fuzzy Sets Syst.* 159 (2008) 720-729.
- [31] T.Z. Xu, J.M. Rassias, W.X. Xu, A generalized mixed Quadratic-Quartic functional equation, *Bull. Malays. Math. Sci. Soc.* 35(3) (2012) 633-649.
- [32] J.M. Rassias, H.-M. Kim, Approximate homomorphisms and derivations between  $C^*$ -ternary algebras. *J. Math. Phys.* 49 (2008), no. 6, 063507, 10 pp.
- [33] M.M. Pourpasha, J.M. Rassias, R. Saadati, S.M. Vaezpour, A fixed point approach to the stability of Pexider quadratic functional equation with involution, *J. Inequal. Appl.* 2010, Art. ID 839639, 18 pp.
- [34] M.E. Gordji, N. Ghobadipour, Generalized Ulam-Hyers stabilities of quartic derivations on Banach algebras, *Proyecciones J. Math.*, 29 (2010) 209-226.
- [35] C. Park, J.M. Rassias, Cubic derivations and quartic derivations on Banach modules, in: "Functional Equations, Difference Inequalities and Ulam Stability Notions" (F. U. N.), Editor: J.M. Rassias, 2010, 119-129, ISBN 978-1-60876-461-7, Nova Science Publishers, Inc.
- [36] J. Rätz, On the orthogonal additive mappings, *Aequationes Math.* 28 (1985) 35-49.

# Existence of positive solutions for summation boundary value problem for a fourth-order difference equations

Thanin Sitthiwirattham

Department of Mathematics, Faculty of Applied Science,  
King Mongkut's University of Technology North Bangkok, Bangkok 10800, Thailand  
E-mail address: tst@kmutnb.ac.th

## Abstract

In this paper, we study the existence of positive solutions to the difference-summation boundary value problem

$$\Delta^4 u(t-2) + a(t)f(u) = 0, \quad t \in \{2, 3, \dots, T\},$$

$$u(0) = \Delta u(0) = \Delta^2 u(0) = 0, \quad u(T+2) = \alpha \sum_{s=4}^{\eta} u(s),$$

where  $f$  is continuous,  $T \geq 5$  is a fixed positive integer,  $\eta \in \{4, 5, \dots, T-1\}$ ,  $0 < \alpha < \frac{4T(T+1)(T+2)}{(\eta-3)(\eta+2)(\eta^2-\eta+4)}$ . We show the existence of at least one positive solution if  $f$  is either superlinear or sublinear by applying Guo-Krasnoselskii fixed point theorem in cones.

*Keywords* : Positive solution; Boundary value problem; Fixed point theorem; Cone  
*2010 Mathematics Subject Classification*: 39A15, 34B15

## 1 Introduction

The existence of solutions for boundary value problems of difference equations has received much attention. For example, see [4-15] and the references therein.

Liang *et al.* in [4] considered the fourth-order boundary value problem of the form

$$\begin{cases} \Delta^4 x(t-2) + a(t)f(x) = 0, & t \in \{2, 3, \dots, T\}, \\ x(0) = x(T+2) = 0, \quad \Delta^2 x(0) = \Delta^2 x(T) = 0, \end{cases}$$

where  $T > 2$ . Existence and uniqueness of solutions are obtained by a fixed point theorem.

Ma *et al.* in [5] considered the fourth-order boundary value problem of the form

$$\begin{cases} \Delta^4 u(t-2) - \lambda f(t, u(t)) = 0, & t \in \{2, 3, \dots, T\}, \\ u(1) = u(T+1) = \Delta^2 u(0) = \Delta^2 u(T) = 0, \end{cases}$$

where  $\lambda$  is a parameter,  $T > 5$ . Existence and uniqueness of solutions are obtained by the theory of fixed-point index in cones.

In this paper, we consider the existence of positive solutions to the equation

$$\Delta^4 u(t-2) + a(t)f(u) = 0, \quad t \in \{2, 3, \dots, T\}, \tag{1.1}$$

with summation boundary condition

$$u(0) = \Delta u(0) = \Delta^2 u(0) = 0, \quad u(T+2) = \alpha \sum_{s=4}^{\eta} u(s), \tag{1.2}$$

where  $f$  is continuous.

The aim of this paper is to give some results for existence of positive solutions to (1.1)-(1.2).

Let  $\mathbb{N}$  be the nonnegative integer, we let  $\mathbb{N}_{i,j} = \{k \in \mathbb{N} \mid i \leq k \leq j\}$  and  $\mathbb{N}_p = \mathbb{N}_{0,p}$ . By the positive solution of (1.1)-(1.2) we mean that a function  $u(t) : \mathbb{N}_{T+2} \rightarrow [0, \infty)$  and satisfies the problem (1.1)-(1.2).

Throughout this paper, we suppose the following conditions hold:

- (H1)  $T \geq 5$  is a fixed positive integer,  $\eta \in \{4, 5, \dots, T-1\}$ , constant  $\alpha > 0$  such that  $0 < \alpha < \frac{4T(T+1)(T+2)}{(\eta-3)(\eta+2)(\eta^2-\eta+4)}$ .
- (H2)  $f \in C([0, \infty), [0, \infty))$ ,  $f$  is either superlinear or sublinear. Set

$$f_0 = \lim_{u \rightarrow 0^+} \frac{f(u)}{u}, \quad f_\infty = \lim_{u \rightarrow \infty} \frac{f(u)}{u}.$$

Then  $f_0 = 0$  and  $f_\infty = \infty$  correspond to the superlinear case, and  $f_0 = \infty$  and  $f_\infty = 0$  correspond to the sublinear case.

- (H3)  $a \in C(\mathbb{N}_{2,T}, [0, \infty))$ ,  $a$  is not identical zero.

Existence of positive solutions for summation boundary value problem ... 3

The proof of the main theorem is based upon an application of the following Guo-Krasnoselskii's fixed point theorem in a cone.

**Theorem 1.1.** *Let  $E$  be a Banach space, and let  $K \subset E$  be a cone. Assume  $\Omega_1, \Omega_2$  are open subsets of  $E$  with  $0 \in \Omega_1, \bar{\Omega}_1 \subset \Omega_2$ , and let*

$$A : K \cap (\bar{\Omega}_2 \setminus \Omega_1) \longrightarrow K$$

*be a completely continuous operator such that*

(i)  $\|Au\| \leq \|u\|, \quad u \in K \cap \partial\Omega_1, \text{ and } \|Au\| \geq \|u\|, \quad u \in K \cap \partial\Omega_2;$  or

(ii)  $\|Au\| \geq \|u\|, \quad u \in K \cap \partial\Omega_1, \text{ and } \|Au\| \leq \|u\|, \quad u \in K \cap \partial\Omega_2.$

*Then  $A$  has a fixed point in  $K \cap (\bar{\Omega}_2 \setminus \Omega_1)$ .*

## 2 Preliminaries

We now state and prove several lemmas before stating our main results.

**Lemma 2.1.** *Suppose that  $y(t) \in C(\mathbb{N}_{2,T})$  and  $y(t) > 0$ . Then the linear boundary value problem*

$$\Delta^4 u(t-2) + y(t) = 0, \quad t \in \mathbb{N}_{2,T}, \tag{2.1}$$

$$u(0) = \Delta u(0) = \Delta^2 u(0) = 0, \quad u(T+2) = \alpha \sum_{s=4}^{\eta} u(s), \tag{2.2}$$

*has a unique solution*

$$u(t) = \frac{t^3}{6\Lambda} \left[ \sum_{s=2}^T (T-s+3)^3 y(s) - \frac{\alpha}{4} \sum_{s=2}^{\eta-2} (\eta-s+2)^4 y(s) \right] - \frac{1}{6} \sum_{s=2}^{t-2} (t-s+1)^3 y(s), \tag{2.3}$$

*where*

$$\Lambda := T(T+1)(T+2) - \frac{\alpha}{4}(\eta-3)(\eta+2)(\eta^2 - \eta + 4). \tag{2.4}$$

**Proof.** In fact, if  $u(t)$  is a solution of problem (2.1), by the discrete Taylor expansion formula, we have

$$u(t) = C_1 t^3 + C_2 t^2 + C_3 t^1 + C_4 - \frac{1}{6} \sum_{s=0}^{t-4} (t-s-1)^3 y(s+2), \quad t \in \mathbb{N}_{T+2}.$$

4

Thanin Sitthiwiratttham

Applying the first boundary condition  $u(0) = \Delta u(0) = \Delta^2 u(0) = 0$  in (2.2), we obtain

$$C_2 = C_3 = C_4 = 0.$$

So,

$$u(t) = C_1 t^3 - \frac{1}{6} \sum_{s=0}^{t-4} (t-s-1)^3 y(s+2), \tag{2.5}$$

From (2.5) and the second boundary condition in (2.2) implies

$$\begin{aligned} \alpha \sum_{s=4}^{\eta} u(s) &= \alpha C_1 \sum_{s=4}^{\eta} (s)^3 - \frac{\alpha}{6} \sum_{s=4}^{\eta} \sum_{\xi=0}^{s-4} (s-\xi-1)^3 y(\xi+2) \\ &= \alpha C_1 \sum_{s=0}^{\eta-4} (s+4)^3 - \frac{\alpha}{6} \sum_{s=0}^{\eta-4} \sum_{\xi=0}^{\eta-s-4} (\xi+3)^3 y(s+2) \\ &= C_1 (T+2)^3 - \frac{1}{6} \sum_{s=0}^{T-2} (T-s+1)^3 y(s+2). \end{aligned}$$

Solving the above equation for a constant  $C_1$ , we get

$$C_1 = \frac{1}{6\Lambda} \sum_{s=0}^{T-2} (T-s+1)^3 y(s+2) - \frac{\alpha}{6\Lambda} \sum_{s=0}^{\eta-4} \sum_{\xi=0}^{\eta-s-4} (\xi+3)^3 y(s+2)$$

where  $\Lambda$  is defined by (2.4)

Therefore, (2.1)-(2.2) has a unique solution

$$\begin{aligned} u(t) &= \frac{t^3}{6\Lambda} \left[ \sum_{s=2}^T (T-s+3)^3 y(s) - \frac{\alpha}{4} \sum_{s=2}^{\eta-2} (\eta-s+2)^4 y(s) \right] \\ &\quad - \frac{1}{6} \sum_{s=2}^{t-2} (t-s+1)^3 y(s). \end{aligned}$$

□

**Lemma 2.2.** *The function*

$$G(t, s) = \frac{1}{6\Lambda} \begin{cases} -\Lambda(t-s+1)^3 + t^3(T-s+3)^3 - \frac{\alpha t^3}{4}(\eta-s+2)^4, & s \in \mathbb{N}_{2,t-2} \cap \mathbb{N}_{2,\eta-2} \\ -\Lambda(t-s+1)^3 + t^3(T-s+3)^3, & s \in \mathbb{N}_{\eta-1,t-2} \\ t^3(T-s+3)^3 - \frac{\alpha t^3}{4}(\eta-s+2)^4, & s \in \mathbb{N}_{t-1,\eta-2} \\ t^3(T-s+3)^3, & s \in \mathbb{N}_{t-1,T} \cap \mathbb{N}_{\eta-1,T} \end{cases} \tag{2.6}$$

Existence of positive solutions for summation boundary value problem ... 5

where  $\Lambda$  is defined by (2.4), is the Green's function of the problem

$$\begin{aligned}
 -\Delta^4 u(t-2) &= 0, \quad t \in \mathbb{N}_{2,T}, \\
 u(0) = \Delta u(0) = \Delta^2 u(0) &= 0, \quad u(T+2) = \alpha \sum_{s=4}^{\eta} u(s). \quad (2.7)
 \end{aligned}$$

**Proof.** Suppose  $t < \eta$ . The unique solution of problem (2.1)-(2.2) can be written

$$\begin{aligned}
 u(t) &= -\frac{1}{6} \sum_{s=2}^{t-2} (t-s+1)^3 y(s) \\
 &\quad + \frac{t^3}{6\Lambda} \left[ \sum_{s=2}^{t-2} (T-s+3)^3 y(s) + \sum_{s=t-1}^{\eta-2} (T-s+3)^3 y(s) + \sum_{s=\eta-1}^T (T-s+3)^3 y(s) \right] \\
 &\quad - \frac{\alpha t^3}{24\Lambda} \left[ \sum_{s=2}^{t-2} (\eta-s+2)^4 y(s) + \sum_{s=t-1}^{\eta-2} (\eta-s+2)^4 y(s) \right] \\
 &= \frac{1}{6\Lambda} \sum_{s=2}^{t-2} \left[ -\Lambda(t-s+1)^3 + t^3(T-s+3)^3 - \frac{\alpha t^3}{4}(\eta-s+2)^4 \right] y(s) \\
 &\quad + \frac{1}{6\Lambda} \sum_{s=t-1}^{\eta-2} \left[ t^3(T-s+3)^3 - \frac{\alpha t^3}{4}(\eta-s+2)^4 \right] y(s) \\
 &\quad + \frac{1}{6\Lambda} \sum_{s=\eta-1}^T t^3(T-s+3)^3 y(s) \\
 &= \sum_{s=2}^T G(t,s)y(s).
 \end{aligned}$$

Suppose  $t \geq \eta$ . The unique solution of problem (2.1)-(2.2) can be written

$$\begin{aligned}
 u(t) &= -\frac{1}{6} \left[ \sum_{s=2}^{\eta-2} (t-s+1)^3 y(s) + \sum_{s=\eta-1}^{t-2} (t-s+1)^3 y(s) \right] \\
 &\quad + \frac{t^3}{6\Lambda} \left[ \sum_{s=2}^{\eta-2} (T-s+3)^3 y(s) + \sum_{s=\eta-1}^{t-2} (T-s+3)^3 y(s) + \sum_{s=t-1}^T (T-s+3)^3 y(s) \right] \\
 &\quad - \frac{\alpha t^3}{24\Lambda} \sum_{s=2}^{\eta-2} (\eta-s+2)^4 y(s)
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{6\Lambda} \sum_{s=2}^{\eta-2} \left[ -\Lambda(t-s+1)^3 + t^3(T-s+3)^3 - \frac{\alpha t^3}{4}(\eta-s+2)^4 \right] y(s) \\
 &\quad + \frac{1}{6\Lambda} \sum_{s=\eta-1}^{t-2} \left[ -\Lambda(t-s+1)^3 + t^3(T-s+3)^3 \right] y(s) \\
 &\quad + \frac{1}{6\Lambda} \sum_{s=t-1}^T t^3(T-s+3)^3 y(s) \\
 &= \sum_{s=2}^T G(t,s)y(s).
 \end{aligned}$$

Then the unique solution of problem (2.1)-(2.2) can be written as  $u(t) = \sum_{s=2}^T G(t,s)y(s)$ . The proof is complete.  $\square$

We observe that the condition  $0 < \alpha < \frac{4T(T+1)(T+2)}{(\eta-3)(\eta+2)(\eta^2-\eta+4)}$  implies  $G(t,s)$  is positive on  $\mathbb{N}_{2,T} \times \mathbb{N}_{2,T}$ .

Let

$$M_1 = \min \left\{ \frac{G(t,s)}{G(t,t)} : t \in \mathbb{N}_{2,T}, s \in \mathbb{N}_{2,T} \right\} \tag{2.8}$$

$$M_2 = \max \left\{ \frac{G(t,s)}{G(t,t)} : t \in \mathbb{N}_{T+2}, s \in \mathbb{N}_{2,T} \right\} \tag{2.9}$$

**Lemma 2.3.** *Let  $(t,s) \in \mathbb{N}_{2,T} \times \mathbb{N}_{2,T}$ . Then we have*

$$G(t,s) \geq M_1 G(t,t) \tag{2.10}$$

where  $0 < M_1 < 1$  is a constant given by

$$\begin{aligned}
 M_1 = \min \left\{ \frac{4(T-\eta+7)^3 - 4\Lambda\left(\frac{\eta-5}{\eta-2}\right) - \alpha\eta^4}{4(T+1)^3 - 24\alpha}, \frac{4(T-\eta+5)^3 - \alpha(\eta+1)^4}{4(T+1)^3 - 24\alpha}, \right. \\
 \left. \frac{6}{4(T+1)^3 - 24\alpha}, \frac{4(T-\eta-5)^3 - 4\Lambda\left(\frac{T-3}{T}\right) - \alpha\eta^4}{4(T-\eta+4)^3}, \right. \\
 \left. \frac{60(\eta-1)^3 - \Lambda(T-\eta+2)^3}{T^3(T-\eta+4)^3}, \frac{6}{(T-\eta+4)^3} \right\} \tag{2.11}
 \end{aligned}$$

**Proof.** In order that 2.10 holds, it is sufficient that  $M_1$  satisfies

$$M_1 \leq \min_{(t,s) \in \mathbb{N}_{2,T} \times \mathbb{N}_{2,T}} \frac{G(t,s)}{G(t,t)}. \tag{2.12}$$

Existence of positive solutions for summation boundary value problem ... 7

Then we may choose

$$M_1 \leq \min \left\{ \min_{(t,s) \in \mathbb{N}_{2,\eta-2} \times \mathbb{N}_{2,T}} \frac{G(t,s)}{G(t,t)}, \min_{(t,s) \in \mathbb{N}_{\eta-1,T} \times \mathbb{N}_{2,T}} \frac{G(t,s)}{G(t,t)} \right\}. \tag{2.13}$$

since

$$\begin{aligned} & \min_{(t,s) \in \mathbb{N}_{2,\eta-2} \times \mathbb{N}_{2,T}} \frac{G(t,s)}{G(t,t)} \\ &= \min_{t \in \mathbb{N}_{2,\eta-2}} \left\{ \min_{s \in \mathbb{N}_{2,t-2}} \frac{-\Lambda(t-s+1)^3 + t^3(T-s+3)^3 - \frac{\alpha t^3}{4}(\eta-s+2)^4}{t^3(T-t+3)^3 - \frac{\alpha t^3}{4}(\eta-t+2)^4}, \right. \\ & \quad \min_{s \in \mathbb{N}_{t-1,\eta-2}} \frac{t^3(T-s+3)^3 - \frac{\alpha t^3}{4}(\eta-s+2)^4}{t^3(T-t+3)^3 - \frac{\alpha t^3}{4}(\eta-t+2)^4}, \\ & \quad \left. \min_{s \in \mathbb{N}_{\eta-1,T}} \frac{t^3(T-s+3)^3}{t^3(T-t+3)^3 - \frac{\alpha t^3}{4}(\eta-t+2)^4} \right\} \\ &\geq \min_{t \in \mathbb{N}_{2,\eta-2}} \left\{ \frac{-\Lambda(t-1)^3 + t^3(T-t+5)^3 - \frac{\alpha t^3}{4}\eta^4}{t^3(T-t+3)^3 - \frac{\alpha t^3}{4}(\eta-t+2)^4}, \frac{(T-\eta+5)^3 - \frac{\alpha}{4}(\eta-t+3)^4}{(T-t+3)^3 - \frac{\alpha}{4}(\eta-t+2)^4}, \right. \\ & \quad \left. \frac{3^3}{(T-t+3)^3 - \frac{\alpha}{4}(\eta-t+2)^4} \right\} \\ &\geq \min \left\{ \frac{-\Lambda \left(1 - \frac{3}{\eta-2}\right) + (T-\eta+7)^3 - \frac{\alpha}{4}(\eta)^4}{(T+1)^3 - \frac{\alpha 4^4}{4}}, \frac{(T-\eta+5)^3 - \frac{\alpha}{4}(\eta+1)^4}{(T+1)^3 - \frac{\alpha 4^4}{4}}, \frac{6}{(T+1)^3 - \frac{\alpha 4^4}{4}} \right\} \\ &= \min \left\{ \frac{4(T-\eta+7)^3 - 4\Lambda \left(\frac{\eta-5}{\eta-2}\right) - \alpha\eta^4}{4(T+1)^3 - 24\alpha}, \frac{4(T-\eta+5)^3 - \alpha(\eta+1)^4}{4(T+1)^3 - 24\alpha}, \frac{6}{4(T+1)^3 - 24\alpha} \right\} \tag{2.14} \end{aligned}$$

Similarly, we get

$$\begin{aligned} & \min_{(t,s) \in \mathbb{N}_{\eta-1,T} \times \mathbb{N}_{2,T}} \frac{G(t,s)}{G(t,t)} \\ &\geq \min \left\{ \frac{4(T-\eta-5)^3 - 4\Lambda \left(\frac{T-3}{T}\right) - \alpha\eta^4}{4(T-\eta+4)^3}, \frac{60(\eta-1)^3 - \Lambda(T-\eta+2)^3}{T^3(T-\eta+4)^3}, \frac{6}{(T-\eta+4)^3} \right\} \tag{2.15} \end{aligned}$$

The (2.11) is immediate from (2.14)-(2.15) □

**Lemma 2.4.** *Let  $(t,s) \in \mathbb{N}_{T+2} \times \mathbb{N}_{2,T}$ . Then we have*

$$G(t,s) \leq M_2 G(t,t) \tag{2.16}$$



where  $M_2 \geq 1$  is a constant given by

$$M_2 = \max \left\{ \frac{24\Lambda + (\eta - 2)^3(T + 1)^3}{\alpha(\eta - 2)^3\eta^4}, \frac{4(T - \eta + 6)^3 - 24\alpha}{4(T - \eta + 6)^3 - \alpha\eta^4}, \frac{4(T - \eta + 4)^3}{4(T - \eta + 5)^3 - \alpha\eta^4}, \frac{4T^3 - 24\alpha(\eta - 1)^3}{6(\eta - 1)^3}, \frac{T^3(T - \eta + 4)^3 - 6\Lambda}{6(\eta - 1)^3}, \frac{(T - \eta + 5)^3}{6} \right\} \quad (2.17)$$

**Proof.** For  $t = 0, 1$ , from (2.6) we get

$$G(0, s) = G(0, 0) = 0; \quad G(1, s) = G(1, 1) = 0.$$

Then we may choose  $M_2 = 1$ . For  $t \in \mathbb{N}_{2,T}$ , if 2.16 holds, it is sufficient that  $M_2$  satisfies

$$M_2 \geq \max_{(t,s) \in \mathbb{N}_{2,T} \times \mathbb{N}_{2,T}} \frac{G(t, s)}{G(t, t)}. \quad (2.18)$$

Then we may choose

$$M_2 \geq \max \left\{ \max_{(t,s) \in \mathbb{N}_{2,\eta-2} \times \mathbb{N}_{2,T}} \frac{G(t, s)}{G(t, t)}, \max_{(t,s) \in \mathbb{N}_{\eta-1,T} \times \mathbb{N}_{2,T}} \frac{G(t, s)}{G(t, t)} \right\}. \quad (2.19)$$

since

$$\begin{aligned} & \max_{(t,s) \in \mathbb{N}_{2,\eta-2} \times \mathbb{N}_{2,T}} \frac{G(t, s)}{G(t, t)} \\ &= \max_{t \in \mathbb{N}_{2,\eta-2}} \left\{ \max_{s \in \mathbb{N}_{2,t-2}} \frac{-\Lambda(t - s + 1)^3 + t^3(T - s + 3)^3 - \frac{\alpha t^3}{4}(\eta - s + 2)^4}{t^3(T - t + 3)^3 - \frac{\alpha t^3}{4}(\eta - t + 2)^4}, \right. \\ & \quad \max_{s \in \mathbb{N}_{t-1,\eta-2}} \frac{t^3(T - s + 3)^3 - \frac{\alpha t^3}{4}(\eta - s + 2)^4}{t^3(T - t + 3)^3 - \frac{\alpha t^3}{4}(\eta - t + 2)^4}, \\ & \quad \left. \max_{s \in \mathbb{N}_{\eta-1,T}} \frac{t^3(T - s + 3)^3}{t^3(T - t + 3)^3 - \frac{\alpha t^3}{4}(\eta - t + 2)^4} \right\} \\ &\leq \max_{t \in \mathbb{N}_{2,\eta-2}} \left\{ \frac{-\Lambda 3^3 + t^3(T + 1)^3 - \frac{\alpha t^3}{4}(\eta - t + 4)^4}{t^3(T - t + 3)^3 - \frac{\alpha t^3}{4}(\eta - t + 2)^4}, \frac{(T - t + 4)^3 - \frac{\alpha 4^4}{4}}{(T - t + 3)^3 - \frac{\alpha}{4}(\eta - t + 2)^4}, \right. \\ & \quad \left. \frac{(T - \eta + 4)^3}{(T - t + 3)^3 - \frac{\alpha}{4}\eta^4} \right\} \\ &\leq \max \left\{ \frac{24\Lambda + (\eta - 2)^3(T + 1)^3}{\alpha(\eta - 2)^3\eta^4}, \frac{4(T - \eta + 6)^3 - 24\alpha}{4(T - \eta + 6)^3 - \alpha\eta^4}, \frac{4(T - \eta + 4)^3}{4(T - \eta + 5)^3 - \alpha\eta^4} \right\} \quad (2.20) \end{aligned}$$

Similarly, we get

$$\begin{aligned} & \max_{(t,s) \in \mathbb{N}_{\eta-1,T} \times \mathbb{N}_{2,T}} \frac{G(t, s)}{G(t, t)} \\ &\leq \max \left\{ \frac{4T^3 - 24\alpha(\eta - 1)^3}{6(\eta - 1)^3}, \frac{T^3(T - \eta + 4)^3 - 6\Lambda}{6(\eta - 1)^3}, \frac{(T - \eta + 5)^3}{6} \right\} \quad (2.21) \end{aligned}$$

For  $t = T + 1, T + 2$  from (2.6) we get

$$\begin{aligned} G(T + 1, s) &< \frac{1}{6\Lambda} \left[ -\Lambda(T - s + 2)^3 + (T + 1)^3 \left( (T - s + 3)^3 - \frac{\alpha}{4}(\eta - s + 2)^4 \right) \right] \\ &= -\frac{\alpha}{24\Lambda} (T + 1)^3 (\eta - s + 2)^4 \\ &< -\frac{\alpha}{24\Lambda} (T + 1)^3 (\eta - T + 1)^4 \\ &= G(T + 1, T + 1), \end{aligned}$$

$$\begin{aligned} G(T + 2, s) &< \frac{1}{6\Lambda} \left[ -\Lambda(T - s + 3)^3 + (T + 2)^3 \left( (T - s + 3)^3 - \frac{\alpha}{4}(\eta - s + 2)^4 \right) \right] \\ &= \frac{1}{6\Lambda} \left[ (T - s + 2)^3 \left( \frac{T - s + 3}{T - s} \right) \left( (T + 2)^3 - \Lambda \right) - \frac{\alpha}{4}(\eta - s + 2)^4 \right] \\ &= -\frac{\alpha}{24\Lambda} (T + 2)^3 (\eta - s + 2)^4 \\ &< -\frac{\alpha}{24\Lambda} (T + 2)^3 (\eta - T)^4 \\ &= G(T + 2, T + 2). \end{aligned}$$

Then we choose  $M_2 = 1$ . So (2.17) is immediate from (2.20)-(2.21). □

### 3 Main Results

Now we are in the position to establish the main result.

**Theorem 3.1.** *Assume (H1) - (H3) hold. Then the problem (1.1)-(1.2) has at least one positive solution.*

**Proof.** In the following, we denote

$$m = \min_{t \in \mathbb{N}_{\eta-1, T}} G(t, t), \quad M = \max_{t \in \mathbb{N}_{T+2}} G(t, t).$$

Then  $0 < m < M$ .

Let  $E$  be the Banach's space defined by  $E = \{u : \mathbb{N}_{T+2} \rightarrow R\}$ . Define

$$K = \{u \in E : u \geq 0, t \in \mathbb{N}_{T+2} \text{ and } \min_{t \in \mathbb{N}_{2, T}} u(t) \geq \sigma \|u\|\}.$$

where  $\sigma = \frac{M_1 m}{M_2 M} \in (0, 1)$ ,  $\|u\| = \max_{t \in \mathbb{N}_{T+2}} |u(t)|$ . It is obvious that  $K$  is a cone in  $E$ .

We define the operator  $A : K \rightarrow E$  by

$$(Au)(t) = \sum_{s=2}^T G(t, s)a(s)f(u(s)), t \in \mathbb{N}_{T+2}.$$

It is clear that problem (1.1)-(1.2) has a solution  $u$  if and only if  $u \in K$  is a fixed point of operator  $A$ . We shall now show that the operator  $A$  maps  $K$  to itself. For this, let  $u \in K$ , from  $(H_2) - (H_3)$ , we get

$$(Au)(t) = \sum_{s=1}^T G(t, s)a(s)f(u(s)) \geq 0, t \in \mathbb{N}_{T+2}. \tag{3.1}$$

from (2.9), we obtain

$$\begin{aligned} (Au)(t) &= \sum_{s=2}^T G(t, s)a(s)f(u(s)) \leq M_2 \sum_{s=2}^T G(t, t)a(s)f(u(s)) \\ &\leq M_2M \sum_{s=2}^T a(s)f(u(s)), \quad t \in \mathbb{N}_{T+2}. \end{aligned}$$

Therefore

$$\| Au \| \leq M_2M \sum_{s=2}^T a(s)f(u(s)). \tag{3.2}$$

Now from  $(H_2), (H_3), (2.8)$  and (3.2), for  $t \in \mathbb{N}_{\eta, T}$ , we have

$$\begin{aligned} (Au)(t) &\geq M_1 \sum_{s=2}^T G(t, t)a(s)f(u(s)) \geq M_1m \sum_{s=2}^T a(s)f(u(s)) \\ &\geq \frac{M_1m}{M_2M} \| Au \| = \sigma \| u \|. \end{aligned}$$

Then

$$\min_{t \in \mathbb{N}_{\eta, T}} (Au)(t) \geq \sigma \| u \|. \tag{3.3}$$

From (3.1)-(3.2), we obtain  $Au \in K$ , Hence  $A(K) \subseteq K$ . So  $F : k \rightarrow K$  is completely continuous.

**Superlinear case.**  $f_0 = 0$  and  $f_\infty = \infty$ . Since  $f_0 = 0$ , we may choose  $H_1 > 0$  so that  $f(u) \leq \epsilon_1 u$ , for  $0 < u \leq H_1$ , where  $\epsilon_1 > 0$  satisfies

$$\epsilon_1 M_2M \sum_{s=2}^T a(s) \leq 1. \tag{3.4}$$

Thus, if we let

$$\Omega_1 = \{u \in E : \|u\| < H_1\},$$

then for  $u \in K \cap \partial\Omega_1$ , we get

$$\begin{aligned} (Au)(t) &\leq M_2 \sum_{s=2}^T G(t, t)a(s)f(u(s)) \leq \epsilon_1 M_2 M \sum_{s=2}^T a(s)u(s) \\ &\leq \epsilon_1 M_2 M \sum_{s=2}^T a(s)\|u\| \leq \|u\|. \end{aligned}$$

Thus  $\|Au\| \leq \|u\|$ ,  $u \in K \cap \partial\Omega_1$ .

Further, since  $f_\infty = \infty$ , there exists  $\widehat{H}_2 > 0$  such that  $f(u) \geq \epsilon_2 u$ , for  $u \geq \widehat{H}_2$ , where  $\epsilon_2 > 0$  satisfies

$$\epsilon_2 M_1 \sigma \sum_{s=\eta-1}^T G(\eta-1, \eta-1)a(s) \geq 1. \tag{3.5}$$

Let  $H_2 = \max\{2H_1, \frac{\widehat{H}_2}{\sigma}\}$  and  $\Omega_2 = \{u \in E : \|u\| < H_2\}$ . Then  $u \in K \cap \partial\Omega_2$  implies

$$\min_{t \in \mathbb{N}_{\eta-1, T}} u(t) \geq \sigma \|u\| \geq \widehat{H}_2.$$

Applying (2.8) and (3.5), we get

$$\begin{aligned} (Au)(\eta-1) &= M_1 \sum_{s=2}^T G(\eta-1, s)a(s)f(u(s)) \geq M_1 \sum_{s=\eta-1}^T G(\eta-1, \eta-1)a(s)f(u(s)) \\ &\geq \epsilon_2 M_1 \sum_{s=\eta-1}^T G(\eta-1, \eta-1)a(s)y(s) \geq \epsilon_2 M_1 \sigma \sum_{s=\eta-1}^T G(\eta-1, \eta-1)a(s)\|u\| \\ &\geq \|u\|. \end{aligned}$$

Hence,  $\|Au\| \geq \|u\|$ ,  $u \in K \cap \partial\Omega_2$ . By the first part of Theorem 1.1,  $A$  has a fixed point in  $K \cap (\overline{\Omega}_2 \setminus \Omega_1)$  such that  $H_1 \leq \|u\| \leq H_2$ .

**Sublinear case.**  $f_0 = \infty$  and  $f_\infty = 0$ . Since  $f_0 = \infty$ , choose  $H_3 > 0$  such that  $f(u) \geq \epsilon_3 u$  for  $0 < u \leq H_3$ , where  $\epsilon_3 > 0$  satisfies

$$\epsilon_3 M_1 \sigma \sum_{s=\eta-1}^T G(\eta-1, \eta-1)a(s) \geq 1. \tag{3.6}$$

Let

$$\Omega_3 = \{u \in E : \|u\| < H_3\},$$

then for  $u \in K \cap \partial\Omega_3$ , we get

$$\begin{aligned} (Au)(\eta - 1) &\geq M_1 \sum_{s=\eta-1}^T G(\eta - 1, \eta - 1)a(s)f(u(s)) \geq \epsilon_3 M_1 \sum_{s=\eta-1}^T G(\eta - 1, \eta - 1)a(s)y(s) \\ &\geq \epsilon_3 M_1 \sigma \sum_{s=\eta-1}^T G(\eta - 1, \eta - 1)a(s)\|u\| \geq \|u\|. \end{aligned}$$

Thus,  $\|Au\| \geq \|u\|$ ,  $u \in K \cap \partial\Omega_3$ .

Now, since  $f_\infty = 0$ , there exists  $\widehat{H}_4 > 0$  so that  $f(u) \leq \epsilon_4 u$  for  $u \geq \widehat{H}_4$ , where  $\epsilon_4 > 0$  satisfies

$$\epsilon_4 M_2 M \sum_{s=\eta-1}^T a(s) \geq 1. \tag{3.7}$$

**Subcase 1.** Suppose  $f$  is bounded,  $f(u) \leq L$  for all  $u \in [0, \infty)$  for some  $L > 0$ .

Let  $H_4 = \max\{2H_3, LM_2M \sum_{s=1}^T a(s)\}$ .

Then for  $u \in K$  and  $\|u\| = H_4$ , we get

$$\begin{aligned} (Au)(\eta) &\leq M_2 \sum_{s=2}^T G(t, t)a(s)f(u(s)) \leq LM_2M \sum_{s=2}^T a(s) \\ &\leq H_4 = \|u\| \end{aligned}$$

Thus  $(Au)(t) \leq \|u\|$ .

**Subcase 2.** Suppose  $f$  is unbounded, there exist  $H_4 > \max\{2H_3, \frac{\widehat{H}_4}{\sigma}\}$  such that  $f(u) \leq f(H_4)$  for all  $0 < u \leq H_4$ . Then for  $u \in K$  with  $\|u\| = H_4$  from (2.9) and (3.7), we have

$$\begin{aligned} (Au)(t) &\leq M_2 \sum_{s=2}^T G(t, t)a(s)f(u(s)) \leq M_2M \sum_{s=2}^T a(s)f(H_4) \\ &\leq \epsilon_4 M_2 M \sum_{s=2}^T a(s)H_4 \leq H_4 = \|u\|. \end{aligned}$$

Thus in both cases, we may put  $\Omega_4 = \{u \in E : \|u\| < H_4\}$ . Then

$$\|Au\| \leq \|u\|, u \in K \cap \partial\Omega_4.$$

By the second part of Theorem 1.1,  $A$  has a fixed point  $u$  in  $K \cap (\overline{\Omega}_4 \setminus \Omega_3)$ , such that  $H_3 \leq \|u\| \leq H_4$ . This completes the sublinear part of the theorem. Therefore, the problem (1.1)-(1.2) has at least one positive solution.  $\square$

### 4 Some examples

In this section, in order to illustrate our result, we consider some examples.

**Example 4.1** Consider the BVP

$$\Delta^4 u(t - 2) + t^2 u^k = 0, \quad t \in N_{2,6}, \tag{4.1}$$

$$u(0) = \Delta u(0) = \Delta^2 u(0) = 0, \quad u(8) = \frac{2}{3} \sum_{s=4}^5 u(s). \tag{4.2}$$

Set  $\alpha = \frac{2}{3}$ ,  $\eta = 5$ ,  $T = 6$ ,  $a(t) = t^2$ ,  $f(u) = u^k$ .

We can show that

$$T(T + 1)(T + 2) - \frac{\alpha}{4}(\eta - 3)(\eta + 2)(\eta^2 - \eta + 4) = 280 > 0.$$

Case I :  $k \in (1, \infty)$ . In this case,  $f_0 = 0$ ,  $f_\infty = \infty$  and (i) of theorem 3.1 holds. Then BVP (4.1)-(4.2) has at least one positive solution.

Case II :  $k \in (0, 1)$ . In this case,  $f_0 = \infty$ ,  $f_\infty = 0$  and (ii) of theorem 3.1 holds. Then BVP (4.1)-(4.2) has at least one positive solution.

**Example 4.2** Consider the BVP

$$\Delta^4 u(t - 2) + e^t t^e \left( \frac{\pi \sin u + 2 \cos u}{u^2} \right) = 0, \quad t \in N_{2,8}, \tag{4.3}$$

$$u(0) = \Delta u(0) = \Delta^2 u(0) = 0, \quad u(10) = \frac{1}{3} \sum_{s=4}^6 u(s), \tag{4.4}$$

Set  $\alpha = \frac{1}{3}$ ,  $\eta = 6$ ,  $T = 8$ ,  $a(t) = e^t t^e$ ,  $f(u) = \frac{\pi \sin u + 2 \cos u}{u^2}$ .

We can show that

$$T(T + 1)(T + 2) - \frac{\alpha}{4}(\eta - 3)(\eta + 2)(\eta^2 - \eta + 4) = 596 > 0.$$

Through a simple calculation we can get  $f_0 = \infty$ ,  $f_\infty = 0$ . Thus, by (ii) of theorem 3.1, we can get BVP (4.3)-(4.4) has at least one positive solution.

**Acknowledgement(s)** : This research (KMUTNB-GOV-57-07) is supported by King Mongkut’s University of Technology North Bangkok, Thailand.

## References

- [1] R.P. Agarwal, Focal Boundary Value Problems for Differential and Difference Equations, Kluwer Academic Publishers, Dordrecht, 1998.
- [2] R.P. Agarwal, D. O'Regan, P.J.Y. Wong, Positive Solutions of Differential, Difference and Integral Equations, Kluwer Academic Publishers, Dordrecht, 1999.
- [3] M.A. Krasnoselskii, Positive Solutions of Operator Equations, Noordhoof, Groningen, 1964.
- [4] R.J. Liang, Y.H. Zhao, J.P. Sun, A new theorem of existence to fourth-order boundary value problem, International Journal of Differential Equations and Applications, 7 (2003) 257-262
- [5] R.Ma, Y. Xu, Existence of positive solution for nonlinear fourth-order difference equations, Computers and Mathematics with Applications 59 (2010) 3770-3777.
- [6] Z.M. He, J.S. Yu, On the existence of positive solutions of fourth-order difference equations, Appl. Math. Comput. 161 (2005) 139-148.
- [7] B.G. Zhang, L.J. Kong, Y.J. Sun, X.H. Deng, Existence of positive solutions for BVPs of fourth-order difference equations, Appl. Math. Comput. 131 (2002). 583-591.
- [8] Q.L. Yao, Existence, multiplicity and infinite solvability of positive solutions to a nonlinear fourth-order periodic boundary value problem, Nonlinear Anal. 63 (2005) 237-246.
- [9] F.Y. Li, Q. Zhang, Z.P. Liang, Existence and multiplicity of solutions of a kind of fourth-order boundary value problem, Nonlinear Anal. 62 (2005). 803-816.
- [10] X.L. Liu, W.T. Li, Existence and multiplicity of solutions for fourth-order boundary value problems with parameters, J. Math. Anal. Appl. 327 (2007). 362-375.
- [11] Y. Yang, J.H. Zhang, Existence of solutions for some fourth-order boundary value problems with parameters, Nonlinear Anal. 69 (2008) 136-1375.
- [12] G.D. Han, Z.B. Xu, Multiple solutions of some nonlinear fourth-order beam equations, Nonlinear Anal. 68 (2008) 3646-3656.

Existence of positive solutions for summation boundary value problem ... 15

- [13] X. Lin, W. Lin, Three positive solutions of a second order difference Equations with Three-Point Boundary Value Problem, *J.Appl. Math. Comput.* 31(2009), 279-288.
- [14] J.Reunsumrit, T. Sitthiwirattam, Positive solutions of difference-summation boundary value problem for a second-order difference equation, *Journal of Computational Analysis and Applications.* (In press)
- [15] T. Sitthiwirattam, J. Tariboon, Positive Solutions to a Generalized Second Order Difference Equation with Summation Boundary Value Problem. *Journal of Applied Mathematics.* Vol.2012, Article ID 569313, 15 pages.



# Similarity measure between generalized intuitionistic fuzzy sets and its application to pattern recognition

Jin Han Park\*

Department of Applied Mathematics, Pukyong National University  
Busan 608-737, South Korea  
jihpark@pknu.ac.kr

Jongchul Hwang, Juhyung Kim, Byeongmuk Park,  
Juyoung Park, Jeongwoo Son, Sihun Lee  
Busanil Science High School, Busan 660-756, South Korea  
oecuo@empal.com (J. Hwang), rlawngudzkzk@naver.com (J. Kim),  
qudanr50@naver.com (B. Park), gabby24@naver.com (J.Park),  
rodeh1945@naver.com (J. Son), pencil98@naver.com (S. Lee)

## Abstract

This paper presents new methods for measuring similarity between generalized intuitionistic fuzzy sets (GIFSs) and its application to pattern recognition. Firstly, the geometrical interpretation of GIFSs is carefully reviewed and then the results of the interpretation is utilized to generate new methods for measuring similarity in order to calculate the degree of similarity between GIFSs. Numerical example is given to illustrate the application of the proposed similarity measures. Finally, we also use the proposed similarity measures to characterize the similarity between linguistic variables.

## 1 Introduction

As a generalization of fuzzy sets, intuitionistic fuzzy sets (IFSs) were presented by Atanassov [1, 2, 3]. Since IFSs can present the degrees of membership and non-membership with a degree of hesitancy, the knowledge and semantic representation become more meaningful and applicable. These IFSs have been widely studied and applied in various areas, such as logic programming [4], decision making [7, 22], pattern recognition [12, 14, 15, 16, 19, 26] and medical diagnosis

---

\*Corresponding author: jihpark@pknu.ac.kr (J.H. Park)  
This work was supported by a Research Grant of Pukyong National University (2014).

[9, 25], and seem to have more popular than fuzzy sets technology. Mondal and Samanta [18] introduced generalized intuitionistic fuzzy sets (GIFSs) as a generalization of IFSs and studied their basic properties. Park et al. [20] proposed a method to calculate the correlation coefficient of GIFSs. There is a little investigation on GIFSs. Similarity assessment plays a fundamental and important role in inference and approximate reasoning in all applications of intuitionistic fuzzy logic [4]. For different purposes different similarity measures should be used. Based on the importance of the problem, the effectiveness and properties of the different similarity measures for IFSs have been compared and examined by many researchers (e.g. Hung and Yang [12], Li and Cheng [14], Li et al. [15], Liang and Shi [16], Mitchell [19], Szmidt and Baldwin [23]). The analysis of similarity is also a fundamental issue while employing GIFSs. Recently, Park et al. [21] proposed and applied similarity measure to compare generalized intuitionistic fuzzy preferences given by individuals (experts) and evaluated an extent of a group agreement. In this paper, we propose new similarity measures based on the geometrical representation for GIFS. The proposed similarity measures depend on the triplet, membership degree, nonmembership degree, and hesitation margin. This paper proves that the proposed similarity measures satisfy the properties of axiomatic definition for similarity measures. Numerical example is given to illustrate the application of the developed similarity measures. Furthermore, we use the proposed similarity measures to characterize the similarity between linguistic variables.

## 2 Brief introduction of GIFSs

In the following, we firstly recall basic notions and definitions of GIFSs which can be found in [18].

Let  $X$  be the universe of discourse. A generalized intuitionistic fuzzy set (GIFS)  $A$  in  $X$  is an object having the form

$$A = \{(x, \mu_A(x), \nu_A(x)) \mid x \in X\} \tag{1}$$

where  $\mu_A, \nu_A : X \rightarrow [0, 1]$  denote membership function and non-membership function, respectively, of  $A$  and satisfy  $\min\{\mu_A(x), \nu_A(x)\} \leq 0.5$  for all  $x \in X$ . Let  $\text{GIFS}(X)$  denote the set of all GIFSs in  $X$ .

For an IFS  $A = \{(x, \mu_A(x), \nu_A(x)) \mid x \in X\}$ , it is observed that  $\mu_A(x) + \nu_A(x) \leq 1$  implies  $\min\{\mu_A(x), \nu_A(x)\} \leq 0.5$  for each  $x \in X$ . Thus, every IFS is GIFS.

For each GIFS  $A$  in  $X$ , we call

$$\phi_A(x) = 1 - \mu_A(x) - \nu_A(x) \tag{2}$$

a generalized intuitionistic fuzzy index (or a hesitation margin) of  $x$  in  $A$  and it expresses a lack/excess of knowledge of whether  $x$  belongs to  $A$  or not. (see, [20]). It is obvious that  $-0.5 \leq \phi_A(x) \leq 1$  for each  $x \in X$ .

Having in mind that for each element  $x$  belonging to a GIFS  $A$ , the values of membership, non-membership and generalized intuitionistic fuzzy index add up to one, i.e.

$$\mu_A(x) + \nu_A(x) + \phi_A(x) = 1 \tag{3}$$

and that each of the membership and non-membership are from  $[0, 1]$  and the generalized intuitionistic fuzzy index is from  $[-0.5, 1]$ , we can imagine a cuboid (Figure 1) inside which there is a polygon  $ADBEGF$  where the above equation is fulfilled. In other words, the polygon  $ADBEGF$  represents a surface where coordinates of any element belonging to a GIFS can be represented. Each point belonging to the polygon  $ADBEGF$  is described via three coordinates:  $(\mu, \nu, \phi)$ . Points  $A$  and  $B$  represent crisp elements. Point  $A(1, 0, 0)$  represents elements fully belonging to a GIFS as  $\mu = 1$ . Point  $B(0, 1, 0)$  represents elements fully not belonging to a GIFS as  $\nu = 1$ . Point  $D(0, 0, 1)$  represents element about which we are not able to say if they belong or not belong to a GIFS (generalized intuitionistic fuzzy index  $\phi = 1$ ). Point  $E(1, 0.5, -0.5)$  represents element about which we can say to belong to a GIFS ( $\phi = -0.5$ ). Point  $F(0.5, 1, -0.5)$  represents element about which we can say to not belong to a GIFS ( $\phi = -0.5$ ). Such an interpretation is intuitively appealing and provides means for the representation of many aspects of imperfect information. Segment  $AB$  (where  $\phi = 0$ ) represents elements belonging to classical fuzzy sets ( $\mu + \nu = 1$ ). Triangle  $ADB$  (where  $0 \leq \phi \leq 1$ ) represents elements belonging to IFSs ( $0 \leq \mu + \nu \leq 1$ ). Any other combination of the values characterizing a GIFS can be represented inside the triangles  $AGF$  and  $BEG$ . In other words, each element belonging to a GIFS can be represented as a point  $(\mu, \nu, \phi)$  belonging to the polygon  $ADBEGF$  (cf. Figure 1).




Figure 1: A geometrical interpretation of a GIFS

It is worth mentioning that the geometrical interpretation is directly related to the definition of a GIFS, and it does not need any additional assumptions. By employing the above geometrical representation, a GIFS  $A$  can be expressed as

$$A = \{(\mu_A(x), \nu_A(x), \phi_A(x)) \mid x \in X\}. \tag{4}$$

Therefore, this representation of a GIFS will be a point of departure for considering our method in calculating the degree of similarity between GIFSs.

For  $A, B \in \text{GIFS}(X)$ , Mondal and Samanta [18] defined the notion of containment as follows:

$$A \subseteq B \Leftrightarrow \mu_A(x) \leq \mu_B(x) \text{ and } \nu_A(x) \geq \nu_B(x) \forall x \in X. \tag{5}$$

As above-mentioned, we can not omit the third parameter (hesitancy degree) in the representation of GIFSs and then redefine the notion of containment as follows:

$$A \subseteq B \Leftrightarrow \mu_A(x) \leq \mu_B(x), \nu_A(x) \geq \nu_B(x) \text{ and } \phi_A(x) \geq \phi_B(x) \forall x \in X. \tag{6}$$

**Definition 2.1** Let  $S : \text{GIFS}(X) \times \text{GIFS}(X) \rightarrow [0, 1]$  be a mapping.  $S(A, B)$  is said to be the degree of similarity between  $A \in \text{GIFS}(X)$  and  $B \in \text{GIFS}(X)$  if  $S(A, B)$  satisfies the properties (SP1)-(SP4):

- (SP1)  $0 \leq S(A, B) \leq 1$ ;
- (SP2)  $S(A, B) = 1$  if and only if  $A = B$ ;
- (SP3)  $S(A, B) = S(B, A)$ ;
- (SP4)  $S(A, C) \leq S(A, B)$  and  $S(A, C) \leq S(B, C)$  if  $A \subseteq B \subseteq C, A, B, C \in \text{GIFS}(X)$ .

**Definition 2.2** Let  $D : \text{GIFS}(X) \times \text{GIFS}(X) \rightarrow [0, 1]$  be a mapping.  $D(A, B)$  is called a distance  $A \in \text{GIFS}(X)$  and  $B \in \text{GIFS}(X)$  if  $D(A, B)$  satisfies the properties (DP1)-(DP4):

- (DP1)  $0 \leq D(A, B) \leq 1$ ;
- (DP2)  $D(A, B) = 0$  if and only if  $A = B$ ;
- (DP3)  $D(A, B) = D(B, A)$ ;
- (DP4)  $D(A, B) \leq D(A, C)$  and  $D(B, C) \leq D(A, C)$  if  $A \subseteq B \subseteq C, A, B, C \in \text{GIFS}(X)$ .

Because distance and similarity measures are complementary concepts, similarity measures can be used to define distance measures and vice verses.

### 3 New similarity measures between GIFSs

In this section, we take into account three parameters describing GIFSs to propose a new similarity measures between GIFSs based on the geometrical representation of GIFSs.

Let  $A = \{(x, \mu_A(x), \nu_A(x)) \mid x \in X\}$  and  $B = \{(x, \mu_B(x), \nu_B(x)) \mid x \in X\}$  be two GIFSs in  $X = \{x_1, x_2, \dots, x_n\}$ . We propose a new similarity measure:

$$S_g(A, B) = 1 - \frac{1}{n} \sum_{i=1}^n \left( \frac{|\mu_A(x_i) - \mu_B(x_i)| + |\nu_A(x_i) - \nu_B(x_i)| + |\phi_A(x_i) - \phi_B(x_i)|}{4} + \frac{\max(|\mu_A(x_i) - \mu_B(x_i)|, |\nu_A(x_i) - \nu_B(x_i)|, |\phi_A(x_i) - \phi_B(x_i)|)}{2} \right), \quad (7)$$

where  $\phi_A(x_i)$  and  $\phi_B(x_i)$  are, respectively, the hesitancy degree of the element  $x_i \in X$  to the sets  $A$  and  $B$ .

**Theorem 3.1**  $S_g(A, B)$  is the similarity measure between two GIFSs  $A$  and  $B$ .

**Proof** For the sake of simplicity, IFSs  $A$  and  $B$  are denoted by  $A = \{(\mu_A(x_i), \nu_A(x_i), \phi(x_i)) \mid x_i \in X\}$  and  $B = \{(\mu_B(x_i), \nu_B(x_i), \phi_B(x_i)) \mid x_i \in X\}$ , respectively. Obviously,  $S_g(A, B)$  satisfies (SP1) and (SP3) of Definition 1. We only need to prove that  $S_g(A, B)$  satisfies (SP2) and (SP4).

(SP2): From (6), we have

$$\begin{aligned} S_g(A, B) &= 1 \\ &\Leftrightarrow \mu_A(x_i) = \mu_B(x_i), \nu_A(x_i) = \nu_B(x_i), \phi_A(x_i) = \phi_B(x_i), \forall x_i \in X \\ &\Leftrightarrow A = B. \end{aligned}$$

(SP4): For any IFS  $C = \{(\mu_C(x_i), \nu_C(x_i), \phi_C(x_i)) \mid x_i \in X\}$ , if  $A \subseteq B \subseteq C$ , then we have

$$S_g(A, C) = 1 - \frac{1}{n} \sum_{i=1}^n \left( \frac{|\mu_A(x_i) - \mu_C(x_i)| + |\nu_A(x_i) - \nu_C(x_i)| + |\phi_A(x_i) - \phi_C(x_i)|}{4} + \frac{\max(|\mu_A(x_i) - \mu_C(x_i)|, |\nu_A(x_i) - \nu_C(x_i)|, |\phi_A(x_i) - \phi_C(x_i)|)}{2} \right).$$

It is easy to see that

$$\begin{aligned} |\mu_A(x_i) - \mu_C(x_i)| &\geq |\mu_A(x_i) - \mu_B(x_i)|, |\nu_A(x_i) - \nu_C(x_i)| \geq |\nu_A(x_i) - \nu_B(x_i)|, \\ |\phi_A(x_i) - \phi_C(x_i)| &\geq |\phi_A(x_i) - \phi_B(x_i)|. \end{aligned}$$

So we have

$$\begin{aligned} &\frac{|\mu_A(x_i) - \mu_C(x_i)| + |\nu_A(x_i) - \nu_C(x_i)| + |\phi_A(x_i) - \phi_C(x_i)|}{4} \\ &\quad + \frac{\max(|\mu_A(x_i) - \mu_C(x_i)|, |\nu_A(x_i) - \nu_C(x_i)|, |\phi_A(x_i) - \phi_C(x_i)|)}{2} \\ &\geq \frac{|\mu_A(x_i) - \mu_B(x_i)| + |\nu_A(x_i) - \nu_B(x_i)| + |\phi_A(x_i) - \phi_B(x_i)|}{4} \\ &\quad + \frac{\max(|\mu_A(x_i) - \mu_B(x_i)|, |\nu_A(x_i) - \nu_B(x_i)|, |\phi_A(x_i) - \phi_B(x_i)|)}{2} \end{aligned}$$

and thus we get  $S_g(A, C) \leq S_g(A, B)$ . By the same reason, we can get  $S_g(A, C) \leq S_g(B, C)$ .

However, the elements in the universe may have different importance in pattern recognition. We should consider the weight of the elements so that we can obtain more reasonable results in pattern recognition.

Assume that the weight of  $x_i$  in  $X$  is  $w_i$ , where  $w_i \in [0, 1]$  ( $i = 1, 2, \dots, n$ ) and  $\sum_{i=1}^n w_i = 1$ . The similarity measure between GIFSs  $A$  and  $B$  can be obtained by the following form:

$$S_{gw}(A, B) = 1 - \sum_{i=1}^n w_i \left( \frac{|\mu_A(x_i) - \mu_B(x_i)| + |\nu_A(x_i) - \nu_B(x_i)| + |\phi_A(x_i) - \phi_B(x_i)|}{4} + \frac{\max(|\mu_A(x_i) - \mu_B(x_i)|, |\nu_A(x_i) - \nu_B(x_i)|, |\phi_A(x_i) - \phi_B(x_i)|)}{2} \right). \quad (8)$$

Likewise, for  $S_{gw}(A, B)$ , the following theorem holds.

**Theorem 3.2**  $S_{gw}(A, B)$  is the similarity measure between two GIFSs  $A$  and  $B$ .

**Proof** The proof is similar to that of Theorem 3.1.

**Remark 3.3** Obviously, if  $w_i = 1/n$  ( $i = 1, 2, \dots, n$ ), (8) becomes (7). So, (7) is a special case of (8).

Now, we propose another new similarity measure between GIFSs  $A = \{(x, \mu_A(x), \nu_A(x)) | x \in X\}$  and  $B = \{(x, \mu_B(x), \nu_B(x)) | x \in X\}$  in  $X = \{x_1, x_2, \dots, x_n\}$  as follows:

Let  $\varphi_{\mu_{AB}}(i) = |\mu_A(x_i) - \mu_B(x_i)|/2$ ,  $\varphi_{\nu_{AB}}(i) = |\nu_A(x_i) - \nu_B(x_i)|/2$ ,  $\varphi_{\phi_{AB}}(i) = |\phi_A(x_i) - \phi_B(x_i)|/2$  and  $x_i \in X$ . Then

$$S_d^p(A, B) = 1 - \frac{1}{\sqrt[p]{n}} \sqrt[p]{\sum_{i=1}^n (\varphi_{\mu_{AB}}(i) + \varphi_{\nu_{AB}}(i) + \varphi_{\phi_{AB}}(i))^p}, \quad (9)$$

where  $1 \leq p < \infty$ .

**Theorem 3.4**  $S_d^p(A, B)$  is the similarity measure between two GIFSs  $A$  and  $B$ .

**Proof** Obviously,  $S_d^p(A, B)$  satisfies (SP1) and (SP3). As to (SP2) and (SP4), we give the following proof.

(SP2): From (8), we have

$$\begin{aligned} S_d^p(A, B) &= 1 \\ &\Leftrightarrow \varphi_{\mu_{AB}}(i) = 0, \varphi_{\nu_{AB}}(i) = 0, \varphi_{\phi_{AB}}(i) = 0, \forall i = 1, \dots, n \\ &\Leftrightarrow \mu_A(x_i) = \mu_B(x_i), \nu_A(x_i) = \nu_B(x_i), \phi_A(x_i) = \phi_B(x_i), \forall x_i \in X \\ &\Leftrightarrow A = B. \end{aligned}$$

(SP4): Since  $A \subseteq B \subseteq C$ , we have  $\mu_A(x_i) \leq \mu_B(x_i) \leq \mu_C(x_i)$ ,  $\nu_A(x_i) \geq \nu_B(x_i) \geq \nu_C(x_i)$  and  $\phi_A(x_i) \geq \phi_B(x_i) \geq \phi_C(x_i)$  for any  $x_i \in X$ . Then we have

$$\begin{aligned} & \varphi_{\mu_{AB}}(i) + \varphi_{\nu_{AB}}(i) + \varphi_{\phi_{AB}}(i) \\ &= |\mu_A(x_i) - \mu_B(x_i)|/2 + |\nu_A(x_i) - \nu_B(x_i)|/2 + |\phi_A(x_i) - \phi_B(x_i)|/2 \\ &\leq |\mu_A(x_i) - \mu_C(x_i)|/2 + |\nu_A(x_i) - \nu_C(x_i)|/2 + |\phi_A(x_i) - \phi_C(x_i)|/2 \\ &= \varphi_{\mu_{AC}}(i) + \varphi_{\nu_{AC}}(i) + \varphi_{\phi_{AC}}(i). \end{aligned}$$

So we have

$$\sqrt[p]{\sum_{i=1}^n (\varphi_{\mu_{AC}}(i) + \varphi_{\nu_{AC}}(i) + \varphi_{\phi_{AC}}(i))^p} \geq \sqrt[p]{\sum_{i=1}^n (\varphi_{\mu_{AB}}(i) + \varphi_{\nu_{AB}}(i) + \varphi_{\phi_{AB}}(i))^p}.$$

Therefore,  $S_d^p(A, C) \leq S_d^p(A, B)$ . In the similar way, it is easy to prove  $S_d^p(A, C) \leq S_d^p(B, C)$ .

Similar to (8), considering the weight  $w_i$  of  $x_i \in X$ , the similarity measure of GIFSs  $A$  and  $B$  can be obtained as following form.

$$S_{dw}^p(A, B) = 1 - \sqrt[p]{\sum_{i=1}^n w_i (\varphi_{\mu_{AB}}(i) + \varphi_{\nu_{AB}}(i) + \varphi_{\phi_{AB}}(i))^p}, \quad (10)$$

where  $1 \leq p < \infty$ .

Likewise, for  $S_{dw}^p(A, B)$ , the following theorem holds.

**Theorem 3.5**  $S_{dw}^p(A, B)$  is the similarity measure between two GIFSs  $A$  and  $B$ .

**Proof** The proof is similar to that of Theorem 3.4.

## 4 An application to pattern recognition problem

Assume that a question related to pattern recognition is given using GIFSs.

Assume that there exist  $m$  patterns which are represented by GIFSs  $A_i = \{(x_i, \mu_A(x_i), \nu_A(x_i)) \mid x_i \in X\}$  ( $i = 1, 2, \dots, m$ ), where  $X = \{x_1, x_2, \dots, x_n\}$ . Suppose that there be a sample to be recognized which is represented by GIFS  $B = \{(x_i, \mu_B(x_i), \nu_B(x_i)) \mid x_i \in X\}$ . Set

$$S(A_{i_0}, B) = \max_{1 \leq i \leq n} \{S(A_i, B)\}, \quad (11)$$

where  $S(A_i, B)$  is the similarity measure between  $A_i$  and  $B$  ( $i = 1, 2, \dots, n$ ) given by (8) or (10).

According to the principle of the maximum degree of similarity between GIFSs, it can be decided that the sample  $B$  belongs to some pattern  $A_{i_0}$ .

**Example 4.1** Assume that there are three patterns denoted with GIFSs in  $X = \{x_1, x_2, x_3\}$ . Three patterns  $A_1, A_2$  and  $A_3$  are denoted as follows:

$$\begin{aligned} A_1 &= \{(x_1, 0.6, 0.3), (x_2, 0.8, 0.3), (x_3, 0.7, 0.4)\}; \\ A_2 &= \{(x_1, 0.5, 0.6), (x_2, 0.5, 0.4), (x_3, 0.7, 0.5)\}; \\ A_3 &= \{(x_1, 0.6, 0.5), (x_2, 0.7, 0.4), (x_3, 0.8, 0.4)\}. \end{aligned}$$

Assume that a sample  $B = \{(x_1, 0.6, 0.3), (x_2, 0.7, 0.5), (x_3, 0.7, 0.5)\}$  is given. Given three kinds of mineral fields, each is featured by the content of three minerals and contains one kind of typical hybrid minerals. The three kinds of typical hybrid minerals are represented by GIFSs  $A_1, A_2$  and  $A_3$  in  $X$ , respectively. Given another kind of hybrid mineral  $B$ , to which field does this kind of mineral  $B$  most probably belong to ?

For convenience, assume that the weight  $w_i$  of  $x_i$  in  $X$  are equal and  $p = 2$ . By (8) and (10), we have

$$\begin{aligned} S_g(A_1, B) &= 0.900, S_g(A_2, B) = 0.800, S_g(A_3, B) = 0.866; \\ S_d^2(A_1, B) &= 0.971, S_d^2(A_2, B) = 0.755, S_d^2(A_3, B) = 0.859. \end{aligned}$$

From this data, the proposed similarity measures  $S_g$  and  $S_d^p$  show the same classification according to the principle of the maximum degree of similarity between GIFSs. That is, the sample  $B$  belongs to the pattern  $A_1$ .

The results of above example indicates the proposed similarity measure to be good in pattern recognition problems. In the following example, the proposed similarity measure is used to characterize the similarity between linguistic variables.

**Example 4.2** Let  $F = \{(x, \mu_F(x), \nu_F(x)) : x \in X\}$  be a GIFS in  $X$ . For any positive real number  $n$ , We define the GIFS  $F^n$  as follows:

$$F^n = \{(x, (\mu_F(x))^n, 1 - (1 - \nu_F(x))^n) \mid x \in X\}.$$

Using the above operation, we also define the concentration and dilation of  $F$  as follows:

- concentration:  $\text{CON}(F) = F^2$ ;
- dilation:  $\text{DIL}(F) = F^{1/2}$ .

Like the fuzzy sets,  $\text{CON}(F)$  and  $\text{DIL}(F)$  may be treated as “very ( $F$ )” and “more or less ( $F$ )”, respectively.

In the next, we consider a GIFS  $F$  in  $X = \{6, 7, 8, 9, 10\}$  defined by

$$F = \{(6, 0.2, 0.9), (7, 0.4, 0.7), (8, 0.7, 0.4), (9, 0.9, 0.1), (10, 1, 0)\}.$$

With taking into account the characterization of linguistic variables, we regard  $F$  as “LARGE” in  $X$ . Using the operations of concentration and dilation



- $F^{1/2}$  may be treated as “More or less LARGE”,
- $F^2$  may be treated as “Very LARGE”,
- $F^4$  may be treated as “Very very LARGE”.

The proposed similarity measure is utilized to calculate the degree of similarity between these GIFSs. The results are summarized in Table 1. In Table 1, L., V.L., V.V.L. and M.L.L. denote LARGE, Very LARGE, Very very LARGE and More or less LARGE, respectively.

Table 1: The values calculated by the proposed similarity measure  $S_d^1$

$S_d^1$	M.L.L.	L.	V.L.	V.V.L.
M.L.L.	1	0.8562	0.7134	0.6022
L.	0.8562	1	0.8540	0.7428
V.L.	0.7134	0.8540	1	0.8848
V.V.L.	0.6022	0.7428	0.8848	1

From the viewpoint of mathematical operations, the similarities between the above GIFSs require the following conditions:

$$S(\text{M.L.L.}, \text{L.}) > S(\text{M.L.L.}, \text{V.L.}) > S(\text{M.L.L.}, \text{V.V.L.}), \quad (12)$$

$$S(\text{L.}, \text{M.L.L.}) > S(\text{L.}, \text{V.L.}) > S(\text{L.}, \text{V.V.L.}), \quad (13)$$

$$S(\text{V.L.}, \text{V.V.L.}) > S(\text{V.L.}, \text{L.}) > S(\text{V.L.}, \text{M.L.L.}), \quad (14)$$

$$S(\text{V.V.L.}, \text{V.L.}) > S(\text{V.V.L.}, \text{L.}) > S(\text{V.V.L.}, \text{M.L.L.}). \quad (15)$$

From Table 1, it can be seen that the proposed similarity measure  $S_d^1$  satisfies the requirements (12)-(15). Therefore, the proposed similarity measure  $S_d^1$  is reliable in applications with compound linguistic variables.

## 5 Conclusions

We apply the principle of maximum degree of similarity measures between GIFSs to solve the pattern recognition problem. Based on the geometrical interpretation of GIFSs, we take into account three parameters (membership, non-membership, hesitation margin) to propose a similarity measure for calculating the degree of similarity between GIFSs. Numerical example is given to illustrate the application of the developed similarity measures. Furthermore, we use the proposed similarity measures to characterize the similarity between linguistic variables.

## References

- [1] K. Atanassov, Intuitionistic fuzzy sets, *Fuzzy Sets and Systems*, 20, 87-96 (1986).
- [2] K. Atanassov, More on intuitionistic fuzzy sets, *Fuzzy Sets and Systems*, 33, 37-46 (1989).
- [3] K. Atanassov, New operations defined over the intuitionistic fuzzy sets, *Fuzzy Sets and Systems*, 61, 1377-142 (1994).
- [4] K. Atanassov and G. Gargov, Intuitionistic fuzzy logic, *CR Acad. Bulg. Soc.*, 43, 9-12 (1990).
- [5] H. Bustince and P. Burillo, Vague sets are intuitionistic fuzzy sets, *Fuzzy Sets and Systems*, 79, 403-405 (1996).
- [6] S.M. Chen, Similarity measure between vague sets and elements, *IEEE Trans. Systems Man Cybernt.*, 27, 153-158 (1997).
- [7] S.M. Chen and J.M. Tan, Handling multi-criteria fuzzy decision-making problems based on vague sets, *Fuzzy Sets and Systems*, 67, 163-172 (1994).
- [8] S.K. De, R. Biswas and A.R. Roy, Some operations on intuitionistic fuzzy sets, *Fuzzy Sets and Systems*, 114, 477-484 (2000).
- [9] S.K. De, R. Biswas and A.R. Roy, An application of intuitionistic fuzzy sets in medical diagnosis, *Fuzzy Sets and Systems*, 117, 209-213 (2001).
- [10] W.L. Gau and D.J. Buehere, Vague sets, *IEEE Trans. Systems Man Cybernt.*, 23, 610-614 (1994).
- [11] D.H. Hong and C. Kim, A note on similarity measures between vague sets and between elements, *Inform. Science*, 115, 83-96 (1999).
- [12] W.L. Hung and M.S. Yang, MS (2004) Similarity measures of intuitionistic fuzzy sets based on Hausdorff distance, *Pattern Recognition Lett.*, 25, 1603-1611 (2004).
- [13] A. Kaufman, *Introduction a la theorie des sous-ensembles flous*, Masson et Cie, Editeurs, 1973.
- [14] D. Li and C. Cheng, New similarity measures of intuitionistic fuzzy fuzzy sets and applications to pattern recognitions, *Pattern Recognition Lett.*, 23, 221-225 (2002).
- [15] Y. Li, D.L. Olson and Z. Qin, Similarity measures between intuitinistic fuzzy (vague) sets: A comparative analysis, *Pattern Recognition Lett.*, 28, 278-285 (2007).
- [16] Z. Liang and P. Shi, Similarity measures on intuitionistic fuzzy fuzzy sets, *Pattern Recognition Lett.*, 24, 2687-2693 (2003).

- [17] B. Loewer and R. Laddaga, Destroying the consensus, In B. Loewer (Guest Ed.): Special Issue on Consensus, *Synthese*, 62 (1985), 79-96.
- [18] T.K. Mondal and S.K. Samanta, Generalized intuitionistic fuzzy sets, *J. Fuzzy Math.*, 10, 839-861 (2002).
- [19] H.B. Mitchell, On the Dengfeng-Chuitian similarity measure and its application to pattern recognition, *Pattern Recognition Lett.*, 24, 3101-3104 (2003).
- [20] J.H. Park, Y.B. Park and K.M. Lim, Correlation coefficient of generalized intuitionistic fuzzy sets by statistical method, *Honam Math. J.*, 28, 317-326 (2006).
- [21] J.H. Park, J.H. Kang, Y.T. Jang, Y.C. Kwun and J.H. Koo, Similarity measure for generalized intuitionistic fuzzy sets and its application to group decision making. in: Proceeding of the International Conference on e-Commerce, e-Administration, e-Society, e-Education, and e-Technology, Macao, China (2010), 2246-2256.
- [22] E. Szmidt, *Applications of intuitionistic fuzzy sets in decision making*, (D. Sc. dissertation) Tech. Univ, Sofia 2000.
- [23] E. Szmidt and J. Baldwin, New similarity measure for intuitionistic fuzzy set theory and mass assignment theory, *Notes on IFSs*, 9, 60-76 (2003).
- [24] E. Szmit and J. Kacprzyk, Distances between intuitionistic fuzzy sets, *Fuzzy Sets and Systems*, 114, 505-518 (2000).
- [25] E. Szmit and J. Kacprzyk, A new concept of a similarity measure for intuitionistic fuzzy sets and its use in group decision making, *LNAI*, 3558, 272-282 (2005).
- [26] I.K. Vlachos and G.D. Sergiadis, Intuitionistic fuzzy information - Applications to pattern recognition, *Pattern Recognition Lett.*, 28, 197-206 (2007).
- [27] L.A. Zadeh, Fuzzy sets, *Inform. and Control*, 8, 338-353 (1965).

# TABLE OF CONTENTS, JOURNAL OF COMPUTATIONAL ANALYSIS AND APPLICATIONS, VOL. 20, NO. 5, 2016

Barnes-Type Peters of the First Kind and Poly-Cauchy of the First Kind Mixed-Type Polynomials, Dae San Kim, Taekyun Kim, Takao Komatsu, and Dmitry V. Dolgy, .....	803
On the Hyperstability of a Functional Equation in Commutative Groups, Muaadh Almahalebi, and Choonkil Park,.....	826
A Fractional Finite Difference Inclusion, Dumitru Baleanu, Shahram Rezapour, and Saeid Salehi,.....	834
Some Implicit Properties of the Second Kind Bernoulli Polynomials of Order $\alpha$ , C. S. Ryoo, and J. Y. Kang,.....	843
An Oscillation of the Solution For a Nonlinear Second-Order Stochastic Differential Equation, Iryna Komashynska, Mohammed AL-Smadi, Ali Atewi, and Ayed Al e'damat,.....	860
Fixed Point Theorems and T-stability of Picard Iteration For Generalized Lipschitz Mappings in Cone Metric Spaces Over Banach Algebras, Huaping Huang, Shaoyuan Xu, Hao Liu, and Stojan Radenovic,.....	869
Int-soft Filters of MTL-Algebras, Young Bae Jun, Seok Zun Song, Eun Hwan Roh, and Sun Shin Ahn,.....	889
Convergence Analysis of New Iterative Approximating Schemes with Errors for Total Asymptotically Nonexpansive Mappings in Hyperbolic Spaces, Ting-jian Xiong, and Heng-you Lan,.....	902
Some Ostrowski Type Integral Inequalities for Double Integral on Time Scales, Wajeeha Irshad, Muhammad Iqbal Bhatti, and Muhammad Muddassar,.....	914
The Henstock-Stieltjes Integral for Fuzzy-Number-Valued Functions on a Infinite Interval, Ke-feng Duan,.....	928
New Weighted $q$ -Chebyshev-Grüss Type Inequalities for Double Integrals, Zhen Liu, and Wengui Yang,.....	938
Quadratic $\rho$ -Functional Inequalities in Normed Spaces, Ikan Choi, Sunghoon Kim, George A. Anastassiou, and Choonkil Park,.....	949

**TABLE OF CONTENTS, JOURNAL OF COMPUTATIONAL  
ANALYSIS AND APPLICATIONS, VOL. 20, NO. 5, 2016**

**(continued)**

Solution of the Ulam Stability Problem for Quartic (a, b)-Functional Equations, Abdullah Alotaibi, John Michael Rassias, and S.A. Mohiuddine,.....957

Existence of Positive Solutions for Summation Boundary Value Problem for a Fourth-Order Difference Equations, Thanin Sitthiwiratham,.....969

Similarity Measure Between Generalized Intuitionistic Fuzzy Sets and Its Application to Pattern Recognition, Jin Han Park, Jongchul Hwang, Juhyung Kim, Byeongmuk Park, Juyoung Park, Jeongwoo Son, and Sihun Lee,.....984