

## Accepted Manuscript

Relaxation and diffusion models with non-singular kernels

HongGuang Sun, Xiaoxiao Hao, Yong Zhang, Dumitru Baleanu

PII: S0378-4371(16)30753-1

DOI: <http://dx.doi.org/10.1016/j.physa.2016.10.066>

Reference: PHYSA 17624

To appear in: *Physica A*

Received date: 28 June 2016

Revised date: 3 October 2016

Please cite this article as: H. Sun, X. Hao, Y. Zhang, D. Baleanu, Relaxation and diffusion models with non-singular kernels, *Physica A* (2016), <http://dx.doi.org/10.1016/j.physa.2016.10.066>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



# Relaxation and diffusion models with non-singular kernels

HongGuang Sun<sup>a</sup>, Xiaoxiao Hao<sup>a</sup>, Yong Zhang<sup>a,b,\*</sup>, Dumitru Baleanu<sup>c,d</sup>

*a. Institute of Soft Matter Mechanics, Department of Engineering Mechanics, Hohai University, 1 XiKang Road, Nanjing, Jiangsu 210098, China*

*b. Department of Geological Sciences, University of Alabama, Tuscaloosa, AL 35487, USA*

*c. Department of Mathematics, Cankaya University, Ankara, Turkey*

*d. Institute of Space Sciences, Magurele-Bucharest, Romania*

*Corresponding author: yzhang264@ua.edu (Yong Zhang)*

---

## Abstract

Anomalous relaxation and diffusion processes have been widely quantified by fractional derivative models, where the definition of the fractional-order derivative remains a historical debate due to its limitation in describing different kinds of non-exponential decays (e.g. stretched exponential decay). Meanwhile, many efforts by mathematicians and engineers have been made to overcome the singularity of power function kernel in its definition. This study first explores physical properties of relaxation and diffusion models where the temporal derivative was defined recently using an exponential kernel. Analytical analysis shows that the Caputo type derivative model with an exponential kernel cannot characterize non-exponential dynamics well-documented in anomalous relaxation and diffusion. A legitimate extension of the previous derivative is then proposed by replacing the exponential kernel with a stretched exponential kernel. Numerical tests show that the Caputo type derivative model with the stretched exponential kernel can describe a much wider range of anomalous diffusion than the exponential kernel, implying the potential applicability of the new derivative in quantifying real-world, anomalous relaxation and diffusion processes.

*Key words:* Anomalous relaxation and diffusion, Non-singular kernel, Stretched exponential function kernel, Memory characterization, Mean squared displacement

## 1. Introduction

In physics, processes involving the basic phenomena of relaxation and diffusion are of great relevance [1, 2]. Many non-equilibrium physical systems and processes can be characterized by relaxation and diffusion equations [3, 4]. However, anomalous relaxation and diffusion processes in various complex systems do not usually follow classical exponential decay or Gaussian statistics, but rather exhibit different relaxation or diffusion patterns. The traditional relaxation equation or Fick's second law therefore fails to describe the related behaviors [5, 6, 7], motivating the development of new mathematical or physical equation models [8, 9, 10]. Fractional derivatives, which can contain the historical memory and global correlation information of complex physical systems, have been considered a successful tool to characterize anomalous dynamics in various physics and engineering processes [11, 12, 13, 14]. Hereby, the fractional-derivative based relaxation equation and the corresponding diffusion equation models (which govern many physical phenomena such as heat, mass, or electron transfer; pollutants or liquid transport through porous media; amorphous semiconductors; colloid or proteins moving in biosystems or even in ecosystems) have been widely investigated in multiple disciplines including physics, mathematics, mechanics, and hydrology and control theory, among many others [7, 12, 15, 16].

From the viewpoint of mathematics, the reason that the fractional derivative model can characterize the system memory and global correlation well is due to its standard power-law memory kernel used to define the fractional derivative. However, the singularity of the power-law kernel is the main problem in numerical computation and applications of fractional-order partial differential equations (PDEs). Recently, a new definition of the fractional derivative with an exponential kernel (hereinafter named Caputo type definition) was recently proposed and further investigated from mathematical and physical aspects by Caputo and Fabrizio [17]. The main advantage of the new definition is that the singular power-law kernel is now replaced by a non-singular exponential kernel, which is easier to use in theoretical analysis, numerical calculations and real-world applications. The mathematical properties of the new definition such as the existence of solution [18] and Laplace transform [19, 21] have been investigated. Potential application of fractional derivative models with an exponential kernel was recently investigated for electromagnetism [22], diffusion [23], and heat transfer [24].

Physical aspects of Caputo type derivative models where the memory

kernel is an exponential function, however, have not been well documented. It motivated this study and further yielded some new thoughts on modeling anomalous relaxation and diffusion with stretched exponential decay. In the following we first analyze the analytical solution and the mean squared displacement for relaxation and diffusion governed by Caputo type derivative models with an exponential kernel. Results show that Caputo type derivative models with an exponential kernel cannot efficiently characterize the non-exponential nature of anomalous relaxation and diffusion. Hereby, a new notion of a Caputo type derivative where the memory kernel takes the form of a stretched exponential function is proposed to overcome the limitations of the existing exponential kernel. For the sake of simplicity, but without the loss of generality, we consider one dimensional relaxation and diffusion equations with a temporal derivative.

## 2. Caputo type derivative with non-singular kernels

The standard Caputo type fractional derivative contains a singular power-law kernel [25, 26]:

$${}^C D_t^\alpha f(t) = \frac{1}{\Gamma(1-\alpha)} \int_0^t \frac{f'(\tau) d\tau}{(t-\tau)^\alpha}, \quad 0 < \alpha < 1, \quad (1)$$

in which  $\alpha$  is the order of the fractional derivative, and  $\Gamma(\cdot)$  denotes the Gamma function.

To overcome the complexity caused by using a singular memory kernel  $(t-\tau)^{-\alpha}$ , Caputo and Fabrizio [17] recently proposed a derivative using the non-singular exponential function:

$${}^{CF1} D^\alpha f(t) = \frac{M(\alpha)}{(1-\alpha)} \int_a^t f'(\tau) \exp\left[-\frac{\alpha(t-\tau)}{1-\alpha}\right] d\tau, \quad (2)$$

where the super-script “CF1” denotes the Caputo and Fabrizio type derivative,  $a \in (-\infty, t)$  is the initial time in the history, and  $M(\alpha)$  is a normalization function such that  $M(0) = M(1) = 1$ . For simplicity, we choose  $M(\alpha) = 1$  in the following sections.

Caputo and Fabrizio [22] also investigated the potential expression of new derivative with a Gaussian-function kernel for gradient and Laplacian operators:

$${}^{CF2} D^\alpha f(t) = \frac{1+\alpha^2}{\sqrt{\pi^\alpha(1-\alpha)}} \int_a^t f'(\tau) \exp\left[-\frac{\alpha(t-\tau)^2}{1-\alpha}\right] d\tau, \quad (3)$$

which will be briefly analyzed in the following.

In addition, another interesting definition with a Mittag-Leffler function kernel has been proposed and further discussed by Atangana and Baleanu [20, 27]. The expression is stated as

$${}^{ABC}D^\alpha f(t) = \frac{B(\alpha)}{(1-\alpha)} \int_a^t f'(\tau) E_\alpha[-\frac{\alpha(t-\tau)^\alpha}{1-\alpha}] d\tau, \quad (4)$$

in which  $B(\alpha)$  is a normalization function, and  $E_\alpha(\cdot)$  denotes the Mittag-Leffler function. In this paper, we focus our attention on the definition with an exponential function kernel.

### 3. Power-law and exponential function kernels in characterizing relaxation and diffusion processes

Here we explore the dynamics of relaxation and diffusion processes characterized by temporal derivative models with known memory kernels.

#### 3.1. Fractional relaxation equation model

The following fractional-order relaxation equation is a simple and typical model used to describe various physical processes:

$$\begin{cases} \frac{d^\alpha u(t)}{dt^\alpha} = -\lambda u(t), 0 < \alpha \leq 1, \\ u(t=0) = U_0. \end{cases} \quad (5)$$

The variable  $u(t)$  can represent concentration in solute transport [10, 28, 29], the relaxation modulus in viscoelasticity [30], or input in control systems [31]. It is well known that this equation with an integer order ( $\alpha = 1$ ) can characterize the phenomena of exponential relaxation in physics.

#### (a) Fractional derivative relaxation equation with a power-law kernel

First we consider a fractional derivative with the traditional power-law function kernel. The Laplace transform of Eq. (5) can be stated as:

$$s\bar{u}(s) - U_0 s^{\alpha-1} = -\lambda \bar{u}(s), 0 < \alpha \leq 1, \quad (6)$$

where  $\bar{u}(s)$  represents the Laplace transform of  $u(t)$ . Inverse Laplace transform of Eq. (6) provides the analytical solution of Eq. (5) [30]:

$$u(t) = U_0 E_\alpha(-\lambda t^\alpha), \quad (7)$$

where  $E_\alpha(\cdot)$  denotes the Mittag-Leffler function [10, 34].

**(b) Relaxation equation model with an exponential kernel**

To obtain the analytical solution of Eq.(5) with an exponential kernel, we first use the Laplace transform of the fractional derivative with an exponential kernel Eq.(2) [17, 19]

$$\begin{aligned} L[{}^{CF1}D^\alpha u(t)](s) &= \frac{1}{1-\alpha} L(u'(t)) L[\exp(\frac{-\alpha t}{1-\alpha})] \\ &= \frac{su(s) - U_0}{s(1-\alpha) + \alpha}. \end{aligned} \quad (8)$$

Then one can get the expression of Eq. (5) after Laplace transform

$$\frac{su(s) - U_0}{s(1-\alpha) + \alpha} = -\lambda u(s). \quad (9)$$

Finally, the analytical solution of Eq. (5) can be obtained via an inverse Laplace transform with the initial condition defined in (5):

$$u(t) = U_0 e^{-\frac{\lambda \alpha t}{1+\lambda-\alpha\lambda}}. \quad (10)$$

A well-known and interesting feature of anomalous relaxation in complex systems is their non-exponential nature [32]. The solution of fractional derivative models with a power-law function kernel exhibits power-law decay when  $t \rightarrow \infty$ , while the solution of that with an exponential function kernel shows an exponential decay similar to the classical integer-order model. In fact, the integer-order relaxation equation with a relaxation coefficient  $\lambda\alpha/(1+\lambda-\alpha\lambda)$  gives the same analytical solution (8), from mathematical viewpoint.

*3.2. Mean squared displacement of the temporal derivative diffusion model*

Mean squared displacement (MSD) offers an important information of diffusive motion in physical and biophysical systems, and serves as an important criterion in anomalous diffusion [10, 33, 34]. To investigate the physical behavior of fractional derivative model with an exponential function kernel, we consider the following time fractional diffusion equation:

$$\begin{cases} \frac{\partial^\alpha c(x, t)}{\partial t^\alpha} = K \frac{\partial^2 c(x, t)}{\partial x^2}, & -\infty < x < +\infty, t > 0, \\ c(x, 0) = \delta(x), c(\pm\infty, t) = 0, \frac{\partial c(\pm\infty, t)}{\partial x} = 0, \end{cases} \quad (11)$$

where  $\delta(x)$  is the Dirac delta function,  $\alpha \in (0, 1]$  denotes the order of the time fractional derivative,  $K$  is a generalized diffusion coefficient,  $c(x, t)$  represents the target concentration.

Previous investigations have shown that the MSD of model (11) with the traditional Caputo fractional derivative expressed by Eq. (1) can be stated as [10]

$$\langle x^2(t) \rangle = \frac{2Kt^\alpha}{\Gamma(\alpha + 1)}. \quad (12)$$

We then derive the MSD for the Caputo type derivative model Eq.(11) with an exponential kernel. At the first step, we take integration of both sides of Eq. (11) with  $\int_{-\infty}^{\infty} x^2 dx$ , which leads to

$$\int_{-\infty}^{\infty} x^2 {}^{CF1}D^\alpha c(x, t) dx = \int_{-\infty}^{\infty} x^2 K \frac{\partial^2 c(x, t)}{\partial x^2} dx. \quad (13)$$

It can be written as the following form by changing the order of integral and differentiation:

$${}^{CF1}D_{0+}^\alpha \langle x^2(t) \rangle = \int_{-\infty}^{\infty} x^2 K \frac{\partial^2 c(x, t)}{\partial x^2} dx. \quad (14)$$

Clearly, Eq. (14) can be simplified into the following form based on the boundary conditions of model Eq. (11)

$${}^{CF1}D_{0+}^\alpha \langle x^2(t) \rangle = 2K. \quad (15)$$

The final expression of the MSD can be obtained by applying the Laplace transform of Eq. (15) and then taking inverse Laplace transform:

$$\langle x^2(t) \rangle = 2K\alpha t. \quad (16)$$

It is obvious that the fractional derivative diffusion equation model with an exponential kernel reduces to normal diffusion based on the criterion of MSD.

#### 4. Fractional derivative model with the memory kernel as a stretched exponential function

It is clear that the Caputo-Fabrizio derivative operator with an exponential kernel has obvious limitations in characterizing anomalous relaxation

and diffusion with non-exponential nature. Meanwhile, previous literature also showed that the memory kernel appearing in the generalized diffusion equation has various potential forms which can describe a wide range of experimental phenomena [35, 36]. Nevertheless, the relaxation and diffusion phenomena with the stretched exponential decay, which cannot be well characterized using the fractional derivative with a power function kernel, have been observed in various heterogeneous media or fully developed turbulence [2, 37]. Hereby, we consider a legitimate extension of the fractional derivative by replacing the exponential function kernel by a stretched exponential function kernel:

$${}^{SE}D^\alpha f(t) = \frac{M(\alpha)}{(1-\alpha)^{1/\alpha}} \int_a^t f'(\tau) \exp\left[-\frac{\alpha(t-\tau)^\alpha}{1-\alpha}\right] d\tau, \quad 0 < \alpha < 1, \quad (17)$$

in which the super-script ‘‘SE’’ represents the stretched exponential type Caputo definition, and  $M(\alpha)$  is a normalization function. For simplicity, here we take  $M(\alpha) = 1/\Gamma(1+\alpha)$ .

Since the above temporal derivatives differ only in the functional form of the memory kernel, we first compare the memory behavior of different kernels. Figure 1 shows that the power-law kernel represents strong memory in long time, compared to the exponential, stretched exponential, and Gaussian function kernels. Although the Caputo type derivative defined by the exponential kernel represents a process with memory, the long-time memory is very weak, which may explain why the solutions of the corresponding relaxation and diffusion equations are similar to classical integer-order derivative models. While the stretched exponential function kernel offers a relatively strong memory in long time, the memory property in small and long-time histories can be balanced in a wide range by changing the derivative order  $\alpha$ . Therefore, from this viewpoint, the stretched exponential function kernel is a better choice to characterize the memory (time) or nonlocal (space) process when using the Caputo type derivative with non-singular kernels.

Next, we investigate the physical behavior of Caputo type derivative relaxation and diffusion equations with a stretched exponential kernel, where the analytical solutions are not available. We use a finite difference numerical scheme to obtain accurate solutions of considered physical quantities [38, 39, 40].

Numerical solutions of the fractional derivative relaxation equation with a power-law or stretched exponential kernel are depicted in Figure 2. On the one hand, the Caputo type derivative relaxation model with a stretched



exponential kernel exhibits much faster decay at small times and much slower decay at large times for a small order of fractional derivative (such as  $\alpha = 0.2$ ). On the other hand, the Caputo type derivative relaxation model with a stretched exponential kernel produces a faster decay than that with a power-law kernel for a large  $\alpha$  (such as 0.8). This implies that the Caputo type derivative model with a stretched exponential kernel can describe a wider range of relaxation phenomena, compared to that with a power-law kernel.

We further investigate the Caputo type derivative diffusion equation with a stretched exponential kernel. Figure 3 shows truncated diffusive behavior of the Caputo type derivative diffusion equation with a stretched exponential kernel. This truncated diffusion behavior is analogous to the result of distributed-order or tempered fractional derivative models which may describe variations of MSD across scales [41, 42]. A nonlinear increasing of the MSD which represents anomalous diffusion is observed at the initial time period, while an approximately linear increasing of the MSD with time is observed at the long time range. It should be noted that a Caputo type derivative model with a stretched exponential or power-law kernel produces slower diffusive motion than a model with an exponential kernel.

Figure 1 shows that the power-law kernel represents a strong memory in long-time history, compared to non-singular kernels. To characterize the strong memory in complex systems using the stretched exponential kernel, we propose a modified definition by using a stretched exponential function kernel as follows:

$${}^{SE2}D^\alpha f(t) = \frac{1}{(1-\alpha)^{1/\alpha}\Gamma(1+\alpha)} \int_a^t f'(\tau) \exp\left[-\frac{\alpha}{1-\alpha} \left(\frac{t-\tau}{t}\right)^\alpha\right] d\tau. \quad (18)$$

Another advantage of definition (18) is that the fractional dimension problem in the application of fractional derivative operators can be solved by introducing the expression of  $(t-\tau)/t$ . Figure 4 shows that the stretched exponential kernel with  $(t-\tau)/t$  exhibits super-slow decay of the system's memory, which may be a better tool to describe relaxation or diffusion processes with extremely strong memory.

## 5. Conclusions

This study found that the Caputo type derivative with an exponential function kernel has limitations in characterizing the complex anomalous relaxation and diffusion processes. Here introduces an alternative notion of the

Caputo type derivative with a stretched exponential function kernel to describe real-world complex relaxation and diffusion processes. In the limiting case we recover the Caputo type derivative with exponential kernel.

There are still many issues which are worth further investigation. For examples, the lower limit of the integral in the Caputo type derivative definition should be  $-\infty$  from the mathematical viewpoint, otherwise the above analysis results are not strict. The new definition therefore should be improved, and the related mathematical analysis such as existence and uniqueness of solutions of obtained fractional ordinary equations should be investigated in a future study. Moreover, the applicability of Caputo type definition with stretched exponential function kernel, should be further verified by using a number of experimental data.

### Acknowledgments

This work was supported by the National Natural Science Foundation of China under grants 11572112, 11528205, and 41628202. Y. Zhang was also partially supported by the National Science Foundation grant DMS-1460319 and the University of Alabama. This paper does not necessarily reflect the views of the funding agencies. The authors thank the reviewers for their constructive suggestions.

### References

- [1] F. Mainardi, Fractional Relaxation-Oscillation and fractional diffusion-wave phenomena, *Chaos Soliton Fract.* 7(9)(1996) 146-1477.
- [2] J. Kakalios, R.A. Street, W.B. Jackson, Stretched-exponential relaxation arising from dispersive diffusion of hydrogen in amorphous silicon, *Phys. Rev. Lett.* 59(9)(1987) 1037.
- [3] V.D. Borman, V.N. Tronin, V.A. Byrkin, Anomalously slow relaxation of the system of liquid clusters in a disordered nanoporous medium according to the self-organized criticality scenario, *Phys. Lett. A* 380(18)(2016) 1615-1620.
- [4] J.H. Jeon, N. Leijnse, L.B. Oddershede, R. Metzler, Anomalous diffusion and power-law relaxation of the time averaged mean squared displacement in worm-like micellar solutions, *New J. Phys.* 15(4)(2013) 045011.

- [5] M.M. Meerschaert, B. Toaldo, Relaxation patterns and semi-Markov dynamics, (2015) arXiv preprint arXiv:1506.02951.
- [6] J.P. Bouchaud, A. Georges, Anomalous diffusion in disordered media: statistical mechanisms, models and physical applications, *Phys. Rep.* 195(4-5)(1990) 127-293.
- [7] R. Metzler, J.H. Jeon, A.G. Cherstvy, E. Barkai, Anomalous diffusion models and their properties: non-stationarity, non-ergodicity, and ageing at the centenary of single particle tracking, *Phys. Chem. Chem. Phys.* 16(44)(2014) 24128-24164.
- [8] C. Li, Y.Q. Chen, J. Kurths, Fractional calculus and its applications, *Phil. trans. R. Soc. A* 371(1990)(2013) 20130037.
- [9] K.L. Ngai, Relaxation and diffusion in complex systems, Springer Science & Business Media, 2011.
- [10] R. Metzler, J. Klafter, The random walk's guide to anomalous diffusion: a fractional dynamics approach, *Phys. Rep.* 339(1)(2000) 1-77.
- [11] R.L. Magin, C. Ingo, L. Colon-Perez, W. Triplett, T.H. Mareci, Characterization of anomalous diffusion in porous biological tissues using fractional order derivatives and entropy, *Micropor. Mesopor. Mat.* 178(2013) 39-43.
- [12] B. Yu, X. Jiang, H. Qi, An inverse problem to estimate an unknown order of a Riemann-Liouville fractional derivative for a fractional Stokes first problem for a heated generalized second grade fluid, *Acta Mech. Sin.* 31(2)(2015) 153-161.
- [13] C. Zeng, Y.Q. Chen, Q. Yang, Almost sure and moment stability properties of fractional order Black-Scholes model, *Fract. Calc. Appl. Anal.* 16(2)(2013) 317-331.
- [14] Z. Tomovski, T. Sandev, R. Metzler, J. Dubbeldam, Generalized space-time fractional diffusion equation with composite fractional time derivative, *Phys. A* 391(8)(2012) 2527-2542.
- [15] G.C. Wu, D. Baleanu, S.D. Zeng, Z.G. Deng, Discrete fractional diffusion equation, *Nonlinear Dynam.* 80(1-2)(2015) 281-286.

- [16] V.V. Uchaikin, R.T. Sibatov, Fractional kinetics in solids: anomalous charge transport in semiconductors, Dielectrics and Nanosystems, World Science, 2013.
- [17] M. Caputo, M. Fabrizio, A new definition of fractional derivative without singular kernel, *Progr. Fract. Differ. Appl.* 1(2)(2015) 1-13.
- [18] A. Atangana, B.S.T. Alkahtani, Analysis of the Keller-Segel model with a fractional derivative without singular kernel, *Entropy* 17(6)(2015) 4439-4453.
- [19] J. Losada, J.J. Nieto, Properties of a new fractional derivative without singular kernel, *Progr. Fract. Differ. Appl.* 1(2)(2015) 87-92.
- [20] A. Atangana, D. Baleanu, New fractional derivatives with nonlocal and non-singular kernel: Theory and application to heat transfer model, *Thermal Sci.* (2016) 18-18.
- [21] A. Atangana, On the new fractional derivative and application to nonlinear Fishers reaction-diffusion equation, *Appl. Math. Comput.* 273(2016) 948-956.
- [22] M. Caputo, M. Fabrizio, Applications of new time and spatial fractional derivatives with exponential kernels, *Progr. Fract. Differ. Appl.* 2(2)(2016) 1-11.
- [23] J.F. Gómez-Aguilar, M.G. López-López, V.M. Alvarado-Martinez, J. Reyes-Reyes, M. Adam-Medina, Modeling diffusive transport with a fractional derivative without singular kernel, *Phys. A* 447(2016) 467-481.
- [24] A. Atangana, D. Baleanu, New fractional derivatives with nonlocal and non-singular kernel: Theory and application to heat transfer model, *Therm. Sci.* (2016) DOI: 10.2298/TSCI160111018A.
- [25] M. Caputo, *Elasticitae Dissipazione*, Zanichelli, Bologna, 1965.
- [26] D. Baleanu, O.P. Agrawal, Fractional Hamilton formalism within Caputo's derivative, *Czech. J. Phys.* 56(10-11)(2006) 1087-1092.

- [27] A. Atangana, I. Koca, Chaos in a simple nonlinear system with Atangana-Baleanu derivatives with fractional order, *Chaos Soliton. Fract.* (2016) <http://dx.doi.org/10.1016/j.chaos.2016.02.012>.
- [28] A.V. Chechkin, V.Y. Gonchar, R. Gorenflo, N. Korabel, I.M. Sokolov, Generalized fractional diffusion equations for accelerating subdiffusion and truncated Lévy flights, *Phys. Rev. E* 78(2)(2008) 021111.
- [29] Y. Zhang, D.A. Benson, D.M. Reeves, Time and space nonlocalities underlying fractional-derivative models: Distinction and literature review of field applications, *Adv. Water Resour.* 32(4)(2009) 561-581.
- [30] F. Mainardi, *Fractional Calculus and Waves in Linear Viscoelasticity*, Imperial College Press, London, 2010.
- [31] H. Sheng, Y. Li, Y.Q. Chen, Application of numerical inverse Laplace transform algorithms in fractional calculus, *J. Franklin Inst.* 348(2)(2011) 315-330.
- [32] A. Stanislavsky, K. Weron, A. Weron, Anomalous diffusion approach to non-exponential relaxation in complex physical systems, *Commun. Nonlinear Sci. Numer. Simulat.* 24(1)(2015) 117-126.
- [33] E. Kepten, A. Weron, G. Sikora, K. Burnecki, Y. Garini, Guidelines for the fitting of anomalous diffusion mean square displacement graphs from single particle tracking experiments, *PloS one* 10(2)(2015) e0117722.
- [34] H.G. Sun, W. Chen, H. Sheng, Y.Q. Chen, On mean square displacement behaviors of anomalous diffusions with variable and random orders, *Phys. Lett. A* 374(7)(2010) 906-910.
- [35] T. Sandev, A. Chechkin, H. Kantz, R. Metzler, Diffusion and Fokker-Planck-Smoluchowski Equations with Generalized Memory Kernel, *Fract. Calc. Appl. Anal.* 18(4)(2015) 1006-1038.
- [36] I.M. Sokolov, J. Klafter, From diffusion to anomalous diffusion: a century after Einstein's Brownian motion, *Chaos* 15(2)(2005) 026103.
- [37] A. La Porta, G.A. Voth, A.M. Crawford, et al. Fluid particle accelerations in fully developed turbulence, *Nature* 409(6823)(2001) 1017-1019.

- [38] K. Diethelm, N.J. Ford, Analysis of fractional differential equations, J. Math. Anal. Appl. 265(2)(2002) 229-248.
- [39] X. Zhao, Z.Z. Sun, Compact Crank-Nicolson schemes for a class of fractional Cattaneo equation in inhomogeneous medium, J. Sci. Comput. 62(3)(2015) 747-771.
- [40] H. Wang, K. Wang, T. Sircar, A direct  $O(N \log 2N)$  finite difference method for fractional diffusion equations, J. Comput. Phys. 229(21)(2010) 8095-8104.
- [41] A.V. Chechkin, R. Gorenflo, I.M. Sokolov, Retarding subdiffusion and accelerating superdiffusion governed by distributed-order fractional diffusion equations, Phys. Rev. E 66(4)(2002) 046129.
- [42] M.M. Meerschaert, Y. Zhang, B. Baeumer, Tempered anomalous diffusion in heterogeneous systems, Geophys. Res. Lett. 35(17)(2008) L17403.




Figure 1: Memory property represented by using different kernels with  $\alpha = 0.5$ : the power-law function kernel is  $(t - \tau)^{-\alpha}$ , the exponential function kernel  $\exp(-\frac{\alpha*(t-\tau)}{1-\alpha})$ , the Gaussian function kernel  $\exp(-\frac{\alpha*(t-\tau)^2}{1-\alpha})$ , and the stretched exponential function kernel  $\exp(-\frac{\alpha*(t-\tau)^\alpha}{1-\alpha})$ .




Figure 2: Comparison of Caputo type derivative relaxation equations with a power-law kernel and a stretched exponential kernel. Here the legend “SE” represents stretched exponential. The blue lines represent concentration relaxation curves drawn by using the stretched exponential function kernel with  $\alpha = 0.2, 0.5$  and  $0.8$  (from top to bottom). The black lines denote concentration relaxation curves drawn by using the power-law function kernel with  $\alpha = 0.2, 0.5$  and  $0.8$  (from top to bottom). In this simulation the relaxation coefficient  $\lambda = 1.0$ .






Figure 3: The mean squared displacement (MSD) of the Caputo type derivative diffusion model with a stretched exponential function, power-law function, and exponential function as the kernel with  $\alpha = 0.5$  and the diffusion coefficient  $K = 1.0$ .




Figure 4: Memory property represented by different kernels with  $\alpha = 0.5$ : the power-law function kernel  $(t - \tau)^{-\alpha}$ , the stretched exponential function kernel  $\exp\left[-\frac{\alpha*(t-\tau)^\alpha}{1-\alpha}\right]$ , and the stretched exponential function(V2) kernel  $\exp\left[-\frac{\alpha}{1-\alpha}\left(\frac{t-\tau}{t}\right)^\alpha\right]$ .

**Highlights**

1. Relaxation model with exponential kernel yields exponential decay.
2. Mean squared displacement of Caputo type derivative diffusion model is offered.
3. A legitimate extension is proposed to characterize non-exponential dynamics.
4. The memory behaviors of different kernels are presented.