

## ADOMIAN-PADÉ APPROXIMATE SOLUTIONS TO THE CONFORMABLE NON-LINEAR HEAT TRANSFER EQUATION

by

**Aliyu Isa ALIYU<sup>a,b</sup>, Mustafa INC<sup>a</sup>, Abdullahi YUSUF<sup>a,b</sup>,  
and Dumitru BALEANU<sup>c,d\*</sup>**

<sup>a</sup> Department of Mathematics Faculty Science, Science Faculty, Firat University, Elazig, Turkey

<sup>b</sup> Department of Mathematics, Faculty of Science, Federal University Dutse, Jigawa, Nigeria

<sup>c</sup> Department of Mathematics, Cankaya University, Ankara, Turkey

<sup>d</sup> Institute of Space Sciences, Magurele, Romania

Original scientific paper

<https://doi.org/10.2298/TSCI181029037A>

*This paper adopts the Adomian decomposition method and the Padé approximation technique to derive the approximate solutions of a conformable heat transfer equation by considering the new definition of the Adomian polynomials. The Padé approximate solutions are derived along with interesting figures showing the approximate solutions.*

Key words: Adomian decomposition method, conformable heat transfer equation, Padé approximation

### Introduction

It is well-known that the majority of the real-world physical phenomena are modeled by mathematical equations, especially PDE [1]. The investigations of the exact and numerical solutions of various PDE have become a very important practice by different scholars [2]. In recent time, fractional derivatives have been applied in various fields of physical sciences such as reaction diffusion, heat transform, control, and many more. Fractional derivatives have many kinds of definitions [3-5]. We consider the non-linear conformable heat transfer equation in the following form [6]:

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \frac{\partial^2 u}{\partial x^2} - 2u^3 \quad (1)$$

with the initial condition:

$$u(x, 0) = \frac{1 + 2x}{x^2 + x + 1} \quad (2)$$

Equation (1) has been solved using different techniques, including the fractional complex transform along with He's variational iteration method [6], homotopy perturbation method [7], sub-equation method [8]. This paper applies the Adomian decomposition method and the Padé approximation technique [9] to search for an approximate solutions of the model problem. Several research results have been report in the last few decades in this avenue [10-20].

\* Corresponding author, e-mail: [dumitru@cankaya.edu.tr](mailto:dumitru@cankaya.edu.tr)

### Conformable fractional derivative

In what follows, we give a brief description of the conformable derivative given by Khalil *et al.* [5]:

*Definition 1.* Let

$$f : [a, b] \times (0, \infty) \rightarrow \mathbb{R} \quad (3)$$

then the conformable fractional derivative of  $f$  is defined:

$${}_t \mathbb{T}_\alpha^a(f)(x, t) = \lim_{\varepsilon \rightarrow 0} \frac{f(x, t + \varepsilon(t - \alpha)^{1-\alpha}) - f(x, t)}{\varepsilon}, \quad \alpha \in (0, 1] \quad (4)$$

for all  $t > 0$ .

When  $a = 0$ , we have  ${}_t \mathbb{T}_\alpha^a(f)(x, t) = {}_t \mathbb{T}_\alpha(f)(x, t)$ . This definition of fractional derivative satisfies the following properties.

*Theorem 2.* Let  $\alpha \in (0, 1]$  and  $a, b \in \mathbb{R}$ , then:

- ${}_t \mathbb{T}_\alpha(au + bv) = a {}_t \mathbb{T}_\alpha(u) + b {}_t \mathbb{T}_\alpha(v)$
- ${}_t \mathbb{T}_\alpha(t^\lambda) = \lambda t^{\lambda-\alpha}, \lambda \in \mathbb{R}$
- ${}_t \mathbb{T}_\alpha(uv) = u {}_t \mathbb{T}_\alpha(v) + v {}_t \mathbb{T}_\alpha(u)$
- ${}_t \mathbb{T}_\alpha\left(\frac{u}{v}\right) = \frac{u {}_t \mathbb{T}_\alpha(v) - v {}_t \mathbb{T}_\alpha(u)}{v^2}$
- ${}_t \mathbb{T}_\alpha(u)(t) = t^{1-\alpha} u'(t), u \in C^1$
- ${}_t \mathbb{T}_\alpha(u)(t) = (t - \alpha)^{1-\alpha} u'(t), u \in C^1$

### Padé approximation

Consider a system of fractional differential equations [9]:

$$\begin{cases} L_1 u_1 + R_1(u_1, u_2, \dots, u_s) + N_1(u_1, u_2, \dots, u_s) = g_1 \\ L_2 u_1 + R_2(u_1, u_2, \dots, u_s) + N_2(u_1, u_2, \dots, u_s) = g_2 \\ \dots \\ L_s u_1 + R_s(u_1, u_2, \dots, u_s) + N_s(u_1, u_2, \dots, u_s) = g_s \end{cases} \quad (5)$$

Subject to the initial or boundary conditions:

$$\begin{aligned} B_1(u_1, \dots, u_s) &= \sum \sum \rho_{i,n}^1 \frac{\partial_{(n\alpha)} u_i(t, x, t, y, \dots)}{\partial t^{n\alpha}} \Big|_{t=b_{i,n}} = p_1(x, y, \dots), \\ B_2(u_1, \dots, u_s) &= \sum \sum \rho_{i,n}^2 \frac{\partial_{(n\alpha)} u_i(t, x, t, y, \dots)}{\partial t^{n\alpha}} \Big|_{t=b_{i,n}} = p_2(x, y, \dots) \end{aligned} \quad (6)$$

Applying the inverse  $L_i^{-1}$  on both sides of eq. (5) gives:

$$\begin{cases} u_1 = \phi_1 + L_1^{-1}g_1 - L_1^{-1}R_1(u_1, u_2, \dots, u_s) - L_1^{-1}N_1(u_1, u_2, \dots, u_s) \\ u_2 = \phi_2 + L_2^{-1}g_2 - L_2^{-1}R_2(u_1, u_2, \dots, u_s) - L_2^{-1}N_2(u_1, u_2, \dots, u_s) \\ \dots \\ u_s = \phi_s + L_s^{-1}g_s - L_s^{-1}R_s(u_1, u_2, \dots, u_s) - L_s^{-1}N_s(u_1, u_2, \dots, u_s) \end{cases} \quad (7)$$

in which the  $\phi_i$  engulf the constants of integration w.r.t the variable  $t$ . The solution  $u_i$  can be dissolve into a series as [9]:

$$u_i = \sum_{m=0}^{\infty} u_{i,m}, \quad i = 1, 2, \dots, s, \quad (8)$$

$$\left[ \sum_{n=0}^{\infty} A_n \right]_{n \geq m} = \left[ \left[ \sum_{n=0}^{\infty} \left[ \frac{1}{n!} \sum_{v=0}^{\infty} (u_v - u_0)^n \frac{\partial^n N(u_0)}{\partial u_0^n} \right]_{v \geq p_1(m)} \right]_{n \geq p_1(m)} \right]_{\sum \tilde{p}_j \geq p_3(m)} \quad (9)$$

and the non-linear terms  $N_i(u_1, u_2, \dots, u_s)$  are assumed to be decomposed into a multivariable Adomian polynomials (AP) given:

$$N_i(u_1, u_2, \dots, u_s) = \sum_{m=0}^{\infty} A_{i,m}(u_{1,0}, \dots, u_{2,0}, \dots, u_{2,m}, u_{s,0}, \dots, u_{s,m}) \lambda^m \quad (10)$$

which are defined by

$$A_{i,m} = \frac{1}{m!} \frac{d^m}{d\lambda^m} N_i^{(m)}[u_1(\lambda), \dots, u_s(\lambda)] \Big|_{\lambda=0}$$

and where  $A_{i,m}$  can be generated for the four classes of the AP as shown by the cases:

– *Class I.* The AP  $A_{i,m}^{(I)}$ , is represented:

$$\theta_{i,m}^{(I)} = \sum_{n=0}^{\infty} \frac{[\phi_{i,m}^{(I)} - u_{i,0}]}{n!} N_i^n(u_i, 0), \quad m \geq 1 \quad (11)$$

and

$$\phi_{i,m}^{(I)} = \sum_{v=0}^{m-1} u_{i,v}^{(I)}, \quad A_{i,0}^{(I)} \equiv \theta_{i,1}^{(I)} \quad (12)$$

for  $m \geq 1: A_{i,m}^{(I)} = \theta_{i,m+1}^{(I)} - \theta_{i,m}^{(I)}$ .

– *Class II.* The AP  $A_{i,m}^{(II)}$  is represented:

$$\theta_{i,m}^{(II)} = \sum_{n=0}^{\infty} \frac{[\phi_{i,m}^{(II)} - u_{i,0}]}{n!} N_i^n(u_i, 0), \quad m \geq 1 \quad (13)$$

and

$$\phi_{i,m}^{(II)} = \sum_{v=0}^{m-1} u_{i,v}^{(II)}, \quad A_{i,0}^{(II)} \equiv \theta_{i,1}^{(II)} \quad (14)$$

for  $m \geq 1: A_{i,m}^{(II)} = \theta_{i,m+1}^{(II)} - \theta_{i,m}^{(II)}$ .

– *Class III.* The AP  $A_{i,m}^{(III)}$  is represented:

$$\theta_{i,m}^{(III)} = \sum_{n=0}^{\infty} \frac{[\phi_{i,m-n+1}^{(III)} - u_{i,0}]}{n!} N_i^n(u_i, 0), \quad m \geq 1 \quad (15)$$

and

$$\phi_{i,m}^{(III)} = \sum_{v=0}^{m-1} u_{i,v}^{(III)}, \quad A_{i,0}^{(III)} \equiv \theta_{i,1}^{(III)} \quad (16)$$

for  $m \geq 1$ :  $A_{i,m}^{(III)} = \theta_{i,m+1}^{(III)} - \theta_{i,m}^{(III)}$ .

– *Class IV.* The AP  $A_{i,m}^{(IV)}$  is represented:

$$\theta_{i,m}^{(IV)} = \left\| \sum_{n=0}^{\infty} \frac{[\phi_{i,m-n+1}^{(IV)} - u_{i,0}]}{n!} N_i^n(u_i, 0) \right\|_{\sum j, j \geq m}, \quad m \geq 1 \quad (17)$$

and

$$\phi_{i,m-n+1}^{(IV)} = \sum_{v=0}^{m-1} u_{i,v}^{(IV)}, \quad A_{i,0}^{(IV)} \equiv \theta_{i,1}^{(IV)} \quad (18)$$

for  $m \geq 1$ :  $A_{i,m}^{(IV)} = \theta_{i,m+1}^{(IV)} - \theta_{i,m}^{(IV)}$ .

Substitution eqs. (8) and (10), and utilizing the Adomian-Rach double decomposition approach, one can get:

$$\begin{cases} \sum_{m=0}^{\infty} u_{1,m} = \sum_{m=0}^{\infty} (\phi_{1,m} + L_1^{-1} g_1 - L_1^{-1} R_1 \left( \sum_{m=0}^{\infty} u_{1,m}, \dots, \sum_{m=0}^{\infty} u_{s,m} \right)) - L_1^{-1} \sum_{m=0}^{\infty} A_{1,m} \\ \sum_{m=0}^{\infty} u_{2,m} = \sum_{m=0}^{\infty} (\phi_{2,m} + L_2^{-1} g_1 - L_2^{-1} R_2 \left( \sum_{m=0}^{\infty} u_{1,m}, \dots, \sum_{m=0}^{\infty} u_{s,m} \right)) - L_2^{-1} \sum_{m=0}^{\infty} A_{2,m} \\ \dots \\ \sum_{m=0}^{\infty} u_{s,m} = \sum_{m=0}^{\infty} (\phi_{s,m} + L_s^{-1} g_1 - L_s^{-1} R_s \left( \sum_{m=0}^{\infty} u_{1,m}, \dots, \sum_{m=0}^{\infty} u_{s,m} \right)) - L_s^{-1} \sum_{m=0}^{\infty} A_{s,m} \end{cases} \quad (19)$$

For the specified initial value problems [12], with  $L = d^\alpha / dt^\alpha$  being the operator,  $k_i - 1 < \alpha \leq k_i$ , we have:

$$\phi_{i,m} = C_{i,1}^m + C_{i,2}^m t + \dots, C_{i,k_i}^m t^{k_i} \quad (20)$$

with  $C_{i,j}^m$ ,  $j = 1, \dots, k_i$ . For fractional IVP we have:

$$\phi_{i,m} = \sum_{l=0}^{k_i-1} t^{l\alpha} C_{i,l}^m \quad (21)$$

in which  $C_{i,j}^m$ ,  $j = 1, \dots, k_i$  are computed using the conditions.

Thus,  $u_{i,m}$  ( $m \geq 0$ ) can be derived from:

$$\begin{cases} u_{i,0} = \phi_{i,0} + L_i^{-1} g_i \\ u_{i,m+1} = \phi_{i,m+1} - R_i(u_{1,m}, \dots, u_{s,m}) - L_i^{-1} A_{i,m} \\ m \geq 0 \end{cases} \quad (22)$$

The solutions of the approximants are:

$$\phi_{i,r} = \sum_{m=0}^{r-1} u_{i,m} \quad (23)$$

From eq. (23), using  $u_{i,0}$  all other solution terms  $u_{i,j}$ ,  $j = 1, 2, \dots$ , can be obtained. Thus, the approximation  $\phi_{i,r}$  for the solutions  $u_i$ ,  $i = 1, 2, \dots, s$  are derived.

*The Padé approximation*

To fasten the convergence of the series solutions, the Padé approximants method is also assimilated into the previous described algorithm given as: The main concept of Padé approximants is to replace a series function  $f(z) = \sum_{n=0}^{\infty} c_n z^n$  [11] by:

$$\frac{A^{[L/M]}(z)}{B^{[L/M]}(z)} = \frac{a_0 + a_1 z + \dots, a_L z^L}{1 + b_1 z + \dots, b_M z^M} \quad (24)$$

given by  $[L/M]$  called the Padé approximants.

For a function  $f(z) = c_0 + c_1 z + \dots, c_2 z^2 + \dots$ , the coefficients  $a_0, a_1, \dots, a_L, b_0, b_1, \dots, b_M$  can be calculated by matching the series coefficients. Therefore, one can acquire the following sequence:


$$\begin{aligned} - [1/1] &= \frac{a_0 + a_1 z}{1 + b_1 z}, \quad \lim_{z \rightarrow \infty} [1/1] = \frac{a_1}{b_1}, \\ - [2/2] &= \frac{a_0 + a_1 z + a_2 z^2}{1 + b_1 z + b_2 z^2}, \quad \lim_{z \rightarrow \infty} [2/2] = \frac{a_2}{b_2}, \\ - [3/3] &= \frac{a_0 + a_1 z + a_2 z^2 + a_3 z^3}{1 + b_1 z + b_2 z^2 + b_3 z^3}, \quad \lim_{z \rightarrow \infty} [3/3] = \frac{a_3}{b_3}, \\ - \dots \\ - \lim_{z \rightarrow \infty} [m/m] &= \frac{a_m}{b_m}. \end{aligned}$$

*Application to the conformable heat transfer equation*


We will consider the Class (IV) AP eq. (17) to compute the truncated series solutions and then compare the exact eq. (2) and approximate solutions. For each of the cases to be considered, we will use Padé approximant sizes of  $[10/10]$  and  $[20/20]$ .

- If we select  $\alpha = 0.5$ , using 7 terms of an Adomian power series, Padé approximant size of  $[10/10]$  and using the Class (IV) AP, we obtain the following expression of the Padé approximant for this case (figs. 1 and 2):

$$\begin{aligned} \text{Pade}_4[10/10] &= 1 - 2.25676\sqrt{t} + 44.34929t^2 - 22.98715t^{3/2} + 8t - \\ &\quad - 1472.2094t^3 + 42956.36098t^{7/2} - 428.22830t^{5/2} \end{aligned} \quad (25)$$



**Figure 1.** Comparison of the Adomian-Padé approximate solutions eq. (25) for the different steps and  $0 \leq t \leq 1$  and  $0 \leq u \leq 100$  (for color image see journal web site)



**Figure 2:** The decomposition solution for the 3<sup>rd</sup>-order series for  $0 \leq x \leq 1$ ,  $0 \leq t \leq 1$  and  $0 \leq u \leq 60$  with eq. (2) (for color image see journal web site)

- If we select  $\alpha = 1$ , using 10 terms of an Adomian power series, Padé approximant size of [10/10] and using the Class (IV) AP, we obtain the following expression of the Padé approximant for this case (figs. 3 and 4):


$$\text{Pade}_4[10/10] = 1 - 2t + 4t^2 - 8t^3 + 16t^4 - 32t^5 + 64t^6 - 128t^7 + 256t^8 - 512t^9 + 1024t^{10} \quad (26)$$

- If we select  $\alpha = 0.8$ , using 10 terms of an Adomian power series, Padé approximant size of [20/20] and using the Class (IV) AP, we obtain the following expression of the Padé approximant for this case (figs. 5 and 6):


$$\text{Pade}_4[20/20] = 1 - 2t + 4t^2 - 8t^3 + 16t^4 - 32t^5 + 64t^6 - 128t^7 + 256t^8 - 512t^9 + 1024t^{10} \quad (27)$$

### Concluding remarks


In this work, we have successfully used the Padé approximation technique to derive the approximate solutions of a conformable heat transfer equation. The result shows that the method is powerful and efficient for solving the non-linear fractional differential equations arising in engineering and science. It is observed that the numerical solutions are in closed agreement with the exact solution. Some interesting figures have been given the comparison approximate solutions for the different steps in figs. 1-6.




**Figure 3.** Comparison of the Adomian-Padé approximate solutions eq. (26) for the different steps  $0 \leq t \leq 1$  and  $0 \leq u \leq 100$  (for color image see journal web site)



**Figure 4.** The decomposition solution for the 10<sup>th</sup>- order series for  $0 \leq x \leq 1$ ,  $0 \leq t \leq 1$  and  $0 \leq u \leq 60$  with eq. (2) (for color image see journal web site)



**Figure 5.** Comparison of the Adomian-Padé approximate solutions eq. (27) for the different steps  $0 \leq t \leq 1$  and  $0 \leq u \leq 100$  (for color image see journal web site)



**Figure 6. The decomposition solution for the 10<sup>th</sup>- order series for  $0 \leq x \leq 1$ ,  $0 \leq t \leq 1$  and  $0 \leq u \leq 60$  with eq. (2) (for color image see journal web site)**

## References

- [1] Whitham, G. B., *Linear and Nonlinear Waves*, John Wiley, New York, USA, 1974
- [2] Ferziger, J. H., A Note on Numerical Accuracy, *Int. J. Num. M. in Fluids*, 8 (1988), Sept., pp. 995-996
- [3] Podlubny, I., *Fractional Differential Equations*, Academic Press, New York, USA, 1999
- [4] Yang, X. J., *Advanced Local Fractional Calculus and its Applications*, World Science Publisher, New York, USA, 2012
- [5] Khalil, R., et al., A New Definition of Fractional Derivative, *J. Comput. Appl. Mat.*, 264 (2014), July, pp. 65-70
- [6] Wang, K. L., Liu, Y., He's Fractional Derivative for Nonlinear Fractional Heat Transfer Equation, *Thermal Science*, 20 (2016), 3, pp. 793-796
- [7] Zhang, M. F., et al., Efficient Homotopy Perturbation Method for Fractional Nonlinear Equations Using Sumudu Transform, *Thermal Science*, 19 (2015), 4, pp. 1167-1171
- [8] Ma . H. C., et al., Exact Solutions of Nonlinear Fractional Partial Differential Equations by Fractional Sub-Equation Method, *Thermal Science*, 19 (2015), 4, pp.1239-1244
- [9] Lin, Y., et al., Symbolic Computation of Analytic Approximate Solutions for Nonlinear Fractional Differential Equations, *Computer Physics Communications*, 184 (2013), 1, pp.130-141
- [10] Rach, R., A New Definition of the Adomian Polynomials, *Kybernetes*, 37 (2008), 7, pp. 910-955
- [11] Hosseini, M., et al., On the Convergence of Adomian Decomposition Method, *Applied Mathematics and Computation*, 182 (2006), 1, pp. 536-543
- [12] Wu, G., Adomian Decomposition Method for Non-Smooth Initial Value Problems, *Mathematical and Computer Modelling*, 54 (2011), 9-10, pp. 2104-2108
- [13] Qin, Z., et al., Analytic Study of Solitons in Non-Kerr Nonlinear Negative Index Materials, *Nonlinear Dynamics*, 86 (2016), 1, pp. 623-638
- [14] Mirzazadeh, M, et al., Optical Solitons with Complex Ginzburg-Landau Equation, *Nonlinear Dynamics*, 85 (2016), 3, pp. 1979-2016
- [15] Bo, L., et al., Soliton Interactions for Optical Switching Systems with Symbolic Computations, *Optik*, 175 (2018), Dec., pp.177-180
- [16] Zubair, A., et al., Analytic Study on Optical Solitons in Parity-Time-Symmetric Mixed Linear and Nonlinear Modulation Lattices with Non-Kerr Nonlinearities, *Optik*, 173 (2018), Nov., pp. 249-262
- [17] Weitian, Y., et al., Periodic Oscillations of Dark Solitons in Nonlinear Optics, *Optik*, 165 (2018), July, pp. 341-344
- [18] Zhou, Q., et al., Optical Solitons in Media with Time-Modulated Nonlinearities and Spatiotemporal Dispersion, *Nonlinear Dynamics*, 80 (2015), 1-2, pp. 983-987
- [19] Zhou, Q., Optical Solitons for Biswas-Milovic Model with Kerr Law and Parabolic Law Nonlinearities, *Nonlinear Dynamics*, 84 (2016), 2, pp. 677-681
- [20] Triki, H., et al., Chirped Dark Solitons in Optical Metamaterials, *Optik*, 158 (2018), Apr., pp. 312-315