

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/339984149>

A note on $(p; q)$ -analogue type of Fubini numbers and polynomials

Article in *AIMS Mathematics* · March 2020

DOI: 10.3934/math.2020177

CITATIONS

9

READS

64

3 authors:



Dr. Waseem Ahmad Khan

Prince Mohammad Bin Fahd University, Al Khobar Saudi Arabia

175 PUBLICATIONS 889 CITATIONS

[SEE PROFILE](#)



Kottakkaran Sooppy Nisar

Prince Sattam bin Abdulaziz University

866 PUBLICATIONS 6,791 CITATIONS

[SEE PROFILE](#)



Dumitru Baleanu

Institute of Space Sciences

1,473 PUBLICATIONS 45,579 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



International Conference on Mathematical Modelling in Applied Sciences, ICMAS'17 [View project](#)



Nonlinear Dynamics and Complexity (Online conference) [View project](#)



Research article

A note on (p, q) -analogue type of Fubini numbers and polynomials

Waseem Ahmad Khan¹, Kottakkaran Sooppy Nisar^{2,*} and Dumitru Baleanu^{3,4,5}

¹ Department of Mathematics and Natural Sciences, Prince Mohammad Bin Fahd University, P.O. Box 1664, Al Khobar 31952, Kingdom of Saudi Arabia

² Department of Mathematics, College of Arts and Sciences, Wadi Aldawaser, 11991, Prince Sattam bin Abdulaziz University, Saudi Arabia

³ Department of Mathematics, Cankaya University, Turkey

⁴ Institute of Space Sciences, 077125 Magurele-Bucharest, Romania

⁵ Department of Medical Research, China Medical University Hospital, China Medical University, Taichung, 40447, Taiwan

* **Correspondence:** Email: n.sooppy@psau.edu.sa; Tel: +966563456976.

Abstract: In this paper, we introduce a new class of (p, q) -analogue type of Fubini numbers and polynomials and investigate some properties of these polynomials. We establish summation formulas of these polynomials by summation techniques series. Furthermore, we consider some relationships for (p, q) -Fubini polynomials associated with (p, q) -Bernoulli polynomials, (p, q) -Euler polynomials and (p, q) -Genocchi polynomials and (p, q) -Stirling numbers of the second kind.

Keywords: (p, q) -calculus; (p, q) -Bernoulli polynomials; (p, q) -Euler polynomials; (p, q) -Genocchi polynomials; (p, q) -Fubini numbers and polynomials; (p, q) -Stirling numbers of the second kind.

Mathematics Subject Classification: 11B68, 11B73, 11B75, 11B83, 05A30.

1. Introduction

Throughout this presentation, we use the following standard notions: $\mathbb{N} = \{1, 2, 3, \dots\}$, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. Here as usual \mathbb{Z} denotes the set of integers, \mathbb{R} denotes the set of real numbers and \mathbb{C} denotes the set of complex numbers.

The quantum calculus or q -calculus has several applications in different branches of physics and mathematics. It has attracted serious attention of researchers due to its different applications. At the beginning of 19th century, Jackson initiated and developed the application of the q -calculus (see [7,8]). Later Chakrabarti and Jagannathan defined Jackson (p, q) -derivative as a generalization of q -derivative

(see [3]). Sadjang [13] developed several properties of the (p, q) -derivatives and the (p, q) -integrals and as an application gave two (p, q) -Taylor formulas for polynomials.

For the expedience, we present some definitions and concepts of (p, q) -calculus that were used in this article by assuming as p and q are fixed number such that $0 < p < q \leq 1$.

The (p, q) -derivative of a function f (with respect to x) is defined by

$$D_{p,q}f(x) = \frac{f(px) - f(qx)}{(p - q)x}, \quad (x \neq 0, p \neq q) \quad (1.1)$$

and $D_{p,q}f(0) = f'(0)$, where f' is the ordinary derivative of f .

The (p, q) -derivative operator holds the following properties

$$D_{p,q}(f(x)g(x)) = g(px)D_{p,q}f(x) + f(qx)D_{p,q}g(x), \quad (1.2)$$

and

$$D_{p,q}\left(\frac{f(x)}{g(x)}\right) = \frac{g(qx)D_{p,q}f(x) - f(qx)D_{p,q}g(x)}{g(px)g(qx)}. \quad (1.3)$$

The (p, q) -analogue of $(x + a)^n$ is given by

$$\begin{aligned} (x + a)_{p,q}^n &= (x + a)(px + aq) \cdots (p^{n-2}x + aq^{n-2})(p^{n-1}x + aq^{n-1}), \quad n \geq 1 \\ &= \sum_{k=0}^n \binom{n}{k}_{p,q} p^{\binom{n}{2}} q^{\binom{n-k}{2}} x^k a^{n-k}, \end{aligned} \quad (1.4)$$

where the (p, q) -Gauss binomial coefficients $\binom{n}{k}_{p,q}$ and (p, q) -factorial $[n]_{p,q}!$ are defined by

$$\binom{n}{k}_{p,q} = \frac{[n]_{p,q}!}{[n-k]_{p,q}![k]_{p,q}!}, \quad (n \geq k) \text{ and } [n]_{p,q}! = [n]_{p,q} \cdots [2]_{p,q} [1]_{p,q}, \quad (n \in \mathbb{N}).$$

The (p, q) -exponential functions are defined by

$$e_{p,q}(x) = \sum_{n=0}^{\infty} \frac{p^{\binom{n}{2}} x^n}{[n]_{p,q}!} \quad \text{and} \quad E_{p,q}(x) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} x^n}{[n]_{p,q}!}, \quad (1.5)$$

under condition

$$e_{p,q}(x)E_{p,q}(-x) = 1. \quad (1.6)$$

It follows from (1.1) and (1.5) that

$$D_{p,q}e_{p,q}(x) = e_{p,q}(px) \quad \text{and} \quad D_{p,q}E_{p,q}(x) = E_{p,q}(qx). \quad (1.7)$$

The definite (p, q) -integral of a function f is defined by (see [3])

$$\int_0^a f(x) d_{p,q}x = (p - q)a \sum_{k=0}^{\infty} \frac{p^k}{q^{k+1}} f\left(a \frac{p^k}{q^{k+1}}\right),$$

with the following property

$$\int_a^b f(x) d_{p,q}x = \int_0^b f(x) d_{p,q}x - \int_0^a f(x) d_{p,q}x. \quad (1.8)$$

The generalized (p, q) -Bernoulli, the generalized (p, q) -Euler and the generalized (p, q) -Genocchi numbers and polynomials are defined by means of the following generating function as follows (see [5]):

$$\left(\frac{t}{e_{p,q}(t) - 1}\right)^\alpha e_{p,q}(xt)E_{p,q}(yt) = \sum_{n=0}^{\infty} B_n^{(\alpha)}(x, y : p, q) \frac{t^n}{[n]_{p,q}!}, \quad |t| < 2\pi \quad (1.9)$$

$$\left(\frac{2}{e_{p,q}(t) + 1}\right)^\alpha e_{p,q}(xt)E_{p,q}(yt) = \sum_{n=0}^{\infty} E_n^{(\alpha)}(x, y : p, q) \frac{t^n}{[n]_{p,q}!}, \quad |t| < \pi \quad (1.10)$$

and

$$\left(\frac{2t}{e_{p,q}(t) + 1}\right)^\alpha e_{p,q}(xt)E_{p,q}(yt) = \sum_{n=0}^{\infty} G_n^{(\alpha)}(x, y : p, q) \frac{t^n}{[n]_{p,q}!}, \quad |t| < \pi \quad (1.11)$$

It is clear that

$$B_n^{(\alpha)}(0, 0 : p, q) = B_n^{(\alpha)}(p, q), \quad E_n^{(\alpha)}(0, 0 : p, q) = E_n^{(\alpha)}(p, q),$$

and

$$G_n^{(\alpha)}(0, 0 : p, q) = G_n^{(\alpha)}(p, q), \quad (n \in \mathbb{N}).$$

Geometric polynomials (also known as Fubini polynomials) are defined as follows (see [1]):

$$F_n(x) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} k! x^k, \quad (1.12)$$

where $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}$ is the Stirling number of the second kind (see [9]).

For $x = 1$ in (1.12), we get the n^{th} Fubini number (called Bell number or geometric number) F_n [2,4,14] in defined by

$$F_n(1) = F_n = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} k!. \quad (1.13)$$

The exponential generating functions of geometric polynomials are given by (see [1]):

$$\frac{1}{1 - x(e^t - 1)} = \sum_{n=0}^{\infty} F_n(x) \frac{t^n}{n!}, \quad (1.14)$$

and related to the geometric series (see [1]):

$$\left(x \frac{d}{dx}\right)^m \frac{1}{1-x} = \sum_{k=0}^{\infty} k^m x^k = \frac{1}{1-x} F_m\left(\frac{x}{1-x}\right), \quad |x| < 1.$$

Let us give a short list of these polynomials and numbers as follows:

$$F_0(x) = 1, F_1(x) = x, F_2(x) = x + 2x^2, F_3(x) = x + 6x^2 + 6x^3, F_4(x) = x + 14x^2 + 36x^3 + 24x^4,$$

and

$$F_0 = 1, F_1 = 1, F_2 = 3, F_3 = 13, F_4 = 75.$$

The out line of this paper is as follows: In section 2, we consider generating functions for (p, q) -analogue type of Fubini numbers and polynomials and give some properties of these numbers and polynomials. In section 3, we derive summation formulas for (p, q) -analogue type of Fubini numbers and polynomials. In section , we give relationships for (p, q) -Fubini polynomials associated with (p, q) -Bernoulli polynomials, (p, q) -Euler polynomials and (p, q) -Genocchi polynomials and (p, q) -Stirling numbers of the second kind.

2. (p, q) -analogue type of Fubini numbers and polynomials

In this section, we introduce (p, q) -Fubini polynomials and obtain some basic properties which give new formulas for $F_n(x, y; z : p, q)$.

Definition 2.1. Let $p, q \in \mathbb{C}$ with $0 < |q| < |p| \leq 1$, the three variable (p, q) -Fubini polynomials $F_n(x, y; z : p, q)$ are defined by means of the following generating function:

$$\frac{1}{1 - z(e_{p,q}(t) - 1)} e_{p,q}(xt) E_{p,q}(yt) = \sum_{n=0}^{\infty} F_n(x, y; z : p, q) \frac{t^n}{[n]_{p,q}!}. \quad (2.1)$$

For $x = y = 0$ and $z = 1$ in (2.1), we have

$$F_n(0, 0; 1 : p, q) = F_n(p, q), \quad (2.2)$$

where $F_n(p, q)$ are called the (p, q) -Fubini numbers.

Obviously that

$$\begin{aligned} F_n(0, 0; z : p, q) &= F_n(z : p, q), \\ F_n(x, y; z : p, q)|_{p=1} &= F_n(x, y; z : q), \quad (\text{see [6]}) \\ \lim_{q \rightarrow 1_{p=1}^-} F_n(x, 0 : p, q) &= F_n(x; z), \quad (\text{see [1]}) \\ \lim_{q \rightarrow 1_{p=1}^-} F_n(0, 0; z : p, q) &= F_n(z), \quad (\text{see [1,4]}). \end{aligned}$$

Theorem 2.1. The following series representation for the (p, q) -Fubini polynomials $F_n(x, y; z : p, q)$ holds true:

$$F_n(x, y; z : p, q) = \sum_{m=0}^n \binom{n}{m}_{p,q} F_{n-m}(z : p, q) (x + y)_{p,q}^m. \quad (2.3)$$

Proof. From (2.1), we have

$$\begin{aligned} \sum_{n=0}^{\infty} F_n(x, y; z : p, q) \frac{t^n}{[n]_{p,q}!} &= \frac{1}{1 - z(e_{p,q}(t) - 1)} e_{p,q}(xt) E_{p,q}(yt) \\ &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} F_n(z : p, q) \frac{t^n}{[n]_{p,q}!} (x + y)_{p,q}^m \frac{t^m}{[m]_{p,q}!}. \end{aligned}$$

Replacing n by $n - m$ in above equation and equating the coefficients of same powers of t in both sides of resultant equation, we get representation (2.3). \square

Theorem 2.2. The following summation formula for the (p, q) -Fubini polynomials $F_n(x, y; z : p, q)$ holds true:

$$F_n(x, y; 0 : p, q) = \sum_{m=0}^n \binom{n}{m}_{p,q} p^{\binom{m}{2}} q^{\binom{n-m}{2}} x^m y^{n-m}, \quad (2.4)$$

$$F_n(x, y; z : p, q) = \sum_{m=0}^n \binom{n}{m}_{p,q} p^{\binom{n-m}{2}} F_m(0, y; z : p, q) x^{n-m}, \quad (2.5)$$

$$F_n(x, y; z : p, q) = \sum_{m=0}^n \binom{n}{m}_{p,q} q^{\binom{n-m}{2}} F_m(x, 0; z : p, q) y^{n-m}. \quad (2.6)$$

Proof. Using Eqs (1.4)-(1.6) in generating function (2.1), the proof can be easily proved. So we omit it. \square

Theorem 2.3. For $n \geq 0$, the following formula for (p, q) -Fubini polynomials holds true:

$$(x + y)_{p,q}^n = F_n(x, y; z : p, q) - zF_n(x + 1, y; z : p, q) + zF_n(x, y; z : p, q). \quad (2.7)$$

Proof. We begin with the definition (2.1) and write

$$\begin{aligned} e_{p,q}(xt)E_{p,q}(yt) &= \frac{1 - z(e_{p,q}(t) - 1)}{1 - z(e_{p,q}(t) - 1)} e_{p,q}(xt)E_{p,q}(yt) \\ &= \frac{e_{p,q}(xt)E_{p,q}(yt)}{1 - z(e_{p,q}(t) - 1)} - \frac{z(e_{p,q}(t) - 1)}{1 - z(e_{p,q}(t) - 1)} e_{p,q}(xt)E_{p,q}(yt). \end{aligned}$$

Then using the definition of (1.4) and (2.1), we have

$$\begin{aligned} &\sum_{n=0}^{\infty} (x + y)_{p,q}^n \frac{t^n}{[n]_{p,q}!} \\ &= \sum_{n=0}^{\infty} [F_n(x, y; z : p, q) - zF_n(x + 1, y; z : p, q) + zF_n(x, y; z : p, q)] \frac{t^n}{[n]_{p,q}!}. \end{aligned}$$

Finally, comparing the coefficients of $\frac{t^n}{n!}$, we get (2.7). \square

Theorem 2.4. The following formula for (p, q) -Fubini polynomials holds true:

$$zF_n(x + 1, y; z : p, q) = (1 + z)F_n(x, y; z : p, q) - (x + y)_{p,q}^n. \quad (2.8)$$

Proof. From (2.1), we have

$$\sum_{n=0}^{\infty} [F_n(x + 1, y; z : p, q) - F_n(x, y; z : p, q)] \frac{t^n}{[n]_{p,q}!}$$

$$\begin{aligned}
&= \frac{e_{p,q}(xt)E_{p,q}(yt)}{1 - z(e_{p,q}(t) - 1)}(e_{p,q}(t) - 1) \\
&= \frac{1}{z} \left[\frac{e_{p,q}(xt)E_{p,q}(yt)}{1 - z(e_{p,q}(t) - 1)} - e_{p,q}(xt)E_{p,q}(yt) \right] \\
&= \frac{1}{z} \sum_{n=0}^{\infty} \left[F_n(x, y; z : p, q) - (x + y)_{p,q}^n \right] \frac{t^n}{[n]_{p,q}!}.
\end{aligned}$$

Comparing the coefficients of $\frac{t^n}{n!}$ on both sides, we obtain (2.8). \square

Theorem 2.5. The following recursive formulas for the (p, q) -Fubini polynomials $F_n(x, y; z : p, q)$ holds true:

$$D_{p,q;x}F_n(x, y; z : p, q) = [n]_{p,q}F_{n-1}(px, y; z : p, q), \quad (2.9)$$

$$D_{p,q;y}F_n(x, y; z : p, q) = [n]_{p,q}F_{n-1}(x, qy; z : p, q). \quad (2.10)$$

Proof. Differentiating generating function (2.1) with respect to x and y with the help of Eq. (1.7) and then simplifying with the help of the Cauchy product rule formulas (2.9) and (2.10) are obtained. \square

Theorem 2.6. The following (p, q) -integral is valid

$$\int_a^b F_n(x, y; z : p, q) d_{p,q}x = p \frac{F_{n+1}(\frac{b}{p}, y; z : p, q) - F_{n+1}(\frac{a}{p}, y; z : p, q)}{[n+1]_{p,q}}, \quad (2.11)$$

$$\int_a^b F_n(x, y; z : p, q) d_{p,q}y = p \frac{F_{n+1}(x, \frac{b}{q}; z : p, q) - F_{n+1}(x, \frac{a}{q}; z : p, q)}{[n+1]_{p,q}}. \quad (2.12)$$

Proof. Since

$$\int_a^b \frac{\delta}{\delta_{p,q}x} F_n(x, y; z : p, q) d_{p,q}x = f(b) - f(a), \text{ (see [13]),}$$

in terms of Eq. (2.9) and Eqs (1.7) and (1.8), we arrive at the asserted result

$$\begin{aligned}
\int_a^b \frac{\delta}{\delta_{p,q}x} F_n(x, y; z : p, q) d_{p,q}x &= \frac{p}{[n+1]_{p,q}} \int_a^b F_n\left(\frac{x}{p}, y; z : p, q\right) d_{p,q}x \\
&= p \frac{F_{n+1}(\frac{b}{p}, y; z : p, q) - F_{n+1}(\frac{a}{p}, y; z : p, q)}{[n+1]_{p,q}}.
\end{aligned}$$

The other can be shown using similar method. Therefore, the complete the proof of this theorem. \square

3. Main results

First, we prove the following result involving the (p, q) -Fubini polynomials $F_n(x, y; z : p, q)$ by using series rearrangement techniques and considered its special case:

Theorem 3.1. The following summation formula for (p, q) -Fubini polynomials $F_n(x, y; z : p, q)$ holds true:

$$F_{k+l}(w, y; z : p, q) = \sum_{n,s=0}^{k,l} \binom{k}{n}_{p,q} \binom{l}{s}_{p,q} p^{\binom{n+s}{2}} (w-x)^{n+s} F_{k+l-n-s}(x, y; z : p, q). \quad (3.1)$$

Proof. Replacing t by $t + u$ in (2.1) and then using the formula [12, p.52(2)]:

$$\sum_{N=0}^{\infty} f(N) \frac{(x+y)^N}{N!} = \sum_{n,m=0}^{\infty} f(n+m) \frac{x^n y^m}{n! m!}, \quad (3.2)$$

in the resultant equation, we find the following generating function for the (p, q) -Fubini polynomials $F_n(x, y; z : p, q)$:

$$\begin{aligned} & \frac{1}{1 - z(e_{p,q}(t+u) - 1)} E_{p,q}(y(t+u)) \\ &= e_{p,q}(-x(t+u)) \sum_{k,l=0}^{\infty} F_{k+l}(x, y; z : p, q) \frac{t^k}{[k]_{p,q}!} \frac{u^l}{[l]_{p,q}!}, \quad (\text{see [10,11]}). \end{aligned} \quad (3.3)$$

Replacing x by w in the above equation and equating the resultant equation to the above equation, we find

$$\begin{aligned} & e_{p,q}((w-x)(t+u)) \sum_{k,l=0}^{\infty} F_{k+l}(x, y; z : p, q) \frac{t^k}{[k]_{p,q}!} \frac{u^l}{[l]_{p,q}!} \\ &= \sum_{k,l=0}^{\infty} F_{k+l}(w, y; z : p, q) \frac{t^k}{[k]_{p,q}!} \frac{u^l}{[l]_{p,q}!}. \end{aligned} \quad (3.4)$$

On expanding exponential function (3.4) gives

$$\begin{aligned} & \sum_{N=0}^{\infty} \frac{[(w-x)(t+u)]^N p^{\binom{N}{2}}}{[N]_{p,q}!} \sum_{k,l=0}^{\infty} F_{k+l}(x, y; z : p, q) \frac{t^k}{[k]_{p,q}!} \frac{u^l}{[l]_{p,q}!} \\ &= \sum_{k,l=0}^{\infty} F_{k+l}(w, y; z : p, q) \frac{t^k}{[k]_{p,q}!} \frac{u^l}{[l]_{p,q}!}, \end{aligned} \quad (3.5)$$

which on using formula (3.2) in the first summation on the left hand side becomes

$$\begin{aligned} & \sum_{n,s=0}^{\infty} \frac{(w-x)^{n+s} t^n u^s p^{\binom{n+s}{2}}}{[n]_{p,q}! [s]_{p,q}!} \sum_{k,l=0}^{\infty} F_{k+l}(x, y; z : p, q) \frac{t^k}{[k]_{p,q}!} \frac{u^l}{[l]_{p,q}!} \\ &= \sum_{k,l=0}^{\infty} F_{k+l}(w, y; z : p, q) \frac{t^k}{[k]_{p,q}!} \frac{u^l}{[l]_{p,q}!}. \end{aligned} \quad (3.6)$$

Now replacing k by $k - n$, l by $l - s$ and using the lemma ([12, p.100(1)]):

$$\sum_{k=0}^{\infty} \sum_{n=0}^{\infty} A(n, k) = \sum_{k=0}^{\infty} \sum_{n=0}^k A(n, k - n), \quad (3.7)$$

in the l.h.s. of (3.6), we find

$$\begin{aligned} \sum_{k,l=0}^{\infty} \sum_{n,s=0}^{k,l} \frac{(w-x)^{n+s} p^{\binom{n+s}{2}}}{[n]_{p,q}! [s]_{p,q}!} F_{k+l-n-s}(x, y; z; p, q) \frac{t^k}{(k-n)_{p,q}!} \frac{u^l}{(l-s)_{p,q}!} \\ = \sum_{k,l=0}^{\infty} F_{k+l}(w, y; z; p, q) \frac{t^k}{[k]_{p,q}!} \frac{u^l}{[l]_{p,q}!}. \end{aligned} \quad (3.8)$$

Finally, on equating the coefficients of the like powers of t and u in the above equation, we get the assertion (3.1) of Theorem 3.1. \square

Remark 3.1. Taking $l = 0$ in assertion (3.1) of Theorem 3.1, we deduce the following consequence of Theorem 3.1.

Corollary 3.1. The following summation formula for (p, q) -Fubini polynomials $F_n(x, y; z; p, q)$ holds true:

$$F_k(w, y; z; p, q) = \sum_{n=0}^k \binom{k}{n}_{p,q} p^{\binom{n+s}{2}} (w-x)^n F_{k-n}(x, y; z; p, q). \quad (3.9)$$

Remark 3.2. Replacing w by $w + x$ in (3.9), we obtain

$$F_k(w + x, y; z; p, q) = \sum_{n=0}^k \binom{k}{n}_{p,q} p^{\binom{n+s}{2}} w^n F_{k-n}(x, y; z; p, q). \quad (3.10)$$

Theorem 3.2. The following summation formula for (p, q) -Fubini polynomials $F_n(x, y; z; p, q)$ holds true:

$$\begin{aligned} F_n(w, u; z; p, q) F_m(W, U; Z; p, q) = \sum_{r,k=0}^{n,m} \binom{n}{r}_{p,q} \binom{m}{k}_{p,q} (w-x+u-y)_{p,q}^r \\ \times F_{n-r}(x, y; z; p, q) (W-X+U-Y)_{p,q}^k F_{m-k}(X, Y; Z; p, q). \end{aligned} \quad (3.11)$$

Proof. Consider the product of the (p, q) -Fubini polynomials, we can be written as generating function (2.1) in the following form:

$$\begin{aligned} \frac{1}{1 - z(e_{p,q}(t) - 1)} e_{p,q}(xt) E_{p,q}(yt) \frac{1}{1 - Z(e_{p,q}(T) - 1)} e_{p,q}(XT) E_{p,q}(YT) \\ = \sum_{n=0}^{\infty} F_n(x, y; z; p, q) \frac{t^n}{[n]_{p,q}!} \sum_{m=0}^{\infty} F_m(X, Y; Z; p, q) \frac{T^m}{[m]_{p,q}!}. \end{aligned} \quad (3.12)$$

Replacing x by w , y by u , X by W and Y by U in (3.12) and equating the resultant to itself,

$$\begin{aligned} & \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} F_n(w, u; z : p, q) F_m(W, U; Z : p, q) \frac{t^n}{[n]_{p,q}!} \frac{T^m}{[m]_{p,q}!} \\ &= e_{p,q}((w-x)t) E_{p,q}((u-y)t) e_{p,q}((W-X)T) E_{p,q}((U-Y)t) \\ & \times \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} F_n(x, y; z : p, q) F_m(X, Y; Z : p, q) \frac{t^n}{[n]_{p,q}!} \frac{T^m}{[m]_{p,q}!}, \end{aligned}$$

which on using the generating function (3.7) in the exponential on the r.h.s., becomes

$$\begin{aligned} & \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} F_n(w, u; z : p, q) F_m(W, U; Z : p, q) \frac{t^n}{[n]_{p,q}!} \frac{T^m}{[m]_{p,q}!} \\ &= \sum_{n,r=0}^{\infty} (w-x+u-y)_{p,q}^r F_n(x, y; z) \frac{t^{n+r}}{[n]_{p,q}! [r]_{p,q}!} \\ & \times \sum_{m,k=0}^{\infty} (W-X+U-Y)_{p,q}^k F_m(X, Y; Z) \frac{T^{m+k}}{[m]_{p,q}! [k]_{p,q}!}. \end{aligned} \quad (3.13)$$

Finally, replacing n by $n-r$ and m by $m-k$ and using Eq. (3.7) in the r.h.s. of the above equation and then equating the coefficients of like powers of t and T , we get assertion (3.11) of Theorem 3.2. \square

Remark 3.3. Replacing u by y and U by Y in assertion (3.11) of Theorem 3.2, we deduce the the following consequence of Theorem 3.2.

Corollary 3.2. The following summation formula for (p, q) -Fubini polynomials $F_n(x, y; z : p, q)$ holds true:

$$\begin{aligned} F_n(w, y; z : p, q) F_m(W, Y; Z : p, q) &= \sum_{r,k=0}^{n,m} \binom{n}{r}_{p,q} \binom{m}{k}_{p,q} (w-x)_{p,q}^r F_{n-r}(x, u; z : p, q) \\ & \times (W-X)_{p,q}^k F_{m-k}(X, U; Z : p, q). \end{aligned} \quad (3.14)$$

Theorem 3.3. The following summation formula for (p, q) -Fubini polynomials $F_n(x, y; z : p, q)$ holds true:

$$F_n(x+1, y; z : p, q) = \sum_{r=0}^n \binom{n}{r}_{p,q} F_{n-r}(x, y; z : p, q). \quad (3.15)$$

Proof. Using the generating function (2.1), we have

$$\begin{aligned} & \sum_{n=0}^{\infty} F_n(x+1, y; z : p, q) \frac{t^n}{[n]_{p,q}!} - \sum_{n=0}^{\infty} F_n(x, y; z : p, q) \frac{t^n}{[n]_{p,q}!} \\ &= \left(\frac{1}{1-z(e_{p,q}(t)-1)} \right) (e_{p,q}(t) - 1) e_{p,q}(xt) E_{p,q}(yt) \end{aligned}$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty} F_n(x, y; z : p, q) \frac{t^n}{[n]_{p,q}!} \left(\sum_{r=0}^{\infty} \frac{t^r}{[r]_{p,q}!} - 1 \right) \\
&= \sum_{n=0}^{\infty} F_n(x, y; z : p, q) \frac{t^n}{[n]_{p,q}!} \sum_{r=0}^{\infty} \frac{t^r}{[r]_{p,q}!} - \sum_{n=0}^{\infty} F_n(x, y; z : p, q) \frac{t^n}{[n]_{p,q}!} \\
&= \sum_{n=0}^{\infty} \sum_{r=0}^n \binom{n}{r}_{p,q} F_{n-r}(x, y; z : p, q) \frac{t^n}{[n]_{p,q}!} - \sum_{n=0}^{\infty} F_n(x, y; z : p, q) \frac{t^n}{[n]_{p,q}!}.
\end{aligned}$$

Finally, equating the coefficients of the like powers of t on both sides, we get (3.15). \square

Theorem 3.4. For $n \geq 0$ and $z_1 \neq z_2$, the following formula for (p, q) -Fubini polynomials holds true:

$$\begin{aligned}
&\sum_{k=0}^n \binom{n}{k}_{p,q} F_{n-k}(x_1, y_1; z_1 : p, q) F_k(x_2, y_2; z_2 : p, q) \\
&= \frac{z_2 F_n(x_1 + x_2, y_1 + y_2; z_2 : p, q) - z_1 F_n(x_1 + x_2, y_1 + y_2; z_1 : p, q)}{z_2 - z_1}. \tag{3.16}
\end{aligned}$$

Proof. The products of (2.1) can be written as

$$\begin{aligned}
&\sum_{n=0}^{\infty} \sum_{k=0}^{\infty} F_n(x_1, y_1; z_1 : p, q) F_k(x_2, y_2; z_2 : p, q) \frac{t^n}{[n]_{p,q}!} \frac{t^k}{[k]_{p,q}!} \\
&= \frac{e_{p,q}(x_1 t) E_{p,q}(y_1 t)}{1 - z_1 (e_{p,q}(t) - 1)} \frac{e_{p,q}(x_2 t) E_{p,q}(y_2 t)}{1 - z_2 (e_{p,q}(t) - 1)} \\
&= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k}_{p,q} F_{n-k}(x_1, y_1; z_1 : p, q) F_k(x_2, y_2; z_2 : p, q) \right) \frac{t^n}{[n]_{p,q}!} \\
&= \frac{z_2}{z_2 - z_1} \frac{e_{p,q}[(x_1 + x_2)t] E_{p,q}[(y_1 + y_2)t]}{1 - z_1 (e_{p,q}(t) - 1)} - \frac{z_1}{z_2 - z_1} \frac{e_{p,q}[(x_1 + x_2)t] E_{p,q}[(y_1 + y_2)t]}{1 - z_2 (e_{p,q}(t) - 1)} \\
&= \sum_{n=0}^{\infty} \left(\frac{z_2 F_n(x_1 + x_2, y_1 + y_2; z_2 : p, q) - z_1 F_n(x_1 + x_2, y_1 + y_2; z_1 : p, q)}{z_2 - z_1} \right) \frac{t^n}{[n]_{p,q}!}.
\end{aligned}$$

By equating the coefficients of $\frac{t^n}{[n]_{p,q}!}$ on both sides, we get (3.16). \square

Theorem 3.5. The following relation for the (p, q) -Fubini polynomials $F_n(x, y; z : p, q)$ holds true:

$$(1 + z) F_n(x, y; z : p, q) = z \sum_{k=0}^n \binom{n}{k}_{p,q} F_{n-k}(x, y; z : p, q) + (x + y)_{p,q}^n. \tag{3.17}$$

Proof. Consider the following identity

$$\frac{1 + z}{(1 - z(e_{p,q}(t) - 1))z e_{p,q}(t)} = \frac{1}{1 - z(e_{p,q}(t) - 1)} + \frac{1}{z e_{p,q}(t)}.$$

Evaluating the following fraction using above identity, we find

$$\begin{aligned} \frac{(1+z)e_{p,q}(xt)E_{p,q}(yt)}{(1-z(e_{p,q}(t)-1))ze_{p,q}(t)} &= \frac{e_{p,q}(xt)E_{p,q}(yt)}{1-z(e_{p,q}(t)-1)} + \frac{e_{p,q}(xt)E_{p,q}(yt)}{ze_{p,q}(t)} \\ &= (1+z) \sum_{n=0}^{\infty} F_n(x, y; z : p, q) \frac{t^n}{[n]_{p,q}!} \\ &= z \sum_{n=0}^{\infty} F_n(x, y; z : p, q) \frac{t^n}{[n]_{p,q}!} \sum_{k=0}^{\infty} \frac{t^k}{[k]_{p,q}!} - \sum_{n=0}^{\infty} (x+y)^n_{p,q} \frac{t^n}{[n]_{p,q}!}. \end{aligned}$$

Applying the Cauchy product rule in the above equation and then equating the coefficients of like powers of t in both sides of the resultant equation, assertion (3.17) follows. \square

4. Relationship between (p, q) -Bernoulli, (p, q) -Euler and (p, q) -Genocchi polynomials and (p, q) -Stirling numbers of the second kind

In this section, we prove some relationships for (p, q) -Fubini polynomials related to (p, q) -Bernoulli polynomials, (p, q) -Euler polynomials and (p, q) -Genocchi polynomials and (p, q) -Stirling numbers of the second kind. We start a following theorem.

Theorem 4.1. Each of the following relationships holds true:

$$\begin{aligned} &F_n(x, y; z : p, q) \\ &= \sum_{s=0}^{n+1} \binom{n+1}{s}_{p,q} \left[\sum_{k=0}^s \binom{s}{k}_{p,q} B_{s-k}(x : p, q) p^{\binom{k}{2}} - B_s(x : p, q) \right] \frac{F_{n+1-s}(0, y; z : p, q)}{[n+1]_{p,q}}, \end{aligned} \quad (4.1)$$

where $B_n(x : p, q)$ are called the (p, q) -Bernoulli polynomials.

Proof. By using definition (2.1), we have

$$\begin{aligned} &\left(\frac{1}{1-z(e_{p,q}(t)-1)} \right) e_{p,q}(xt)E_{p,q}(yt) \\ &= \left(\frac{1}{1-z(e_{p,q}(t)-1)} \right) \frac{t}{e_{p,q}(t)-1} \frac{e_{p,q}(t)-1}{t} e_{p,q}(xt)E_{p,q}(yt) \\ &= \frac{1}{t} \sum_{n=0}^{\infty} \left(\sum_{k=0}^s \binom{s}{k}_{p,q} B_{s-k}(x : p, q) p^{\binom{k}{2}} \right) \frac{t^s}{[s]_{p,q}!} \sum_{n=0}^{\infty} F_n(0, y; z : p, q) \frac{t^n}{[n]_{p,q}!} \\ &\quad - \frac{1}{t} \sum_{s=0}^{\infty} B_s(x : p, q) \frac{t^s}{[s]_{p,q}!} \sum_{n=0}^{\infty} F_n(0, y; z : p, q) \frac{t^n}{[n]_{p,q}!} \\ &= \frac{1}{t} \sum_{n=0}^{\infty} \left[\sum_{s=0}^n \binom{n}{s}_{p,q} \sum_{k=0}^s \binom{s}{k}_{p,q} B_{s-k}(x : p, q) p^{\binom{k}{2}} \right] F_{n-s}(0, y; z : p, q) \frac{t^n}{[n]_{p,q}!} \\ &\quad - \frac{1}{t} \sum_{n=0}^{\infty} \left[\sum_{s=0}^n \binom{n}{s}_{p,q} B_s(x : p, q) \right] F_{n-s}(0, y; z : p, q) \frac{t^n}{[n]_{p,q}!}. \end{aligned}$$

By using Cauchy product and comparing the coefficients of $\frac{t^n}{[n]_{p,q}!}$, we arrive at the required result (4.1). \square

Theorem 4.2. Each of the following relationships holds true:

$$\begin{aligned}
 & F_n(0, y; z; u : p, q) \\
 &= \sum_{s=0}^n \binom{n}{s}_{p,q} \left[\sum_{k=0}^s \binom{s}{k}_{p,q} E_{s-k}(x : p, q) p^{\binom{k}{2}} + E_s(x : p, q) \right] \frac{F_{n-s}(0, y; z : p, q)}{[2]_{p,q}}, \quad (4.2)
 \end{aligned}$$

where $E_n(x; p, q)$ are called the (p, q) -Euler polynomials.

Proof. By using definition (2.1), we have

$$\begin{aligned}
 & \left(\frac{1}{1 - z(e_{p,q}(t) - 1)} \right) e_{p,q}(xt) E_{p,q}(yt) \\
 &= \left(\frac{1}{1 - z(e_{p,q}(t) - 1)} \right) \frac{[2]_{p,q} e_{p,q}(t) + 1}{e_{p,q}(t) + 1} \frac{1}{[2]_{p,q}} e_{p,q}(xt) E_{p,q}(yt) \\
 &= \frac{1}{[2]_{p,q}} \left[\sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k}_{p,q} E_{n-k}(x : p, q) p^{\binom{k}{2}} \right) \frac{t^n}{[n]_{p,q}!} + \sum_{n=0}^{\infty} E_n(x : p, q) \frac{t^n}{[n]_{p,q}!} \right] \\
 &\times \sum_{n=0}^{\infty} F_n(0, y; z : p, q) \frac{t^n}{[n]_{p,q}!} \\
 &= \frac{1}{[2]_{p,q}} \sum_{n=0}^{\infty} \left[\sum_{s=0}^n \binom{n}{s}_{p,q} \sum_{k=0}^s \binom{s}{k}_{p,q} E_{s-k}(x : p, q) p^{\binom{k}{2}} + \sum_{s=0}^n \binom{n}{s}_{p,q} E_s(x : p, q) \right] \\
 &\times F_{n-s}(0, y; z : p, q) \frac{t^n}{[n]_{p,q}!}.
 \end{aligned}$$

Comparing the coefficients of $\frac{t^n}{[n]_{p,q}!}$, we arrive at the desired result (4.2). \square

Theorem 4.3. Each of the following relationships holds true:

$$\begin{aligned}
 & F_n(x, y; z : p, q) \\
 &= \sum_{s=0}^n \binom{n+1}{s}_{p,q} \left[\sum_{k=0}^s \binom{s}{k}_{p,q} G_{s-k}(x : p, q) p^{\binom{k}{2}} + G_s(x : p, q) \right] \frac{F_{n-s}(0, y; z : p, q)}{[2]_{p,q} [n+1]_{p,q}}, \quad (4.3)
 \end{aligned}$$

where $G_n(x; p, q)$ are called the (p, q) -Genocchi polynomials.

Proof. By using definition (2.1), we have

$$\begin{aligned}
& \left(\frac{1}{1 - z(e_{p,q}(t) - 1)} \right) e_{p,q}(xt) E_{p,q}(yt) \\
&= \left(\frac{1}{1 - z(e_{p,q}(t) - 1)} \right) e_{p,q}(xt) E_{p,q}(yt) \frac{[2]_{p,q} t}{e_{p,q}(t) + 1} \frac{e_{p,q}(t) + 1}{[2]_{p,q} t} e_{p,q}(xt) E_{p,q}(yt) \\
&= \frac{1}{[2]_{p,q} t} \left[\sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k}_{p,q} G_{n-k}(x : p, q) p^{\binom{k}{2}} \right) \frac{t^n}{[n]_{p,q}!} + \sum_{n=0}^{\infty} G_n(x : p, q) \frac{t^n}{[n]_{p,q}!} \right] \\
&\times \sum_{n=0}^{\infty} F_n(0, y; z : p, q) \frac{t^n}{[n]_{p,q}!} \\
&= \frac{1}{[2]_{p,q}} \sum_{n=0}^{\infty} \left[\sum_{s=0}^n \binom{n}{s}_{p,q} \sum_{k=0}^s \binom{s}{k}_{p,q} G_{s-k}(x : p, q) p^{\binom{k}{2}} + \sum_{s=0}^n \binom{n}{s}_{p,q} G_s(x : p, q) \right] \\
&\times F_{n+1-s}(0, y; z : p, q) \frac{t^n}{[n+1]_{p,q}!}.
\end{aligned}$$

Comparing the coefficients of $\frac{t^n}{[n]_{p,q}!}$, then we have the asserted result (4.3). \square

Theorem 4.4. For $n \geq 0$, the following formula for (p, q) -Fubini polynomials holds true:

$$F_n(x, y; z : p, q) = \sum_{l=0}^n \binom{n}{l}_{p,q} (x+y)_{p,q}^{n-l} \sum_{k=0}^l z^k k! S_2(l, k : p, q). \quad (4.4)$$

Proof. From (2.1), we have

$$\begin{aligned}
\sum_{n=0}^{\infty} F_n(x, y; z : p, q) \frac{t^n}{[n]_{p,q}!} &= \frac{1}{1 - z(e_{p,q}(t) - 1)} e_{p,q}(xt) E_{p,q}(yt) \\
&= e_{p,q}(xt) E_{p,q}(yt) \sum_{k=0}^{\infty} z^k (e_{p,q}(t) - 1)^k \\
&= e_{p,q}(xt) E_{p,q}(yt) \sum_{k=0}^{\infty} z^k \sum_{l=k}^{\infty} k! S_2(l, k : p, q) \frac{t^l}{[l]_{p,q}!} \\
&= \sum_{n=0}^{\infty} (x+y)_{p,q}^n \frac{t^n}{[n]_{p,q}!} \sum_{l=0}^{\infty} z^l \sum_{k=0}^l k! S_2(l, k : p, q) \frac{t^l}{[l]_{p,q}!}.
\end{aligned}$$

Replacing n by $n - l$ in above equation, we get

$$\begin{aligned}
& \sum_{n=0}^{\infty} F_n(x, y; z : p, q) \frac{t^n}{[n]_{p,q}!} \\
&= \sum_{n=0}^{\infty} \left(\sum_{l=0}^n \binom{n}{l}_{p,q} (x+y)_{p,q}^{n-l} \sum_{k=0}^l z^k k! S_2(l, k : p, q) \right) \frac{t^n}{[n]_{p,q}!}.
\end{aligned}$$

Comparing the coefficients of $\frac{t^n}{[n]_{p,q}!}$ in both sides, we get (4.4). \square

Theorem 4.5. For $n \geq 0$, the following formula for (p, q) -Fubini polynomials holds true:

$$F_n(x + r, y; z : p, q) = \sum_{l=0}^n \binom{n}{l}_{p,q} (x + y)_{p,q}^{n-l} \sum_{k=0}^l z^k k! S_2(l + r, k + r : p, q). \quad (4.5)$$

Proof. Replacing x by $x + r$ in (2.1), we have

$$\begin{aligned} \sum_{n=0}^{\infty} F_n(x + r, y; z : p, q) \frac{t^n}{[n]_{p,q}!} &= \frac{1}{1 - z(e_{p,q}(t) - 1)} e_{p,q}((x + r)t) E_{p,q}(yt) \\ &= e_{p,q}(xt) E_{p,q}(yt) e_{p,q}(rt) \sum_{k=0}^{\infty} z^k (e_{p,q}(t) - 1)^k \\ &= e_{p,q}(xt) E_{p,q}(yt) e_{p,q}(rt) \sum_{k=0}^{\infty} z^k \sum_{l=k}^{\infty} k! S_2(l, k : p, q) \frac{t^l}{[l]_{p,q}!} \\ &= \sum_{n=0}^{\infty} (x + y)_{p,q}^n \frac{t^n}{[n]_{p,q}!} \sum_{l=0}^{\infty} z^l \sum_{k=0}^l k! S_2(l + r, k + r : p, q) \frac{t^l}{[l]_{p,q}!}. \end{aligned}$$

Replacing n by $n - l$ in above equation, we get

$$\begin{aligned} &\sum_{n=0}^{\infty} F_n(x + r, y; z : p, q) \frac{t^n}{[n]_{p,q}!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{l=0}^n \binom{n}{l}_{p,q} (x + y)_{p,q}^{n-l} \sum_{k=0}^l z^k k! S_2(l + r, k + r : p, q) \right) \frac{t^n}{[n]_{p,q}!}. \end{aligned}$$

Comparing the coefficients of $\frac{t^n}{[n]_{p,q}!}$ in both sides, we get (4.5). \square

Acknowledgments

The author K.S. Nisar express his thanks to the Deanship of Scientific Research (DSR), Prince Sattam bin Abdulaziz University, Saudi Arabia for providing facilities and support.

Conflict of interest

The authors declare no conflict of interest.

References

1. K. N. Boyadzhiev, *A series transformation formula and related polynomials*, Int. J. Math., **23** (2005), 3849–3866.
2. L. Carlitz, *Eulerian numbers and polynomials*, Mathematics Magazine, **32** (1959), 164–171.
3. R. Chakrabarti, R. Jagannathan, *A (p, q) -oscillator realization of two parameter quantum algebras*, Journal of Physics A: Mathematical and General, **24** (1991), L711–L718.

4. A. Dil, V. Kurt, *Investing geometric and exponential polynomials with Euler-Seidel matrices*, Journal of Integer Sequences, **14** (2011), 1–12.
5. U. Duran, M. Acikgoz, *Apostal type (p, q) -Bernoulli, (p, q) -Euler and (p, q) -Genocchi polynomials and numbers*, Comput. Appl. Math., **8** (2017), 7–30.
6. U. Duran, S. Araci, M. Acikgoz, *A note on q -Fubini polynomials*, Advanced Studies in Contemporary Mathematics, **29** (2019), 211–224.
7. F. H. Jackson, *On q -definite integrals*, Quart. J. Pure Appl. Math., **41** (1910), 193–203.
8. F. H. Jackson, *On q -functions and a certain difference operator*, Transactions of the Royal Society of Edinburgh, **46** (1909), 253–281.
9. L. Kargin, *Some formulae for products of Fubini polynomials with applications*, arXiv:1701.01023, 2016.
10. W. A. Khan, I. A. Khan, J. Y. Kang, *On higher order (p, q) -Frobenius Genocchi numbers and polynomials*, J. Appl. Math. and Informatics, **37**(2019), 297–307.
11. W. A. Khan, I. A. Khan, *A note on (p, q) analogue type of Frobenius Genocchi numbers and polynomials*, East Asian Math. J., **36** (2020), 013–024.
12. H. M. Srivastava, H. L. Manocha, *A treatise on generating functions*, Ellis Horwood, New York, 1984.
13. P. N. Sadjang, *On the fundamental theorem of (p, q) -calculus and some (p, q) -Taylor formulas*, arXiv:1309.3934, 2013.
14. S. M. Tanny, *On some numbers related to Bell numbers*, Canad. Math. Bull., **17** (1974), 733–738.



© 2020 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)