


RESEARCH

Open Access



Oblique explicit wave solutions of the fractional biological population (BP) and equal width (EW) models

Abdel-Haleem Abdel-Aty^{1,2*} , Mostafa M.A. Khater^{3,4}, Dumitru Baleanu^{5,6,7}, S.M. Abo-Dahab^{8,9},
Jamel Bouslimi^{10,11} and M. Omri¹²

*Correspondence:

amabdelaty@ub.edu.sa

¹Department of Physics, College of Sciences, University of Bisha, P.O. Box 344, Bisha 61922, Saudi Arabia

²Physics Department, Faculty of Science, Al-Azhar University, Assiut 71524, Egypt

Full list of author information is available at the end of the article

Abstract

This research uses the extended $\exp(-\varphi(\vartheta))$ -expansion and the Jacobi elliptical function methods to obtain a fashionable explicit format for solutions to the fragmented biological population and the same width models that depict popular logistics because of deaths or births. In mathematical terminology, the linear, hyperbolic, and trigonometric equation solutions that have been found describe several innovative aspects from the two models. Sketching these solutions in different types is used to give them more details. The accuracy and performance of the method adopted show their ability to be applied to various nonlinear developmental equations.

MSC: 35E05; 35C08; 35Q51; 37L50

Keywords: Computational recent schemes; Fractional operator; Fractional nonlinear BP equation; Fractional nonlinear EW equation

1 Introduction

Previously, a system of nonlinear evolutionary equations has been used to formulate a fraction of a population in specific fields [1–5]. In numerous distinct branches of science such as mathematics, chemistry, biology, ecology, chaos syncing, mechanics engineering, physics and anomalous spreads, and so on [6–8], many researchers have investigated analytical, semiautomatic, and numerical solutions of fractional models. Such phenomena have been modeled by the fractional mathematical models based on experimental results to demonstrate their nonlocal properties, where this form of property [9–13] cannot be expressed by nonlinear partial differential equations with an integer order.

Based on the ability to form multiple complex phenomena in diverse fields such as biology, plasma physics, hydrodynamics, fluid mechanics, optics, and so forth, several precise and computational schemes such as [14–23] have been developed. In the treatment to these problems, electronic and technical developments are known to be of essential value across derivative structures. These systems have been recently considered to be basic tools in the development of various waveform travel formulas such as [24–30]. These are com-

© The Author(s) 2020. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

plex phenomena. In the nonlinear partial differential equation (NLPDE), several scientists have struggled to extract and formulate several complicated anomalies in an integer order [31, 32]. Therefore, fractional is assumed to be an effective solution for this question when a nonlocal property is found that is not NLPDE dependent with [33, 34] an integer order. This makes many fractional models and definitions of derivatives represented and formulated as in [35–47].

In this research, we study two primary models in biological science. These models are named with the fractional BP model, and fractional EW equation is given by

- Fractional BP model [48–51]:

This paradigm illustrates community dynamics and is given by

$$D_t^\kappa \mathcal{H} = D_{xx}^{2\kappa} \mathcal{H}^2 + D_{yy}^{2\kappa} \mathcal{H}^2 + v(\mathcal{H}^2 - s), \tag{1}$$

where \mathcal{H} is the function of the population density, while $v(\mathcal{H}^2 - s)$ represents the population logistics according to deaths and births. Additionally, v, s, κ ($0 < \kappa \leq 1$) are arbitrary constants.

The population model describes the number of organisms of the same species (human, animal, and plant) living simultaneously in a particular geographic area with interbreeding capacity.

- Fractional EW equation [52–55]:

It is an alternative form of nonlinear dispersive waves first introduced by Morrison et al. and formulated as follows:

$$D_t^\kappa \Lambda + 2h\Lambda D_x^\kappa \Lambda - rD_{xxt}^{3\kappa} \Lambda = 0, \tag{2}$$

where h, r, κ ($0 < \kappa \leq 1$) are arbitrary constants. Without deformation, waves may spread in a wave medium that is non-dispersive. Electromagnetic waves with unbounded free space are both non-dispersive and non-dissipative, thus can spread over astronomical distances. Sound waves in air are also virtually non-dispersive in the ultrasound range. If not, if high-frequency notes (e.g. piccolo) and low-frequency notes (e.g. basis) spread at different speeds, they could reach the ears at different times. However, the majority of the waves in material media are scattering, and initially established wave forms will change so that wave power can spread or disperse more spatially.

Implementation of the following conformable derivative definitions (for further definition and properties of the conformable fractional derivatives, see the [Appendix](#)) on Eqs. (1) and (2) with the following respective order $\mathcal{H}(x, y, t) = \mathcal{H}(\vartheta)$, $\vartheta = \varrho \frac{x^\kappa}{\kappa} + i\varrho \frac{y^\kappa}{\kappa} + \frac{ct^\kappa}{\kappa}$, $\Lambda(x, t) = \Lambda(\vartheta)$, $\vartheta = \frac{x^\kappa}{\kappa} + \frac{ct^\kappa}{\kappa}$, where ϱ, c, κ are arbitrary constants, transforms the fractional PDE into integer order ODE which are given by

$$c\mathcal{H}' - v\mathcal{H}^2 + vs = 0, \tag{3}$$

$$c\Lambda + h\Lambda^2 - rc\Lambda'' = 0. \tag{4}$$

The remaining sections in our research paper are ordered as follows: Sect. 2 manipulates the extended $\exp(-\varphi(\vartheta))$ -expansion method [56–60] and the Jacobi elliptical function method [61–63] to procure novel solitary solutions of both suggested models. Section 5 gives the conclusion of this research.

2 Applications

This section applies the extended $\exp(-\varphi(\vartheta))$ -expansion and the Jacobi elliptical function techniques on the considered fractional models.

Balancing the terms in Eqs. (3), (4) to get the balance value of each of them leads to respectively $n + 1 = m$ & $n + 2 = m$, where n, m are arbitrary constants. Supposing the value of $n = 1$ and according to the general solution that is suggested by the extended $\exp(-\varphi(\vartheta))$ -expansion method and the Jacobi elliptical function method, the general solution of Eqs. (3) is given respectively by

$$\mathcal{H}(\vartheta) = \frac{\sum_{i=0}^m a_i e^{-i\varphi(\vartheta)}}{\sum_{j=0}^n b_j e^{-j\varphi(\vartheta)}} = \frac{a_1 e^{-\varphi(\vartheta)} + a_2 e^{-2\varphi(\vartheta)} + a_0}{b_1 e^{-\varphi(\vartheta)} + b_0}, \tag{5}$$

$$\mathcal{H}(\vartheta) = \sum_{i=1}^n a_i \phi(\vartheta)^i + \sum_{i=1}^n b_i \phi(\vartheta)^{-i} + a_0 = a_1 \phi(\vartheta) + a_0 + \frac{b_1}{\phi(\vartheta)}, \tag{6}$$

while the general solution of Eqs. (4) is given by

$$\Lambda(\vartheta) = \frac{\sum_{i=0}^m a_i e^{-i\varphi(\vartheta)}}{\sum_{j=0}^n b_j e^{-j\varphi(\vartheta)}} = \frac{a_1 e^{-\varphi(\vartheta)} + a_2 e^{-2\varphi(\vartheta)} + a_3 e^{-3\varphi(\vartheta)} + a_0}{b_1 e^{-\varphi(\vartheta)} + b_0}, \tag{7}$$

$$\begin{aligned} \Lambda(\vartheta) &= \sum_{i=1}^n a_i \phi(\vartheta)^i + \sum_{i=1}^n b_i \phi(\vartheta)^{-i} + a_0 \\ &= a_2 \phi(\vartheta)^2 + a_1 \phi(\vartheta) + a_0 + \frac{b_2}{\phi(\vartheta)^2} + \frac{b_1}{\phi(\vartheta)}, \end{aligned} \tag{8}$$

where a_i, b_j ($i, j = 0, 1, 2, \dots$). Additionally, $\varphi(\vartheta)$ is the solution function of $[\varphi'(\vartheta) = \chi + \gamma e^{\varphi(\vartheta)} + \frac{1}{e^{\varphi(\vartheta)}} \ \& \ \phi'(\vartheta) = \sqrt{p\phi(\vartheta)^2 + q\phi(\vartheta)^4 + \rho}]$, where χ, γ, r, p, q are arbitrary constants.

2.1 Fractional BP model

2.1.1 Extended $\exp(-\varphi(\vartheta))$ -expansion method

Employing Eqs. (5) and (3) in the framework of the extended $\exp(-\varphi(\vartheta))$ -expansion method to solve Eq. (3) yields the following.

Family I

$$\left[a_0 \rightarrow \frac{b_0 \chi \sqrt{s}}{\sqrt{\chi^2 - 4\gamma}}, a_1 \rightarrow \frac{2b_0 \sqrt{s}}{\sqrt{\chi^2 - 4\gamma}} + \frac{b_1 \chi \sqrt{s}}{\sqrt{\chi^2 - 4\gamma}}, a_2 \rightarrow \frac{2b_1 \sqrt{s}}{\sqrt{\chi^2 - 4\gamma}}, c \rightarrow -\frac{2\sqrt{sv}}{\sqrt{\chi^2 - 4\gamma}} \right].$$

Consequently, the explicit wave solutions of Eq. (1) are given by:

In case of $[\chi^2 - 4\gamma > 0 \ \& \ \gamma \neq 0]$,

$$\mathcal{H}_1(x, y, t) = \frac{\sqrt{s}(\chi^2 - 4\gamma + \chi \sqrt{\chi^2 - 4\gamma} \tanh(\frac{-2\sqrt{sv}t^\kappa + \sqrt{\chi^2 - 4\gamma}(\eta\kappa + \varrho x^\kappa + i\varrho y^\kappa)}{2\kappa}))}{\sqrt{\chi^2 - 4\gamma}(\chi + \sqrt{\chi^2 - 4\gamma} \tanh(\frac{-2\sqrt{sv}t^\kappa + \sqrt{\chi^2 - 4\gamma}(\eta\kappa + \varrho x^\kappa + i\varrho y^\kappa)}{2\kappa}))}, \tag{9}$$

$$\mathcal{H}_2(x, y, t) = \frac{\sqrt{s}(\chi^2 - 4\gamma + \chi \sqrt{\chi^2 - 4\gamma} \coth(\frac{-2\sqrt{sv}t^\kappa + \sqrt{\chi^2 - 4\gamma}(\eta\kappa + \varrho x^\kappa + i\varrho y^\kappa)}{2\kappa}))}{\sqrt{\chi^2 - 4\gamma}(\chi + \sqrt{\chi^2 - 4\gamma} \coth(\frac{-2\sqrt{sv}t^\kappa + \sqrt{\chi^2 - 4\gamma}(\eta\kappa + \varrho x^\kappa + i\varrho y^\kappa)}{2\kappa}))}. \tag{10}$$

In case of $[\chi^2 - 4\gamma > 0 \ \& \ \gamma = 0 \ \& \ \chi \neq 0]$,

$$\mathcal{H}_3(x, y, t) = \frac{\chi \sqrt{s} \coth\left(\frac{\chi(\eta\kappa - \frac{2\sqrt{svt^\kappa}}{\sqrt{\chi^2}} + \varrho(x^\kappa + iy^\kappa))}{2\kappa}\right)}{\sqrt{\chi^2}}. \tag{11}$$

In case of $[\chi^2 - 4\gamma = 0 \ \& \ \gamma \neq 0 \ \& \ \chi \neq 0]$,

$$\mathcal{H}_4(x, y, t) = -\frac{2\kappa \chi \sqrt{s}}{2\chi \sqrt{svt^\kappa} - \sqrt{\chi^2 - 4\gamma}(\kappa(\eta\chi + 2) + \chi \varrho(x^\kappa + iy^\kappa))}. \tag{12}$$

In case of $[\chi^2 - 4\gamma < 0 \ \& \ \gamma \neq 0]$,

$$\mathcal{H}_5(x, y, t) = \frac{\sqrt{s}(\chi^2 - 4\gamma - \chi \sqrt{4\gamma - \chi^2} \tan\left(\frac{\sqrt{4\gamma - \chi^2}(\eta\kappa - \frac{2\sqrt{svt^\kappa}}{\sqrt{\chi^2 - 4\gamma}} + \varrho x^\kappa + i\varrho y^\kappa)}{2\kappa}\right))}{\sqrt{\chi^2 - 4\gamma}(\chi - \sqrt{4\gamma - \chi^2} \tan\left(\frac{\sqrt{4\gamma - \chi^2}(\eta\kappa - \frac{2\sqrt{svt^\kappa}}{\sqrt{\chi^2 - 4\gamma}} + \varrho x^\kappa + i\varrho y^\kappa)}{2\kappa}\right))}, \tag{13}$$

$$\mathcal{H}_6(x, y, t) = \frac{\sqrt{s}(\chi^2 - 4\gamma - \chi \sqrt{4\gamma - \chi^2} \cot\left(\frac{\sqrt{4\gamma - \chi^2}(\eta\kappa - \frac{2\sqrt{svt^\kappa}}{\sqrt{\chi^2 - 4\gamma}} + \varrho x^\kappa + i\varrho y^\kappa)}{2\kappa}\right))}{\sqrt{\chi^2 - 4\gamma}(\chi - \sqrt{4\gamma - \chi^2} \cot\left(\frac{\sqrt{4\gamma - \chi^2}(\eta\kappa - \frac{2\sqrt{svt^\kappa}}{\sqrt{\chi^2 - 4\gamma}} + \varrho x^\kappa + i\varrho y^\kappa)}{2\kappa}\right))}. \tag{14}$$

Family II

$$\left[a_0 \rightarrow \frac{b_1 \chi^2 \sqrt{s}}{2\sqrt{\chi^2 - 4\gamma}}, a_1 \rightarrow \frac{2b_1 \chi \sqrt{s}}{\sqrt{\chi^2 - 4\gamma}}, a_2 \rightarrow \frac{2b_1 \sqrt{s}}{\sqrt{\chi^2 - 4\gamma}}, b_0 \rightarrow \frac{b_1 \chi}{2}, c \rightarrow -\frac{2\sqrt{sv}}{\sqrt{\chi^2 - 4\gamma}} \right].$$

Consequently, the explicit wave solutions of Eq. (1) are given by:

In case of $[\chi^2 - 4\gamma > 0 \ \& \ \gamma \neq 0]$,

$$\mathcal{H}_7(x, y, t) = \frac{\sqrt{s}(\chi^2 - 4\gamma + \chi \sqrt{\chi^2 - 4\gamma} \tanh\left(\frac{-2\sqrt{svt^\kappa} + \sqrt{\chi^2 - 4\gamma}(\eta\kappa + \varrho x^\kappa + i\varrho y^\kappa)}{2\kappa}\right))}{\sqrt{\chi^2 - 4\gamma}(\chi + \sqrt{\chi^2 - 4\gamma} \tanh\left(\frac{-2\sqrt{svt^\kappa} + \sqrt{\chi^2 - 4\gamma}(\eta\kappa + \varrho x^\kappa + i\varrho y^\kappa)}{2\kappa}\right))}, \tag{15}$$

$$\mathcal{H}_8(x, y, t) = \frac{\sqrt{s}(\chi^2 - 4\gamma + \chi \sqrt{\chi^2 - 4\gamma} \coth\left(\frac{-2\sqrt{svt^\kappa} + \sqrt{\chi^2 - 4\gamma}(\eta\kappa + \varrho x^\kappa + i\varrho y^\kappa)}{2\kappa}\right))}{\sqrt{\chi^2 - 4\gamma}(\chi + \sqrt{\chi^2 - 4\gamma} \coth\left(\frac{-2\sqrt{svt^\kappa} + \sqrt{\chi^2 - 4\gamma}(\eta\kappa + \varrho x^\kappa + i\varrho y^\kappa)}{2\kappa}\right))}. \tag{16}$$

In case of $[\chi^2 - 4\gamma > 0 \ \& \ \gamma = 0 \ \& \ \chi \neq 0]$,

$$\mathcal{H}_9(x, y, t) = \frac{\chi \sqrt{s} \coth\left(\frac{\chi(\eta\kappa - \frac{2\sqrt{svt^\kappa}}{\sqrt{\chi^2}} + \varrho(x^\kappa + iy^\kappa))}{2\kappa}\right)}{\sqrt{\chi^2}}. \tag{17}$$

In case of $[\chi^2 - 4\gamma = 0 \ \& \ \gamma \neq 0 \ \& \ \chi \neq 0]$,

$$\mathcal{H}_{10}(x, y, t) = -\frac{2\kappa \chi \sqrt{s}}{2\chi \sqrt{svt^\kappa} - \sqrt{\chi^2 - 4\gamma}(\kappa(\eta\chi + 2) + \chi \varrho(x^\kappa + iy^\kappa))}. \tag{18}$$

In case of $[\chi^2 - 4\gamma < 0 \ \& \ \gamma \neq 0]$,

$$\mathcal{H}_{11}(x, y, t) = \frac{\sqrt{s}(\chi^2 - 4\gamma - \chi\sqrt{4\gamma - \chi^2} \tan(\frac{\sqrt{4\gamma - \chi^2}(\eta\kappa - \frac{2\sqrt{sv}t^\kappa}{\sqrt{\chi^2 - 4\gamma}} + \rho x^\kappa + i\rho y^\kappa)}{2\kappa}))}{\sqrt{\chi^2 - 4\gamma}(\chi - \sqrt{4\gamma - \chi^2} \tan(\frac{\sqrt{4\gamma - \chi^2}(\eta\kappa - \frac{2\sqrt{sv}t^\kappa}{\sqrt{\chi^2 - 4\gamma}} + \rho x^\kappa + i\rho y^\kappa)}{2\kappa}))}, \tag{19}$$

$$\mathcal{H}_{12}(x, y, t) = \frac{\sqrt{s}(\chi^2 - 4\gamma - \chi\sqrt{4\gamma - \chi^2} \cot(\frac{\sqrt{4\gamma - \chi^2}(\eta\kappa - \frac{2\sqrt{sv}t^\kappa}{\sqrt{\chi^2 - 4\gamma}} + \rho x^\kappa + i\rho y^\kappa)}{2\kappa}))}{\sqrt{\chi^2 - 4\gamma}(\chi - \sqrt{4\gamma - \chi^2} \cot(\frac{\sqrt{4\gamma - \chi^2}(\eta\kappa - \frac{2\sqrt{sv}t^\kappa}{\sqrt{\chi^2 - 4\gamma}} + \rho x^\kappa + i\rho y^\kappa)}{2\kappa}))}. \tag{20}$$

2.1.2 Jacobi elliptical function method

Substituting Eq. (6) and its derivative into Eq. (3) leads to a system of equations. Solving them yields

$$[a_0 \rightarrow -\sqrt{s}, a_1 \rightarrow 0, b_1 \rightarrow 0]$$

and

$$[a_0 \rightarrow \sqrt{s}, a_1 \rightarrow 0, b_1 \rightarrow 0].$$

This result shows the failure of the Jacobi elliptical function method which is considered as a proof of the fact that there exists no unified computational method that can be used on all nonlinear evolution equation.

2.2 Fractional EW model

2.2.1 Extended $\exp(-\varphi(\vartheta))$ -expansion method

Replacing Eq. (7) into Eq. (4) and collecting all terms with the same power for $[e^{-\varepsilon\varphi(\vartheta)}, \varepsilon = 0, 1, 2, \dots]$ lead to a system of algebraic equation. Solving this system yields the following.

Family I

$$\left[\begin{aligned} a_0 &\rightarrow \frac{a_3 b_0 (-\sqrt{(\chi^2 - 4\gamma)^2} + \chi^2 + 8\gamma)}{12b_1}, \\ a_1 &\rightarrow \frac{1}{12} a_3 \left(\frac{12b_0\chi}{b_1} - \sqrt{(\chi^2 - 4\gamma)^2} + \chi^2 + 8\gamma \right), a_2 \rightarrow \frac{a_3(b_1\chi + b_0)}{b_1}, \\ r &\rightarrow \frac{1}{\sqrt{\chi^4 - 8\chi^2\gamma + 16\gamma^2}}, h \rightarrow \frac{6b_1c}{a_3\sqrt{(\chi^2 - 4\gamma)^2}}. \end{aligned} \right].$$

Consequently, the explicit wave solutions of Eq. (2) are given by:

In case of $[\chi^2 - 4\gamma > 0 \ \& \ \gamma \neq 0]$,

$$\begin{aligned} \Lambda_1(x, y, t) &= \frac{1}{12b_1(\sqrt{\chi^2 - 4\gamma} \tanh(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}) + \chi)^2} \\ &\times \left[a_3 \operatorname{sech}^2\left(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}\right) \right] \\ &\times \left(-(\sqrt{(\chi^2 - 4\gamma)^2} - \chi^2 + 4\gamma) \right) \end{aligned}$$

$$\begin{aligned}
 & \times \left(\chi \sqrt{\chi^2 - 4\gamma} \sinh\left(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right. \\
 & \left. + (\chi^2 - 2\gamma) \cosh\left(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right) \\
 & \left. - 2\gamma(\sqrt{(\chi^2 - 4\gamma)^2} + 5\chi^2 - 20\gamma) \right], \tag{21} \\
 \Lambda_2(x, y, t) = & \frac{1}{12b_1(\sqrt{\chi^2 - 4\gamma} \coth(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}) + \chi)^2} \\
 & \times \left[a_3 \operatorname{csch}^2\left(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}\right) \right. \\
 & \times \left(2\gamma(\sqrt{(\chi^2 - 4\gamma)^2} + 5\chi^2 - 20\gamma) \right. \\
 & \left. - (\sqrt{(\chi^2 - 4\gamma)^2} - \chi^2 + 4\gamma) \left(\chi \sqrt{\chi^2 - 4\gamma} \right. \right. \\
 & \left. \left. \times \sinh\left(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right. \right. \\
 & \left. \left. + (\chi^2 - 2\gamma) \cosh\left(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right) \right]. \tag{22}
 \end{aligned}$$

In case of $[\chi^2 - 4\gamma > 0 \ \& \ \gamma = 0 \ \& \ \chi \neq 0]$,

$$\Lambda_3(x, y, t) = \frac{a_3(3\chi^2 \operatorname{csch}^2(\frac{\chi(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}) - \sqrt{\chi^4 + \chi^2})}{12b_1}. \tag{23}$$

In case of $[\chi^2 - 4\gamma = 0 \ \& \ \gamma \neq 0 \ \& \ \chi \neq 0]$,

$$\Lambda_4(x, y, t) = \frac{a_3(\chi^2(\frac{12\kappa^2}{(c\chi t^\kappa + \eta\kappa\chi + 2\kappa + \chi x^\kappa)^2} - 2) - \sqrt{(\chi^2 - 4\gamma)^2} + 8\gamma)}{12b_1}. \tag{24}$$

In case of $[\chi^2 - 4\gamma = 0 \ \& \ \gamma = 0 \ \& \ \chi = 0]$,

$$\Lambda_5(x, y, t) = \frac{a_3\kappa^2}{b_1(ct^\kappa + \eta\kappa + x^\kappa)^2}. \tag{25}$$

In case of $[\chi^2 - 4\gamma < 0 \ \& \ \gamma \neq 0]$,

$$\begin{aligned}
 \Lambda_6(x, y, t) = & \frac{1}{12b_1(\chi - \sqrt{4\gamma - \chi^2} \tan(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}))^2} \\
 & \times \left[a_3 \sec^2\left(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}\right) \right. \\
 & \times \left(-(\sqrt{(\chi^2 - 4\gamma)^2} - \chi^2 + 4\gamma) \right. \\
 & \left. \times \left((\chi^2 - 2\gamma) \cos\left(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right) \right]
 \end{aligned}$$

$$\begin{aligned}
 & -\chi\sqrt{4\gamma - \chi^2} \sin\left(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \\
 & - 2\gamma(\sqrt{(\chi^2 - 4\gamma)^2 + 5\chi^2 - 20\gamma}) \Big], \tag{26}
 \end{aligned}$$

$$\begin{aligned}
 \Lambda_7(x, y, t) = & \frac{1}{12b_1(\chi - \sqrt{4\gamma - \chi^2} \cot(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}))^2} \\
 & \times \left[a_3 \csc^2\left(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}\right) \right. \\
 & \times \left((\sqrt{(\chi^2 - 4\gamma)^2 - \chi^2 + 4\gamma} \right. \\
 & \times \left(\chi\sqrt{4\gamma - \chi^2} \sin\left(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right. \\
 & \left. \left. + (\chi^2 - 2\gamma) \cos\left(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right) \right. \\
 & \left. - 2\gamma(\sqrt{(\chi^2 - 4\gamma)^2 + 5\chi^2 - 20\gamma}) \right]. \tag{27}
 \end{aligned}$$

Family II

$$\begin{aligned}
 & \left[b_1 \rightarrow \frac{2b_0}{\chi}, a_0 \rightarrow \frac{1}{24}a_3\chi(\sqrt{(\chi^2 - 4\gamma)^2 + \chi^2 + 8\gamma}), \right. \\
 & a_1 \rightarrow \frac{1}{12}a_3(\sqrt{(\chi^2 - 4\gamma)^2 + 7\chi^2 + 8\gamma}), \\
 & \left. a_2 \rightarrow \frac{3a_3\chi}{2}, r \rightarrow -\frac{1}{\sqrt{\chi^4 - 8\chi^2\gamma + 16\gamma^2}}, h \rightarrow -\frac{12b_0c}{a_3\chi\sqrt{(\chi^2 - 4\gamma)^2}} \right].
 \end{aligned}$$

Consequently, the explicit wave solutions of Eq. (2) are given by:

In case of $[\chi^2 - 4\gamma > 0 \ \& \ \gamma \neq 0]$,

$$\begin{aligned}
 \Lambda_8(x, y, t) = & \frac{1}{24b_0(\sqrt{\chi^2 - 4\gamma} \tanh(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}) + \chi)^2} \\
 & \times \left[a_3\chi \operatorname{sech}^2\left(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}\right) \right. \\
 & \times \left((\sqrt{(\chi^2 - 4\gamma)^2 + \chi^2 - 4\gamma} \right. \\
 & \times \left(\chi\sqrt{\chi^2 - 4\gamma} \sinh\left(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right. \\
 & \left. \left. + (\chi^2 - 2\gamma) \cosh\left(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right) \right. \\
 & \left. + 2\gamma(\sqrt{(\chi^2 - 4\gamma)^2 - 5\chi^2 + 20\gamma}) \right], \tag{28}
 \end{aligned}$$

$$\begin{aligned} \Lambda_9(x, y, t) = & \frac{1}{24b_0(\sqrt{\chi^2 - 4\gamma} \coth(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}) + \chi)^2} \\ & \times \left[a_3\chi \operatorname{csch}^2\left(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}\right) \right. \\ & \times \left((\sqrt{(\chi^2 - 4\gamma)^2} + \chi^2 - 4\gamma) \right. \\ & \times \left(\chi\sqrt{\chi^2 - 4\gamma} \sinh\left(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right. \\ & \left. \left. \left. + (\chi^2 - 2\gamma) \cosh\left(\frac{\sqrt{\chi^2 - 4\gamma}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right) \right) \right. \\ & \left. \left. - 2\gamma(\sqrt{(\chi^2 - 4\gamma)^2} - 5\chi^2 + 20\gamma) \right) \right]. \end{aligned} \tag{29}$$

In case of $[\chi^2 - 4\gamma > 0 \ \& \ \gamma = 0 \ \& \ \chi \neq 0]$,

$$\Lambda_{10}(x, y, t) = \frac{a_3\chi(3\chi^2 \operatorname{csch}^2(\frac{\chi(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}) + \sqrt{\chi^4 + \chi^2})}{24b_0}. \tag{30}$$

In case of $[\chi^2 - 4\gamma = 0 \ \& \ \gamma \neq 0 \ \& \ \chi \neq 0]$,

$$\Lambda_{11}(x, y, t) = \frac{a_3\chi(\chi^2(\frac{12\kappa^2}{(c\chi t^\kappa + \eta\kappa\chi + 2\kappa + \chi x^\kappa)^2} - 2) + \sqrt{(\chi^2 - 4\gamma)^2} + 8\gamma)}{24b_0}. \tag{31}$$

In case of $[\chi^2 - 4\gamma < 0 \ \& \ \gamma \neq 0]$,

$$\begin{aligned} \Lambda_{12}(x, y, t) = & \frac{1}{24b_0(\chi - \sqrt{4\gamma - \chi^2} \tan(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}))^2} \\ & \times \left[a_3\chi \operatorname{sec}^2\left(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}\right) \right. \\ & \times \left((\sqrt{(\chi^2 - 4\gamma)^2} + \chi^2 - 4\gamma) \right. \\ & \times \left((\chi^2 - 2\gamma) \cos\left(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right. \\ & \left. \left. - \chi\sqrt{4\gamma - \chi^2} \sin\left(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right) \right. \\ & \left. \left. \times \gamma(\sqrt{(\chi^2 - 4\gamma)^2} - 5\chi^2 + 20\gamma) \right) \right], \end{aligned} \tag{32}$$

$$\begin{aligned} \Lambda_{13}(x, y, t) = & \frac{-1}{24b_0(\chi - \sqrt{4\gamma - \chi^2} \cot(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}))^2} \\ & \times \left[a_3\chi \operatorname{csc}^2\left(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{2\kappa}\right) \right. \\ & \times \left((\sqrt{(\chi^2 - 4\gamma)^2} + \chi^2 - 4\gamma) \right. \end{aligned}$$

Table 1 Some solutions of auxiliary equation of Eq. 8

ρ	p	q	$\phi(\vartheta)$
1	$-(1 + m^2)$	m^2	$sn(\vartheta)$
$1 - m^2$	$2m^2 - 1$	$-m^2$	$cn(\vartheta)$
$m^2 - 1$	$2 - m^2$	-1	$dn(\vartheta)$
$\frac{1}{4}$	$\frac{1-2m^2}{2}$	$\frac{1}{4}$	$ns(\vartheta) \pm cs(\vartheta)$ or $\frac{sn(\vartheta)}{1 \pm cn(\vartheta)}$
$\frac{1-m^2}{4}$	$\frac{1+m^2}{2}$	$\frac{1-m^2}{4}$	$nc(\vartheta) \pm sc(\vartheta)$ or $\frac{cn(\vartheta)}{1 \pm sn(\vartheta)}$
1	$2 - m^2$	$1 - m^2$	$sc(\vartheta)$
$1 - m^2$	$2 - m^2$	1	$cs(\vartheta)$

$$\begin{aligned}
 & \times \left(\chi \sqrt{4\gamma - \chi^2} \sin\left(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right. \\
 & \left. + (\chi^2 - 2\gamma) \cos\left(\frac{\sqrt{4\gamma - \chi^2}(ct^\kappa + \eta\kappa + x^\kappa)}{\kappa}\right) \right) \\
 & \left. - 2\gamma (\sqrt{(\chi^2 - 4\gamma)^2 - 5\chi^2 + 20\gamma}) \right]. \tag{33}
 \end{aligned}$$

2.2.2 Jacobi elliptical function method

Substituting Eq. (8) and its derivative into Eq. (4) leads to a system of equations. Solving them yields

$$\begin{aligned}
 & \left[a_0 \rightarrow \frac{1}{2} \left(\frac{cp}{h\sqrt{p^2 - 3q\rho}} - \frac{c}{h} \right), a_1 \rightarrow 0, a_2 \rightarrow \frac{6cq}{h\sqrt{16p^2 - 48q\rho}}, \right. \\
 & \left. b_1 \rightarrow 0, b_2 \rightarrow 0, r \rightarrow \frac{1}{\sqrt{16p^2 - 48q\rho}} \right].
 \end{aligned}$$

Using these values and Table 1 leads to formulating the explicit wave solutions of Eq. (2) in the following formats:

$$\Lambda_{14}(x, y, t) |_{\rho \rightarrow 1, p \rightarrow -2, q \rightarrow 1} = -\frac{3c \operatorname{sech}^2\left(\frac{ct^\kappa + x^\kappa}{\kappa}\right)}{2h}, \tag{34}$$

$$\Lambda_{15}(x, y, t) |_{\rho \rightarrow \frac{1}{4}, p \rightarrow -\frac{1}{2}, q \rightarrow \frac{1}{4}} = \frac{3c((\coth(\frac{ct^\kappa + x^\kappa}{\kappa}) \pm \operatorname{csch}(\frac{ct^\kappa + x^\kappa}{\kappa}))^2 - 1)}{2h}, \tag{35}$$


$$\Lambda_{16}(x, y, t) |_{\rho \rightarrow 0, p \rightarrow 1, q \rightarrow 1} = \frac{3c \operatorname{csch}^2\left(\frac{ct^\kappa + x^\kappa}{\kappa}\right)}{2h}, \tag{36}$$

$$\Lambda_{16}(x, y, t) |_{\rho \rightarrow \frac{1}{4}, p \rightarrow \frac{1}{2}, q \rightarrow \frac{1}{4}} = \frac{c(3(\operatorname{csc}(\frac{ct^\kappa + x^\kappa}{\kappa}) \pm \cot(\frac{ct^\kappa + x^\kappa}{\kappa}))^2 + 1)}{2h}, \tag{37}$$

$$\Lambda_{18}(x, y, t) |_{\rho \rightarrow \frac{1}{4}, p \rightarrow \frac{1}{2}, q \rightarrow \frac{1}{4}} = \frac{c(3(\sec(\frac{ct^\kappa + x^\kappa}{\kappa}) \pm \tan(\frac{ct^\kappa + x^\kappa}{\kappa}))^2 + 1)}{2h}, \tag{38}$$

$$\Lambda_{19}(x, y, t) |_{\rho \rightarrow 1, p \rightarrow 2, q \rightarrow 1} = \frac{c(3(\sec(\frac{ct^\kappa + x^\kappa}{\kappa}) \pm \tan(\frac{ct^\kappa + x^\kappa}{\kappa}))^2 + 1)}{2h}, \tag{39}$$

$$\Lambda_{20}(x, y, t) |_{\rho \rightarrow 1, p \rightarrow 2, q \rightarrow 1} = \frac{c(3 \cot^2\left(\frac{ct^\kappa + x^\kappa}{\kappa}\right) + 1)}{2h}. \tag{40}$$




3 Figures representation

This section explains the shown figures in our paper. All these plots depend on individual values of the indicated parameters in the obtained solution. Now, we discuss the representation of the shown figures with their settings as follows:

- Fig. 1 represents a cone wave solution of Eq. (9) when $\eta = 1$, $\kappa = 0.5$, $\lambda = 3$, $\mu = 2$, $s = 4$, $\nu = 6$, $\gamma = 7$, $\varrho = 5$ in a three-dimensional sketch.
- Fig. 2 shows a contour plot of Eq. (9) when $\eta = 1$, $\kappa = 0.5$, $\lambda = 3$, $\mu = 0$, $s = 4$, $\nu = 6$, $\gamma = 7$, $\varrho = 5$ that explains the surface is symmetric and peaks in the center.
- Fig. 3 represents a dark wave solution of Eq. (21) when $(a_3 = 3)$; $b_1 = 2$, $c = 5$, $\eta = 1$, $\kappa = 0.5$, $\lambda = 3$, $\mu = 2$, $s = 4$ in three distinct types of plots.
- Fig. 4 shows dark wave solution of Eq. (23) when $a_3 = 3$, $b_1 = 2$, $c = 5$, $\eta = 1$, $\kappa = 0.5$, $\lambda = 3$, $\mu = 2$, $s = 4$ in three different plots.
- Fig. 5 shows dark wave solution of Eq. (34) when $a_3 = 3$, $b_1 = 2$, $c = 5$, $\eta = 1$, $\kappa = 0.5$, $\lambda = 3$, $\mu = 0$, $s = 4$ in three various kind of plots.
- Fig. 6 shows bright wave solution of Eq. (35) when $c = 3$, $h = 2$, $\kappa = 0.5$ in three different forms of sketches.

4 Results and discussion


This segment displays our approaches and their latest functions. Check out how close we are and how the ideas we have found can be contrasted with the previously published papers. The key elements of our debate are the methodological approach employed and the approaches produced.



1. *The computational schemes used:*

We used two computational methods (the extended $\exp(-\varphi(\vartheta))$ -expansion method and the Jacobi elliptical function method) for the fractional BP model and the EW model for constructing the exact traveling and solitary wave solutions. A compliant fractional operator was employed to transform fractional aspects of the equation to a nonlinear ordinary differential equation. This fractional operator enables the classification schemes to be extended to the transformed shape. All systems have the desired approach.

$$\begin{aligned}
 \mathcal{H}(\vartheta) &= \Lambda(\vartheta) \\
 &= \begin{cases} \frac{\sum_{i=0}^m a_i e^{-i\phi(\vartheta)}}{\sum_{i=0}^n b_i e^{-i\phi(\vartheta)}}, \\ \phi'(\vartheta) \rightarrow \lambda + \mu e^{\phi(\vartheta)} + \frac{1}{e^{\phi(\vartheta)}}, \\ \sum_{i=1}^n a_i \phi(\vartheta)^i + \sum_{i=1}^n b_i \phi(\vartheta)^{-i} + a_0, \\ \phi'(\vartheta) \rightarrow \sqrt{p\phi(\vartheta)^2 + q\phi(\vartheta)^4 + \rho}, \end{cases} \quad (41)
 \end{aligned}$$




where $[n, m]$ are arbitrary constants to be determined by using the homogeneous balance rule, while $[\lambda, \mu, p, q, \rho]$ are arbitrary constants to be determined by solving the obtained algebraic system of equations. Moreover, applying two different schemes on one model shows its ability to be used for other various schemes where there is no unified method that is applied for all nonlinear evolution equations, and that is well demonstrated in our paper.

2. *The obtained solutions:*

This part gives a comparison between our obtained solutions and those obtained in previously published papers. In [64], Mostafa M. A. Khater, Raghda A. M. Attia, and Dianchen Lu applied the modified Khater method to a fractional biological population model, fractional equal width model, and fractional modified equal width equation. They got many distinct types of solutions for these fractional biological models.

- Eq. (27, [64]) is equal to Eq. (17) when $[\chi = 2, \lambda = \nu, n\kappa = 0, \varrho = \mu\sqrt{-\alpha\sigma}]$.
- All other obtained solutions of the fractional BP model are different from those obtained in [64].
- All our obtained solutions of the fractional EW model are new and different from those obtained in [64].



5 Conclusion

This paper examined two nonlinear BP and EW fractional models via $\exp(-\varphi(\vartheta))$ -expansion and the Jacobi elliptical function method. The conformable fractional derivative has been employed to convert the nonlinear partial differential equations to an ordinary differential equation with an integer order. Many distinct wave solutions have been obtained and have been represented in three-, two-dimensional, and contour plots. These solutions were explained by different illustrations which clarify the new features of the fractional models in question. Our solutions have been explained in terms of precision and innovation. The novelty of our solutions is shown by comparing our solution with that obtained in previous research papers. The powerful and effective application of the used method is examined and tested to show its ability to be applied to other nonlinear evolution equations.

Appendix

Given function $f : [0, \infty) \rightarrow R$. Then the (conformable fractional derivative) of f of order α is defined by

$$T_\alpha(f)(t) = \lim_{\varepsilon \rightarrow 0} \frac{f(t + \varepsilon t^{1-\alpha}) - f(t)}{\varepsilon}. \tag{42}$$




Figure 5 Dark wave solution of (34) in three (a), two-dimensional (b), and contour plots (c)

For all $t > 0$, $\alpha \in (0, 1)$ if f is α -differentiable in some $(0, \alpha)$, $\alpha > 0$ and $\lim_{t \rightarrow 0} f^\alpha(t)$ exists, then define $f^\alpha(0) = \lim_{t \rightarrow 0} f^\alpha(t)$.

We sometimes write $f^\alpha(t)$ for $T_\alpha(f)(t)$ to denote the conformable fractional derivatives of f of order α . In addition, if the conformable fractional derivative of f of order α exists, then we say f is α -differentiable. We should take into consideration that $T_\alpha(t^p) = pt^{p-\alpha}$. Further, this definition coincides with the same of traditional definition of Riemann–Liouville and of Caputo on polynomials (up to a constant multiple).

The conformable fractional properties:

- $T_\alpha(e^{ct}) = ct^{1-\alpha}e^{ct}$.
- $T_\alpha(\sin(at)) = at^{1-\alpha}\cos(at)$.
- $T_\alpha \sin(at) = -at^{1-\alpha}\sin(at)$.
- $T_\alpha(\tan(at)) = at^{1-\alpha}\sec^2(at)$.
- $T_\alpha(\cot(at)) = -at^{1-\alpha}\csc^2(at)$.
- $T_\alpha(\sec(at)) = at^{1-\alpha}\sec(at)\tan(at)$.
- $T_\alpha(\csc(at)) = -at^{1-\alpha}\csc(at)\cot(at)$.
- $T_\alpha\left(\frac{t^\alpha}{\alpha}\right) = 1$.




Figure 6 Bright wave solution of (35) in three (a), two-dimensional (b), and contour plots (c)

Acknowledgements

This project was funded by the Deanship of Scientific Research (DSR), King Abdulaziz University, Jeddah, under grant No. (D-671-305-1441). The authors, therefore, gratefully acknowledge DSR technical and financial support.

Funding

This project was funded by the Deanship of Scientific Research (DSR), King Abdulaziz University, Jeddah, under grant No. (D-671-305-1441). The authors, therefore, gratefully acknowledge DSR technical and financial support.

Availability of data and materials

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors conceived of the study, participated in its design and coordination, drafted the manuscript, participated in the sequence alignment, and read and approved the final manuscript.

Author details

¹Department of Physics, College of Sciences, University of Bisha, P.O. Box 344, Bisha 61922, Saudi Arabia. ²Physics Department, Faculty of Science, Al-Azhar University, Assiut 71524, Egypt. ³Department of Mathematics, Faculty of Science, Jiangsu University, 212013 Zhenjiang, China. ⁴Department of Mathematics, Obour Institutes, 11828 Cairo, Egypt. ⁵Department of Mathematics, Cankaya University, Ankara, Turkey. ⁶Institute of Space Sciences, Magurele-Bucharest, Romania. ⁷Department of Medical Research, China Medical University Hospital, China Medical University, Taichung, Taiwan. ⁸Department of Mathematics, Faculty of Science, Taif University, Taif 888, Saudi Arabia. ⁹Department of Mathematics, Faculty of Science, South Valley University, Qena 83523, Egypt. ¹⁰Department of Engineering Physics and Instrumentation, National Institute of Applied Sciences and Technology, Carthage University, Tunis, Tunisia. ¹¹Department of Physics, Faculty of Science, Taif University, Taif 888, Saudi Arabia. ¹²Deanship of Scientific Research, King Abdulaziz University, Jeddah, Saudi Arabia.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 18 July 2020 Accepted: 24 September 2020 Published online: 06 October 2020

References

1. Khater, M.M.A., Attia, R.A.M., Abdel-Aty, A.-H., Abdel-Khalek, S., Al-Hadeethi, Y., Lu, D.: On the computational and numerical solutions of the transmission of nerve impulses of an excitable system (the neuron system). *J. Intell. Fuzzy Syst.* **38**, 2603–2610 (2020)
2. Abdalla, M.S., Abdel-Aty, M., Obada, A.-S.F.: Degree of entanglement for anisotropic coupled oscillators interacting with a single atom. *J. Opt. B, Quantum Semiclass. Opt.* **4**, 396–401 (2002)
3. Yang, X.-J., Tenreiro Machado, J.: A new fractal nonlinear Burgers' equation arising in the acoustic signals propagation. *Math. Methods Appl. Sci.* **42**(18), 7539–7544 (2019)
4. Janaki, M., Kanagarajan, K., Elsayed, E.M.: A note on nonlinear implicit neutral Katugampola fractional differential equations with impulse effects and finite delay. *Sohag J. Math.* **6**(2), 29–39 (2019)
5. Alharbi, S.A., Rambely, A.S.: Dynamic behaviour and stabilisation to boost the immune system by complex interaction between tumour cells and vitamins intervention. *Adv. Differ. Equ.* **2020**(1), 412 (2020)
6. Yang, X.-J.: *General Fractional Derivatives: Theory, Methods and Applications*, vol. 1. CRC Press, Boca Raton (2019)
7. Yang, X.-J., Feng, Y.-Y., Cattani, C., Inc, M.: Fundamental solutions of anomalous diffusion equations with the decay exponential kernel. *Math. Methods Appl. Sci.* **42**(11), 4054–4060 (2019)
8. Rizvi, S.R., Afzal, I., Ali, K., Younis, M.: Stationary Solutions for Nonlinear Schrödinger Equations by Lie Group Analysis. *Acta Phys. Pol. A* **136**, 187–189 (2019)
9. Khater, M.M.A., Attia, R.A.M., Abdel-Aty, A.: Computational analysis of a nonlinear fractional emerging telecommunication model with higher-order dispersive cubic–quintic. *Inf. Sci. Lett.* **9**, 83–93 (2020)
10. Arif, A., Younis, M., Imran, M., Tantawy, M., Rizvi, S.T.R.: Solitons and lump wave solutions to the graphene thermophoretic motion system with a variable heat transmission. *Eur. Phys. J. Plus* **134**(6), 303 (2019)
11. Liu, J.-G., Yang, X.-J., Feng, Y.-Y.: On integrability of the time fractional nonlinear heat conduction equation. *J. Geom. Phys.* **144**, 190–198 (2019)
12. Wu, B., Sun, C.: Multiple positive solutions for a continuous fractional boundary value problem with fractional q -differences. *Sohag J. Math.* **7**(2), 43–48 (2020)
13. Liang, Y., Yang, H., Li, H.: Existence of positive solutions for the fractional q -difference boundary value problem. *Adv. Differ. Equ.* **2020**(1), 416 (2020)
14. Rizvi, S.T.R., Afzal, I., Ali, K.: Chirped optical solitons for Triki–Biswas equation. *Mod. Phys. Lett. B* **33**(22), 1950264 (2019)
15. Liu, J.-G., Yang, X.-J., Feng, Y.-Y., Cui, P.: On group analysis of the time fractional extended $(2 + 1)$ -dimensional Zakharov–Kuznetsov equation in quantum magneto-plasmas. *Math. Comput. Simul.* **178**, 407–421 (2020)
16. Alshammari, M., Al-Smadi, M., Alshammari, S., Abu Arqub, O., Hashim, I., Alias, M.: An attractive analytic-numeric approach for the solutions of uncertain Riccati differential equations using residual power series. *Appl. Math. Inf. Sci.* **14**, 177–190 (2020)
17. Owyed, S., Abdou, M., Abdel-Aty, A.-H., Alharbi, W., Nekhili, R.: Numerical and approximate solutions for coupled time fractional nonlinear evolutions equations via reduced differential transform method. *Chaos Solitons Fractals* **131**, 109474 (2020)
18. Jiang, Y., Ge, Y.: An explicit fourth-order compact difference scheme for solving the 2D wave equation. *Adv. Differ. Equ.* **2020**(1), 415 (2020)
19. Lu, D., Seadawy, A.R., Khater, M.M.: Structures of exact and solitary optical solutions for the higher-order nonlinear Schrödinger equation and its applications in mono-mode optical fibers. *Mod. Phys. Lett. B* **33**(23), 1950279 (2019)
20. Golmankhaneh, A.K., Tunc, C., Nia, S.M., Golmankhaneh, A.K.: A review on local and non-local fractal calculus. *Numer. Comput. Methods Sci. Eng.* **1**, 19–31 (2019)
21. Khater, M.M., Lu, D., Attia, R.A.: Dispersive long wave of nonlinear fractional Wu–Zhang system via a modified auxiliary equation method. *AIP Adv.* **9**(2), 025003 (2019)
22. Khater, M.M., Lu, D., Attia, R.A.: Erratum: “Dispersive long wave of nonlinear fractional Wu–Zhang system via a modified auxiliary equation method” [AIP Adv. 9, 025003 (2019)]. *AIP Adv.* **9**(4), 049902 (2019)
23. Khater, M.M., Lu, D., Attia, R.A.: Lump soliton wave solutions for the $(2 + 1)$ -dimensional Konopelchenko–Dubrovsky equation and KdV equation. *Mod. Phys. Lett. B* **2019**, 1950199 (2019)
24. Panda, S.K., Abdeljawad, T., Ravichandran, C.: Novel fixed point approach to Atangana–Baleanu fractional and LP-Fredholm integral equations. *Alex. Eng. J.* **59**(4), 1959–1970 (2020)
25. Yang, S., Deng, M., Ren, R.: Stochastic resonance of fractional-order Langevin equation driven by periodic modulated noise with mass fluctuation. *Adv. Differ. Equ.* **2020**(1), 1 (2020)
26. Liu, J.-G., Yang, X.-J., Feng, Y.-Y.: On integrability of the extended $(3 + 1)$ -dimensional Jimbo–Miwa equation. *Math. Methods Appl. Sci.* **43**(4), 1646–1659 (2020)
27. Al-Refai, M., Pal, K.: New aspects of Caputo–Fabrizio fractional derivative. *Prog. Fract. Differ. Appl.* **5**(2), 157–166 (2019)
28. Chatzarakis, G., Deepa, M., Nagajothi, N., Sadhasivam, V.: Oscillatory properties of a certain class of mixed fractional differential equations. *Appl. Math. Inf. Sci.* **14**, 123–131 (2020)
29. Khater, M.M., Attia, R.A., Abdel-Aty, A.-H., Abdou, M., Eleuch, H., Lu, D.: Analytical and semi-analytical ample solutions of the higher-order nonlinear Schrödinger equation with the non-Kerr nonlinear term. *Results Phys.* **16**, 103000 (2020)
30. Park, C., Khater, M.M.A., Abdel-Aty, A.-H., Attia, R.A.M., Rezazadeh, H., Zidan, A.M., Mohamed, A.-B.A.: Dynamical analysis of the nonlinear complex fractional emerging telecommunication model with higher-order dispersive cubic–quintic. *Alex. Eng. J.* **59**(3), 1425–1433 (2020)
31. Cattani, C., Rushchitsky, J., Sinchilo, S.: Physical constants for one type of nonlinearly elastic fibrous micro-and nanocomposites with hard and soft nonlinearities. *Int. Appl. Mech.* **41**(12), 1368–1377 (2005)
32. Singh, J., Kumar, D., Hammouch, Z., Atangana, A.: A fractional epidemiological model for computer viruses pertaining to a new fractional derivative. *Appl. Math. Comput.* **316**, 504–515 (2018)
33. Cattani, C., Pierro, G.: On the fractal geometry of DNA by the binary image analysis. *Bull. Math. Biol.* **75**(9), 1544–1570 (2013)

34. Zhang, J.-G.: The Fourier–Yang integral transform for solving the 1-D heat diffusion equation. *Therm. Sci.* **21**(1), 63–69 (2017)
35. Liu, J.-G., Yang, X.-J., Feng, Y.-Y., Zhang, H.-Y.: Analysis of the time fractional nonlinear diffusion equation from diffusion process. *J. Appl. Anal. Comput.* **10**, 1060–1072 (2020)
36. Hasanov, A., Choi, J.: Note on Euler–Bernoulli equation. *Sohag J. Math.* **7**(2), 33–36 (2020)
37. Babaei, A., Jafari, H., Liya, A.: Mathematical models of HIV/AIDS and drug addiction in prisons. *Eur. Phys. J. Plus* **135**(5), 395 (2020)
38. Wang, S., Li, Y., Shao, Y., Cattani, C., Zhang, Y., Du, S.: Detection of dendritic spines using wavelet packet entropy and fuzzy support vector machine. *CNS Neurol. Disord. Drug Targets* **16**(2), 116–121 (2017)
39. Widyan, A.M.: Chance constrained approach for treating multicriterion inventory model with random variables in the constraints. *J. Stat. Appl. Pro.* **9**(2), 207–213 (2020)
40. Kanth, A.R., Garg, N.: Computational simulations for solving a class of fractional models via Caputo–Fabrizio fractional derivative. *Proc. Comput. Sci.* **125**, 476–482 (2018)
41. Du, Q., Hesthaven, J.S., Li, C., Shu, C.-W., Tang, T.: Preface to the Focused Issue on Fractional Derivatives and General Nonlocal Models. *Commun. Appl. Math. Comput. Sci.* **1**(4), 503–504 (2019)
42. Abdelhakem, M., Ahmed, A., El-kady, M.: Spectral monic Chebyshev approximation for higher order differential equations. *Math. Sci. Lett.* **8**, 11–17 (2019)
43. Giusti, A.: A comment on some new definitions of fractional derivative. *Nonlinear Dyn.* **93**(3), 1757–1763 (2018)
44. Al-Saif, A.S.J., Abdul-Wahab, M.S.: Application of new simulation scheme for the nonlinear biological population model. *Numer. Comput. Methods Sci. Eng.* **1**, 89–99 (2019)
45. Abdeljawad, T., Al-Mdallal, Q.M., Jarad, F.: Fractional logistic models in the frame of fractional operators generated by conformable derivatives. *Chaos Solitons Fractals* **119**, 94–101 (2019)
46. Martinez, L., Rosales, J., Carreño, C., Lozano, J.: Electrical circuits described by fractional conformable derivative. *Int. J. Circuit Theory Appl.* **46**(5), 1091–1100 (2018)
47. Nchama, G.A.M., Mecias, A.L., Richard, M.R.: The Caputo–Fabrizio fractional integral to generate some new inequalities. *Inf. Sci. Lett.* **8**, 73–80 (2019)
48. Shakeel, M., Iqbal, M.A., Mohyud-Din, S.T.: Closed form solutions for nonlinear biological population model. *J. Biol. Syst.* **26**(01), 207–223 (2018)
49. Bushnaq, S., Ali, S., Shah, K., Arif, M.: Exact solution to non-linear biological population model with fractional order. *Therm. Sci.* **22**(1), 317–327 (2018)
50. Xu, Y., Lu, D.: Study on Approximate Solution of Fractional Order Biological Population Model. *Arch. Curr. Res. Int.* **15**, 1–11 (2018)
51. Wu, C., Rui, W.: Method of separation variables combined with homogenous balanced principle for searching exact solutions of nonlinear time-fractional biological population model. *Commun. Nonlinear Sci. Numer. Simul.* **63**, 88–100 (2018)
52. Goswami, A., Singh, J., Kumar, D., et al.: An efficient analytical approach for fractional equal width equations describing hydro–magnetic waves in cold plasma. *Phys. A, Stat. Mech. Appl.* **524**, 563–575 (2019)
53. Zafar, A.: Rational exponential solutions of conformable space–time fractional equal-width equations. *Nonlinear Eng.* **8**(1), 350–355 (2019)
54. Hassan, S., Abdelrahman, M.A.: Solitary wave solutions for some nonlinear time–fractional partial differential equations. *Pramana* **91**(5), 67 (2018)
55. Ray, S.S.: Invariant analysis and conservation laws for the time fractional $(2 + 1)$ -dimensional Zakharov–Kuznetsov modified equal width equation using Lie group analysis. *Comput. Math. Appl.* **76**(9), 2110–2118 (2018)
56. Baskonus, H.M., Bulut, H., Atangana, A.: On the complex and hyperbolic structures of the longitudinal wave equation in a magneto–electro–elastic circular rod. *Smart Mater. Struct.* **25**(3), 035022 (2016)
57. Koçak, Z.F., Bulut, H., Koc, D.A., Baskonus, H.M.: Prototype traveling wave solutions of new coupled Konno–Oono equation. *Optik* **127**(22), 10786–10794 (2016)
58. Baskonus, H.M., Bulut, H., Belgacem, F.B.M.: Analytical solutions for nonlinear long–short wave interaction systems with highly complex structure. *J. Comput. Appl. Math.* **312**, 257–266 (2017)
59. Hafez, M., Akbar, M.: An exponential expansion method and its application to the strain wave equation in microstructured solids. *Ain Shams Eng. J.* **6**(2), 683–690 (2015)
60. Ayatollahi, M., Safavi-Naeini, S.: A fast analysis method based on exponential expansion of Green’s function for large multilayer structures. In: *IEEE Antennas and Propagation Society International Symposium. 2001 Digest. Held in Conjunction with: USNC/URSI National Radio Science Meeting (Cat. No. 01CH37229)*, vol. 4, pp. 850–853 (2001)
61. Tala-Tebue, E., Djoufack, Z., Fendzi-Donfack, E., Kenfack-Jiotsa, A., Kofané, T.: Exact solutions of the unstable nonlinear Schrödinger equation with the new Jacobi elliptic function rational expansion method and the exponential rational function method. *Optik* **127**(23), 11124–11130 (2016)
62. Tasbozan, O., Çenesiz, Y., Kurt, A.: New solutions for conformable fractional Boussinesq and combined KdV–mKdV equations using Jacobi elliptic function expansion method. *Eur. Phys. J. Plus* **131**(7), 244 (2016)
63. Feng, Q., Meng, F.: Traveling wave solutions for fractional partial differential equations arising in mathematical physics by an improved fractional Jacobi elliptic equation method. *Math. Methods Appl. Sci.* **40**(10), 3676–3686 (2017)
64. Khater, M., Attia, R., Lu, D.: Modified Auxiliary Equation Method versus Three Nonlinear Fractional Biological Models in Present Explicit Wave Solutions. *Math. Comput. Appl.* **24**(1), 1 (2019)

Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com