



Research article

Analysis of positivity results for discrete fractional operators by means of exponential kernels

Pshtiwan Othman Mohammed^{1,*}, Donal O'Regan², Aram Bahroz Brzo³, Khadijah M. Abualnaja⁴ and Dumitru Baleanu^{5,6,7,*}

¹ Department of Mathematics, College of Education, University of Sulaimani, Sulaimani 46001, Iraq

² School of Mathematical and Statistical Sciences, National University of Ireland, Galway, Ireland

³ Department of Physics, College of Education, University of Sulaimani, Sulaimani 46001, Kurdistan Region, Iraq

⁴ Department of Mathematics and Statistics, College of Science, Taif University, P.O. Box 11099, Taif 21944, Saudi Arabia

⁵ Department of Mathematics, Cankaya University, 06530 Balgat, Ankara, Turkey

⁶ Institute of Space Sciences, R76900 Magurele-Bucharest, Romania

⁷ Department of Mathematics, King Abdul Aziz University, Jeddah 21577, Saudi Arabia

* **Correspondence:** Email: pshtiwansangawi@gmail.com, dumitru@cankaya.edu.tr.

Abstract: In this study, we consider positivity and other related concepts such as α -convexity and α -monotonicity for discrete fractional operators with exponential kernel. Namely, we consider discrete Δ fractional operators in the Caputo sense and we apply efficient initial conditions to obtain our conclusions. Note positivity results are an important factor for obtaining the composite of double discrete fractional operators having different orders.

Keywords: discrete fractional calculus; discrete CF-fractional operators; monotonicity analyses

2020 Mathematics Subject Classification: 26A48, 26A51, 33B10, 39A12, 39B62

1. Introduction

Recent years have witnessed the development of distributed discrete fractional operators based on singular and nonsingular kernels with the aim of solving a large variety of discrete problems arising in different application fields such as biology, physics, robotics, economic sciences and engineering (see for example [1–9]). These operators depend on their corresponding kernels overcoming some limits of the order of discrete operators, for example the most popular operators are Riemann-Liouville

and Caputo with standard kernels, Caputo-Fabrizio with exponential kernels, Attangana-Baleanu with Mittag-Leffler kernels (see for example [10–13]). We also refer the reader to [14–18] for discrete fractional operators. Modeling and positivity simulations have been developed or adapted for discrete fractional operators, ranging from continuous fractional models to discrete fractional frameworks; see for example [1, 19, 20]). For other results on positivity and monotonicity we refer the reader to [21–25] and for discrete fractional models with monotonicity and positivity which is important in the context of discrete fractional calculus we refer the reader to [26–29].

In this work, we are interested in finding positivity and monotonicity results for the following single and composition of delta fractional difference equations:

$$\left({}^{CFC} \Delta^{\nu} \mathbf{G}\right)(t)$$

and

$$\left({}^{CFC} \Delta^{\nu} {}^{CFC} \Delta^{\mu} \mathbf{G}\right)(t),$$

where we will assume that \mathbf{G} is defined on $\mathbb{N}_a := \{a, a + 1, \dots\}$, and ν and μ are two different positive orders.

The paper is structured as follows. The mathematical backgrounds and preliminaries needed are given in Section 2. Section 3 presents the problem statement and the main results. Conclusions are provided in Section 4.

2. Mathematical backgrounds and preliminaries

Let us start this section by recalling the notions of discrete delta Caputo-Fabrizio fractional operators that we will need.

Definition 2.1 (see [30, 31]). Let $(\Delta \mathbf{G})(t) = \mathbf{G}(t + 1) - \mathbf{G}(t)$ be the forward difference operator. Then for any function \mathbf{G} defined on \mathbb{N}_a with $a \in \mathbb{R}$, the discrete delta Caputo-Fabrizio fractional difference in the Caputo sense and Caputo-Fabrizio fractional difference in the Riemann sense are defined by

$$\begin{aligned} \left({}^{CFC} \Delta^{\alpha} \mathbf{G}\right)(t) &= \frac{\mathbf{B}(\alpha)}{1 - \alpha} \sum_{\kappa=a}^{t-1} (\Delta_{\kappa} \mathbf{G})(\kappa)(1 + \lambda)^{t-\kappa-1} \\ &= \frac{\mathbf{B}(\alpha)}{1 - 2\alpha} \sum_{\kappa=a}^{t-1} (\Delta_{\kappa} \mathbf{G})(\kappa)(1 + \lambda)^{t-\kappa}, \quad [\forall t \in \mathbb{N}_{a+1}], \end{aligned} \quad (2.1)$$

and

$$\begin{aligned} \left({}^{CFR} \Delta^{\alpha} \mathbf{G}\right)(t) &= \frac{\mathbf{B}(\alpha)}{1 - \alpha} \Delta_t \sum_{\kappa=a}^{t-1} \mathbf{G}(\kappa)(1 + \lambda)^{t-\kappa-1} \\ &= \frac{\mathbf{B}(\alpha)}{1 - 2\alpha} \Delta_t \sum_{\kappa=a}^{t-1} \mathbf{G}(\kappa)(1 + \lambda)^{t-\kappa}, \quad [\forall t \in \mathbb{N}_{a+1}], \end{aligned} \quad (2.2)$$

respectively, where $\lambda = -\frac{\alpha}{1-\alpha}$ for $\alpha \in [0, 1)$, and $\mathbf{B}(\alpha)$ is a normalizing positive constant.

Moreover, for the higher order case when $q < \alpha < q + 1$ with $q \geq 0$, we have

$$\left({}^{CFC} \Delta^{\alpha} \mathbf{G}\right)(t) = \left({}^{CFC} \Delta^{\alpha-q} \Delta^q \mathbf{G}\right)(t), \quad [\forall t \in \mathbb{N}_{a+1}]. \quad (2.3)$$

Remark 2.1. It should be noted that

$$0 < 1 + \lambda = \frac{1 - 2\alpha}{1 - \alpha} < 1,$$

if $\alpha \in (0, \frac{1}{2})$, where (as above) $\lambda = -\frac{\alpha}{1-\alpha}$.

Definition 2.2 (see [29, 32]). Let G be defined on \mathbb{N}_a and $\alpha \in [1, 2]$. Then G is α -convex iff (ΔG) is $(\alpha - 1)$ -monotone increasing. That is,

$$G(t + 1) - \alpha G(t) + (\alpha - 1)G(t - 1) \geq 0, \quad [\forall t \in \mathbb{N}_{a+1}].$$

3. Convexity and positivity results

This section deals with convexity and positivity of the Caputo-Fabrizio operator in the Riemann sense (2.2). We first present some necessary lemmas.

Lemma 3.1. Let $G : \mathbb{N}_a \rightarrow \mathbb{R}$ be a function satisfying

$$\left({}^{CFC}_a \Delta^\alpha \Delta G\right)(t) \geq 0$$

and

$$(\Delta G)(a) \geq 0,$$

for $\alpha \in (0, \frac{1}{2})$ and t in \mathbb{N}_{a+2} . Then $(\Delta G)(t) \geq 0$, for every t in \mathbb{N}_{a+1} .

Proof. From Definition 2.1, we have for each $t \in \mathbb{N}_{a+2}$:

$$\begin{aligned} \left({}^{CFC}_a \Delta^\alpha \Delta G\right)(t) &= \frac{B(\alpha)}{1 - 2\alpha} \sum_{\kappa=a}^{t-1} \left(\Delta_\kappa^2 f\right)(\kappa)(1 + \lambda)^{t-\kappa} \\ &= \frac{B(\alpha)}{1 - 2\alpha} \left[\sum_{\kappa=a}^{t-1} (\Delta G)(\kappa + 1)(1 + \lambda)^{t-\kappa} - \sum_{\kappa=a}^{t-1} (\Delta G)(\kappa)(1 + \lambda)^{t-\kappa} \right] \\ &= \frac{B(\alpha)}{1 - 2\alpha} \left[(1 + \lambda)(\Delta G)(t) + \lambda \sum_{\kappa=a}^{t-1} (\Delta G)(\kappa)(1 + \lambda)^{t-\kappa} \right] \\ &= \frac{B(\alpha)}{1 - 2\alpha} \left[(1 + \lambda)(\Delta G)(t) - (1 + \lambda)^{t-a}(\Delta G)(a) + \lambda \sum_{\kappa=a+1}^{t-1} (\Delta G)(\kappa)(1 + \lambda)^{t-\kappa} \right]. \end{aligned} \quad (3.1)$$

Since $\frac{B(\alpha)}{1-2\alpha} > 0$, $1 + \lambda > 0$ and $\left({}^{CFC}_a \Delta^\alpha \Delta G\right)(t) \geq 0$ for all $t \in \mathbb{N}_{a+2}$, then (3.1) gives us

$$(\Delta G)(t) \geq (1 + \lambda)^{t-a-1}(\Delta G)(a) - \frac{\lambda}{1 + \lambda} \sum_{\kappa=a+1}^{t-1} (\Delta G)(\kappa)(1 + \lambda)^{t-\kappa}. \quad (3.2)$$

We will now show that $(\Delta G)(a+i+1) \geq 0$ if we assume that $(\Delta G)(a+i) \geq 0$ for some $i \in \mathbb{N}_1$. Note from our assumption we have that $(\Delta G)(a) \geq 0$. But then from the lower bound for $(\Delta G)(a+i+1)$ in (3.2) and our assumption we have

$$(\Delta G)(a+i+1) \geq \underbrace{(1+\lambda)^i (\Delta G)(a)}_{\geq 0} - \underbrace{\frac{\lambda}{1+\lambda} \sum_{\kappa=a+1}^{a+i} (\Delta G)(\kappa) (1+\lambda)^{a+i+1-\kappa}}_{\geq 0} \geq 0,$$

where we used $\frac{\lambda}{1+\lambda} < 0$. Thus, the result follows by induction. \square

Lemma 3.2. Let G be defined on \mathbb{N}_a and

$$\left({}^{CFC} \Delta^\alpha G\right)(t) \geq 0 \quad \text{with the initial values} \quad G(a+1) \geq G(a) \geq 0,$$

for $\alpha \in \left(1, \frac{3}{2}\right)$ and $t \in \mathbb{N}_{a+1}$. Then G is monotone increasing, positive and $\left(\frac{1}{2-\alpha}\right)$ -convex on \mathbb{N}_a .

Proof. From the definition with $q = 1$ we have

$$0 \leq \left({}^{CFC} \Delta^\alpha G\right)(t) = \left({}^{CFC} \Delta^{\alpha-1} \Delta G\right)(t), \quad [\forall t \in \mathbb{N}_{a+1}].$$

Since $(\Delta G)(a) \geq 0$ is given we have

$$(\Delta G)(t) \geq 0, \quad [\forall t \in \mathbb{N}_{a+1}],$$

by Lemma 3.1. This implies that G is a monotone increasing function. Therefore,

$$G(t) \geq G(t-1) \geq \dots \geq G(a+1) \geq G(a) \geq 0, \quad [\forall t \in \mathbb{N}_{a+1}],$$

and hence G is positive.

From the idea in Lemma 3.1 we have (here $\lambda = -\frac{\alpha-1}{2-\alpha}$ for $\alpha \in \left(1, \frac{3}{2}\right)$),

$$\begin{aligned} (\Delta G)(t) &\geq (1+\lambda)^{t-a-1} (\Delta G)(a) - \frac{\lambda}{1+\lambda} \sum_{\kappa=a+1}^{t-1} (\Delta G)(\kappa) (1+\lambda)^{t-\kappa} \\ &= \underbrace{(1+\lambda)^{t-a-1} (\Delta G)(a)}_{\geq 0} - \lambda (\Delta G)(t-1) - \underbrace{\frac{\lambda}{1+\lambda} \sum_{\kappa=a+1}^{t-2} (\Delta G)(\kappa) (1+\lambda)^{t-\kappa}}_{\geq 0 \text{ since } (\Delta G)(t) \geq 0} \\ &\geq -\lambda (\Delta G)(t-1) \\ &= \left(\frac{\alpha-1}{2-\alpha}\right) (\Delta G)(t-1) = \left(\frac{1}{2-\alpha} - 1\right) (\Delta G)(t-1). \end{aligned}$$

Consequently we have that G is $\left(\frac{1}{2-\alpha}\right)$ -convex on the set \mathbb{N}_a . \square

Lemma 3.3. Let G be defined on \mathbb{N}_a and

$$\left({}^{CFC} \Delta^\alpha G\right)(t) \geq 0 \quad \text{with} \quad (\Delta^2 G)(a) \geq 0,$$

for $\alpha \in \left(2, \frac{5}{2}\right)$ and $t \in \mathbb{N}_{a+1}$. Then, Then $(\Delta^2 G)(t) \geq 0$, for all $t \in \mathbb{N}_a$. Furthermore, one has G convex on the set \mathbb{N}_a .

Proof. Let $({}^{CFC}_a\Delta^\alpha \mathbf{G})(t) := \mathbf{F}(t)$ for each $t \in \mathbb{N}_{a+1}$. Since $\alpha \in (2, \frac{5}{2})$, we have:

$$({}^{CFC}_a\Delta^\alpha \mathbf{G})(t) = ({}^{CFC}_a\Delta^{\alpha-2} \Delta^2 \mathbf{G})(t) = ({}^{CFC}_a\Delta^{\alpha-2} \Delta \mathbf{F})(t) \geq 0,$$

for each $t \in \mathbb{N}_{a+1}$, and by assumption we have

$$(\Delta \mathbf{F})(a) = (\Delta^2 \mathbf{G})(a) \geq 0.$$

Then, using Lemma 3.2, we get

$$(\Delta \mathbf{F})(t) = (\Delta^2 \mathbf{G})(t) \geq 0$$

for each $t \in \mathbb{N}_{a+1}$. Hence, \mathbf{G} is convex on \mathbb{N}_a . □

Lemma 3.4. *Let \mathbf{G} be defined on \mathbb{N}_a and*

$$\Delta^2 ({}^{CFC}_a\Delta^\alpha \mathbf{G})(t) \geq 0$$

and

$$(\Delta \mathbf{G})(a+1) \geq (\Delta \mathbf{G})(a) \geq 0,$$

for $\alpha \in (0, \frac{1}{2})$ and $t \in \mathbb{N}_{a+1}$. Then $(\Delta^2 \mathbf{G})(t) \geq 0$, for each $t \in \mathbb{N}_a$.

Proof. For $t \in \mathbb{N}_{a+1}$, we have

$$\begin{aligned} \Delta ({}^{CFC}_a\Delta^\alpha \mathbf{G})(t) &= \frac{\mathbf{B}(\alpha)}{1-2\alpha} \Delta \left[\sum_{\kappa=a}^{t-1} (\Delta_\kappa \mathbf{G})(\kappa)(1+\lambda)^{t-\kappa} \right] \\ &= \frac{\mathbf{B}(\alpha)}{1-2\alpha} \left[\sum_{\kappa=a}^t (\Delta_\kappa \mathbf{G})(\kappa)(1+\lambda)^{t+1-\kappa} - \sum_{\kappa=a}^{t-1} (\Delta_\kappa \mathbf{G})(\kappa)(1+\lambda)^{t-\kappa} \right] \\ &= \frac{\mathbf{B}(\alpha)}{1-2\alpha} \left[(1+\lambda)(\Delta \mathbf{G})(t) + \sum_{\kappa=a}^{t-1} (\Delta_\kappa \mathbf{G})(\kappa)(1+\lambda)^{t+1-\kappa} - \sum_{\kappa=a}^{t-1} (\Delta_\kappa \mathbf{G})(\kappa)(1+\lambda)^{t-\kappa} \right] \\ &= \frac{\mathbf{B}(\alpha)}{1-2\alpha} \left[(1+\lambda)(\Delta \mathbf{G})(t) + \lambda \sum_{\kappa=a}^{t-1} (\Delta_\kappa \mathbf{G})(\kappa)(1+\lambda)^{t-\kappa} \right], \end{aligned} \tag{3.3}$$

where $\lambda = -\frac{\alpha}{1-\alpha}$. It follows from (3.3) that,

$$\begin{aligned} \Delta^2 ({}^{CFC}_a\Delta^\alpha \mathbf{G})(t) & \\ &= \frac{\mathbf{B}(\alpha)}{1-2\alpha} \Delta \left[(1+\lambda)(\Delta \mathbf{G})(t) + \lambda \sum_{\kappa=a}^{t-1} (\Delta_\kappa \mathbf{G})(\kappa)(1+\lambda)^{t-\kappa} \right] \\ &= \frac{(1+\lambda)\mathbf{B}(\alpha)}{1-2\alpha} (\Delta^2 \mathbf{G})(t) + \frac{\lambda \mathbf{B}(\alpha)}{1-2\alpha} \left[\sum_{\kappa=a}^t (\Delta_\kappa \mathbf{G})(\kappa)(1+\lambda)^{t+1-\kappa} - \sum_{\kappa=a}^{t-1} (\Delta_\kappa \mathbf{G})(\kappa)(1+\lambda)^{t-\kappa} \right] \\ &= \frac{(1+\lambda)\mathbf{B}(\alpha)}{1-2\alpha} (\Delta^2 \mathbf{G})(t) + \frac{\lambda \mathbf{B}(\alpha)}{1-2\alpha} \left[(1+\lambda)^{t+1-a} (\Delta \mathbf{G})(a) + \sum_{\kappa=a}^{t-1} (\Delta_\kappa \mathbf{G})(\kappa+1)(1+\lambda)^{t-\kappa} \right] \end{aligned} \tag{3.4}$$

$$\begin{aligned}
& - \sum_{\kappa=a}^{t-1} (\Delta_{\kappa} \mathbf{G})(\kappa)(1+\lambda)^{t-\kappa} \Big] \\
& = \frac{(1+\lambda)\mathbf{B}(\alpha)}{1-2\alpha} (\Delta^2 \mathbf{G})(t) + \frac{\lambda \mathbf{B}(\alpha)}{1-2\alpha} \left[(1+\lambda)^{t+1-a} (\Delta \mathbf{G})(a) + \sum_{\kappa=a}^{t-1} (\Delta_{\kappa}^2 \mathbf{G})(\kappa)(1+\lambda)^{t-\kappa} \right]. \quad (3.5)
\end{aligned}$$

Due to the nonnegativity of $\frac{(1+\lambda)\mathbf{B}(\alpha)}{1-2\alpha}$, from (3.4) we deduce

$$(\Delta^2 \mathbf{G})(t) \geq -\frac{\lambda}{1+\lambda} \left[(1+\lambda)^{t+1-a} (\Delta \mathbf{G})(a) + \sum_{\kappa=a}^{t-1} (\Delta_{\kappa}^2 \mathbf{G})(\kappa)(1+\lambda)^{t-\kappa} \right]. \quad (3.6)$$

By substituting $t = a + 1$ into (3.6), we get

$$\begin{aligned}
(\Delta^2 \mathbf{G})(a+1) & \geq -\frac{\lambda}{1+\lambda} \left[(1+\lambda)^2 (\Delta \mathbf{G})(a) + (\Delta^2 \mathbf{G})(a)(1+\lambda) \right] \\
& = \underbrace{\frac{\alpha}{(1-\alpha)}}_{>0} \left[\underbrace{(1+\lambda)(\Delta \mathbf{G})(a)}_{\geq 0} + \underbrace{(\Delta^2 \mathbf{G})(a)}_{\geq 0} \right] \geq 0.
\end{aligned}$$

Also, if we substitute $t = a + 2$ into (3.6), we obtain

$$\begin{aligned}
(\Delta^2 \mathbf{G})(a+2) & \geq -\frac{\lambda}{1+\lambda} \left[(1+\lambda)^3 (\Delta \mathbf{G})(a) + (1+\lambda)^2 (\Delta^2 \mathbf{G})(a) + (1+\lambda) (\Delta^2 \mathbf{G})(a+1) \right] \\
& = \underbrace{\frac{\alpha}{(1-\alpha)}}_{>0} \left[\underbrace{(1+\lambda)^2 (\Delta \mathbf{G})(a)}_{\geq 0} + \underbrace{(1+\lambda) (\Delta^2 \mathbf{G})(a)}_{\geq 0} + \underbrace{(\Delta^2 \mathbf{G})(a+1)}_{\geq 0} \right] \\
& \geq 0.
\end{aligned}$$

By continuing this process, we obtain that $(\Delta^2 \mathbf{G})(t) \geq 0$ for each $t \in \mathbb{N}_a$ as desired. \square

Now, we are in a position to state the first result on convexity. Furthermore, three representative results associated to different subregions in the space of (μ, ν) -parameter will be provided.

Theorem 3.1. Let \mathbf{G} be defined on \mathbb{N}_a with $\nu \in (0, \frac{1}{2})$ and $\mu \in (2, \frac{5}{2})$, and

$$\left({}^{CFC}_{a+1} \Delta^{\nu} {}^{CFC}_a \Delta^{\mu} \mathbf{G} \right) (t) \geq 0$$

and

$$(\Delta^2 \mathbf{G})(a+1) \geq (\Delta^2 \mathbf{G})(a) \geq 0,$$

for each $t \in \mathbb{N}_{a+1}$. Then \mathbf{G} is convex on the set \mathbb{N}_a .

Proof. Let $\left({}^{CFC}_a \Delta^{\mu} \mathbf{G} \right) (t) := \mathbf{F}(t)$ for each $t \in \mathbb{N}_{a+1}$. Then, by assumption we have

$$\left({}^{CFC}_{a+1} \Delta^{\nu} {}^{CFC}_a \Delta^{\mu} \mathbf{G} \right) (t) = \left({}^{CFC}_{a+1} \Delta^{\nu} \mathbf{F} \right) (t) \geq 0,$$

for each $t \in \mathbb{N}_{a+1}$. From the definition with $q = 2$ we have

$$\begin{aligned} F(a+1) &= \left({}^{CFC} \Delta_a^\mu \mathbf{G}\right)(a+1) = \left({}^{CFC} \Delta_a^{\mu-2} \Delta^2 \mathbf{G}\right)(a+1) \\ &= \frac{\mathbf{B}(\mu-2)}{5-2\mu} \sum_{\kappa=a}^a (\Delta^\kappa \mathbf{G})(\kappa) (1+\lambda_\mu)^{a-\kappa} \\ &= \frac{\mathbf{B}(\mu-2)}{5-2\mu} \underbrace{(\Delta^3 \mathbf{G})(a)}_{\geq 0 \text{ by assumption}} \\ &\geq 0, \end{aligned}$$

where $\lambda_\mu = -\frac{\mu-2}{3-\mu}$. Since $(\Delta^2 \mathbf{G})(a) \geq 0$, we find that $(\Delta^2 \mathbf{G})(t) \geq 0$ for each $t \in \mathbb{N}_a$. Furthermore, we see that \mathbf{G} is convex on \mathbb{N}_a from Lemma 3.3. \square

Theorem 3.2. Let \mathbf{G} be defined on \mathbb{N}_a with $\nu \in \left(1, \frac{3}{2}\right)$ and $\mu \in \left(2, \frac{5}{2}\right)$, and

$$\begin{aligned} \left({}^{CFC} \Delta_{a+1}^\nu {}^{CFC} \Delta_a^\mu \mathbf{G}\right)(t) &\geq 0, \\ (\Delta^2 \mathbf{G})(a+2) &\geq \frac{1}{3-\mu} (\Delta \mathbf{G}^2)(a+1) \geq 0, \end{aligned}$$

and

$$(\Delta^2 \mathbf{G})(a+1) \geq (\Delta^2 \mathbf{G})(a) \geq 0,$$

for each $t \in \mathbb{N}_{a+1}$. Then \mathbf{G} is convex on \mathbb{N}_a .

Proof. Let $F(t) := \left({}^{CFC} \Delta_a^\mu \mathbf{G}\right)(t)$. Note that:

$$\left({}^{CFC} \Delta_{a+1}^\nu {}^{CFC} \Delta_a^\mu \mathbf{G}\right)(t) = \left({}^{CFC} \Delta_{a+1}^\nu F\right)(t) \geq 0,$$

for $t \in \mathbb{N}_{a+1}$. Then we have

$$\begin{aligned} F(a+1) &= \left({}^{CFC} \Delta_a^{\mu-2} \Delta^2 \mathbf{G}\right)(a+1) \\ &= \frac{\mathbf{B}(\mu-2)}{5-2\mu} \sum_{\kappa=a}^a (\Delta^3 \mathbf{G})(\kappa) (1+\lambda_\mu)^{a+1-\kappa} \\ &= \frac{\mathbf{B}(\mu-2)}{5-2\mu} (1+\lambda_\mu) (\Delta^3 \mathbf{G})(a) \geq 0, \end{aligned} \tag{3.7}$$

and

$$\begin{aligned} F(a+2) &= \left({}^{CFC} \Delta_a^{\mu-2} \Delta^2 \mathbf{G}\right)(a+2) \\ &= \frac{\mathbf{B}(\mu-2)}{5-2\mu} \sum_{\kappa=a}^{a+1} (\Delta^3 \mathbf{G})(\kappa) (1+\lambda_\mu)^{a+2-\kappa} \\ &= \frac{\mathbf{B}(\mu-2)}{5-2\mu} \left[(1+\lambda_\mu)^2 (\Delta^3 \mathbf{G})(a) + (1+\lambda_\mu) (\Delta^3 \mathbf{G})(a+1) \right] \end{aligned}$$

$$\begin{aligned}
&= \frac{(1 + \lambda_\mu)\mathbf{B}(\mu - 2)}{5 - 2\mu} \left[(1 + \lambda_\mu) \left[(\Delta^2\mathbf{G})(a + 1) - (\Delta^2\mathbf{G})(a) \right] \right. \\
&\quad \left. + \left[(\Delta^2\mathbf{G})(a + 2) - (\Delta^2\mathbf{G})(a + 1) \right] \right] \\
&\geq \frac{(1 + \lambda_\mu)\mathbf{B}(\mu - 2)}{5 - 2\mu} \left[\frac{1}{3 - \mu} - 1 \right] \geq 0,
\end{aligned} \tag{3.8}$$

where $\lambda_\mu = -\frac{\mu-2}{5-2\mu}$. On the other hand, one has

$$\begin{aligned}
\mathbf{F}(a + 2) - \mathbf{F}(a + 1) &= \frac{(1 + \lambda_\mu)\mathbf{B}(\mu - 2)}{5 - 2\mu} \left[(1 + \lambda_\mu)(\Delta^3\mathbf{G})(a) + (\Delta^3\mathbf{G})(a + 1) - (\Delta^3\mathbf{G})(a) \right] \\
&\geq \frac{(1 + \lambda_\mu)\mathbf{B}(\mu - 2)}{5 - 2\mu} \left(\lambda_\mu - 1 + \frac{1}{3 - \mu} \right) (\Delta^2\mathbf{G})(a + 1) \geq 0.
\end{aligned} \tag{3.9}$$

Then, from Eqs (3.7)–(3.9), we see that $\mathbf{F}(a + 2) \geq \mathbf{F}(a + 1) \geq 0$. Therefore, Lemma 3.2 gives

$$\mathbf{F}(t) = \left({}^{CFC}_{a+1}\Delta^\nu \mathbf{G} \right) (t) \geq 0$$

for all $t \in \mathbb{N}_{a+1}$. Moreover, by considering $(\Delta^2\mathbf{G})(a) \geq 0$ in Lemma 3.3, we can deduce that \mathbf{G} is convex on the set \mathbb{N}_a . \square

Theorem 3.3. Let \mathbf{G} be defined on \mathbb{N}_a with $\nu \in \left(2, \frac{5}{2}\right)$ and $\mu \in \left(0, \frac{1}{2}\right)$, and

$$\begin{aligned}
&\left({}^{CFC}_{a+1}\Delta^\nu {}^{CFC}_a\Delta^\mu \mathbf{G} \right) (t) \geq 0, \\
&(\Delta\mathbf{G})(a + 2) \geq \frac{1}{1 - \mu} (\Delta\mathbf{G})(a + 1) \geq 0,
\end{aligned}$$

and

$$(\Delta\mathbf{G})(a + 1) \geq (\Delta\mathbf{G})(a) \geq 0,$$

for each $t \in \mathbb{N}_{a+1}$. Then we have that \mathbf{G} is convex on \mathbb{N}_a .

Proof. Again, we write $\mathbf{F}(t) := \left({}^{CFC}_a\Delta^\mu \mathbf{G} \right) (t)$, and therefore, $\left({}^{CFC}_{a+1}\Delta^\nu \mathbf{F} \right) (t) \geq 0$ by assumption, for each $t \in \mathbb{N}_{a+1}$. Then, we see that

$$\begin{aligned}
(\Delta^2\mathbf{F})(a + 1) &\equiv \Delta^2 \left({}^{CFC}_a\Delta^\mu \mathbf{G} \right) (a + 1) \\
&\stackrel{\text{by}}{\underset{(3.4)}{=}} \frac{(1 + \lambda_\mu)\mathbf{B}(\mu)}{1 - 2\mu} (\Delta^2\mathbf{G})(a + 1) \\
&\quad + \frac{\lambda_\mu \mathbf{B}(\mu)}{1 - 2\mu} \left[(1 + \lambda_\mu)^2 (\Delta\mathbf{G})(a) + \sum_{\kappa=a}^a \left(\Delta_\kappa^2 \mathbf{G} \right) (\kappa) (1 + \lambda_\mu)^{a+1-\kappa} \right] \\
&= \frac{(1 + \lambda_\mu)\mathbf{B}(\mu)}{1 - 2\mu} \left[(\Delta^2\mathbf{G})(a + 1) + \lambda_\mu (1 + \lambda_\mu) (\Delta\mathbf{G})(a) + \lambda_\mu (\Delta^2\mathbf{G})(a) \right] \\
&= \frac{(1 + \lambda_\mu)\mathbf{B}(\mu)}{1 - 2\mu} \left[(\Delta\mathbf{G})(a + 2) + (\lambda_\mu - 1) (\Delta\mathbf{G})(a + 1) + \lambda_\mu^2 (\Delta\mathbf{G})(a) \right]
\end{aligned}$$

$$\begin{aligned}
&\geq \frac{(1 + \lambda_\mu)\mathbf{B}(\mu)}{1 - 2\mu} \left[\frac{1}{1 - \mu}(\Delta\mathbf{G})(a + 1) + (\lambda_\mu - 1)(\Delta\mathbf{G})(a + 1) + \underbrace{\lambda_\mu^2(\Delta\mathbf{G})(a)}_{\geq 0} \right] \\
&\geq \frac{(1 + \lambda_\mu)\mathbf{B}(\mu)}{1 - 2\mu} \left[\frac{1}{1 - \mu} + \lambda_\mu - 1 \right] (\Delta\mathbf{G})(a + 1) \\
&\geq 0,
\end{aligned}$$

where $\lambda_\mu = -\frac{\mu}{1-\mu}$. It follows that,

$$(\Delta^2\mathbf{F})(t) = \Delta^2 \left({}^{CF C} \Delta^\mu \mathbf{G} \right) (t) \geq 0,$$

for each $t \in \mathbb{N}_a$ by Lemma 3.3. Considering, $(\Delta^2\mathbf{G})(a) \geq 0$, we can deduce that \mathbf{G} is convex on \mathbb{N}_a by Lemma 3.4. \square

In Figure 1, we demonstrate the regions of the (μ, ν) -parameter space in which the above three Theorems 3.1–3.3 are applied.




Figure 1. Three different regions concerning Theorems 3.1–3.3.

4. Conclusions

In this study, we present some new positivity results for discrete fractional operators with exponential kernels in the sense of Caputo. In particular new positivity, α -convexity and α -monotonicity were presented. We now refer the reader to observations for discrete generalized fractional operators in [33] which combined with this paper may motivate future work.

Author's contributions

Conceptualization, P.O.M., D.O., A.B.B. and D.B.; methodology, P.O.M., D.O.; software, D.O., D.B., K.M.A., A.B.B.; validation, P.O.M., D.O., D.B. and A.B.B.; formal analysis, K.M.A.; investigation, P.O.M., D.O., K.M.A.; resources, A.B.B.; writing-original draft preparation, P.O.M., D.O., D.B., K.M.A., A.B.B.; writing-review and editing, D.O., D.B. and A.B.B.; funding acquisition, D.B. and K.M.A. All authors read and approved the final manuscript.

Acknowledgements

This Research was supported by Taif University Researchers Supporting Project Number (TURSP-2020/217), Taif University, Taif, Saudi Arabia.

Conflict of interest

The authors declare that they have no conflicts of interest.

References

1. C. Goodrich, A. C. Peterson, *Discrete fractional calculus*, Springer, New York, 2015. <https://doi.org/10.1007/978-3-319-25562-0>
2. F. M. Atici, P. W. Eloe, A transform method in discrete fractional calculus, *Int. J. Differ. Equ.*, **2** (2007), 165–176.
3. H. M. Srivastava, P. O. Mohammed, C. S. Ryou, Y. S. Hamed, Existence and uniqueness of a class of uncertain Liouville-Caputo fractional difference equations, *J. King Saud Univ. Sci.*, **33** (2021), 101497. <https://doi.org/10.1016/j.jksus.2021.101497>
4. C. Lizama, The Poisson distribution, abstract fractional difference equations, and stability, *Proc. Amer. Math. Soc.*, **145** (2017), 3809–3827. <https://doi.org/10.1090/proc/12895>
5. C. Lizama, M. Murillo-Arcila, Well posedness for semidiscrete fractional Cauchy problems with finite delay, *J. Comput. Appl. Math.*, **339** (2018), 356–366. <https://doi.org/10.1016/j.cam.2017.07.027>
6. H. M. Srivastava, P. O. Mohammed, J. L. G. Guirao, Y. S. Hamed, Link theorem and distributions of solutions to uncertain Liouville-Caputo difference equations, *Discrete Contin. Dyn. Syst.*, **15** (2021), 427–440. <http://dx.doi.org/10.3934/dcdss.2021083>
7. İlhan, E. Analysis of the spread of Hookworm infection with Caputo-Fabrizio fractional derivative, *Turk. J. Sci.*, **7** (2022), 43–52.
8. M. A. Ragusa, Parabolic Herz spaces and their applications, *Appl. Math. Lett.*, **25** (2022), 1270–1273. <https://doi.org/10.1016/j.aml.2011.11.022>
9. S. Rezapour, A. Boulfoul, B. Tellab, M. E. Samei, S. Etemad, R. George, Fixed point theory and the Liouville-Caputo integro-differential FBVP with multiple nonlinear terms, *J. Funct. Spaces*, **2022** (2022), 6713533. <https://doi.org/10.1155/2022/6713533>
10. T. Abdeljawad, On Riemann and Caputo fractional differences, *Comput. Math. Appl.*, **62** (2011), 1602–1611. <https://doi.org/10.1016/j.camwa.2011.03.036>
11. T. Abdeljawad, Q. M. Al-Mdallal, M. A. Hajji, Arbitrary order fractional difference operators with discrete exponential kernels and applications, *Discrete Dyn. Nat. Soc.*, **2017** (2017), 4149320. <https://doi.org/10.1155/2017/4149320>
12. T. Abdeljawad, F. Jarad, A. Atangana, P. O. Mohammed, On a new type of fractional difference operators on h-step isolated time scales, *J. Fract. Calc. Nonlinear Syst.*, **1** (2020), 46–74.

13. T. Abdeljawad, Different type kernel h -fractional differences and their fractional h -sums, *Chaos Soliton. Fract.*, **116** (2018), 146–156. <https://doi.org/10.1016/j.chaos.2018.09.022>
14. R. A. C. Ferreira, D. F. M. Torres, Fractional h -difference equations arising from the calculus of variations, *Appl. Anal. Discrete Math.*, **5** (2011), 110–121.
15. C. R. Chen, M. Bohner, B. G. Jia, Ulam-Hyers stability of Caputo fractional difference equations, *Math. Meth. Appl. Sci.*, **42** (2019), 7461–7470. <https://doi.org/10.1002/mma.5869>
16. G. Wu, D. Baleanu, Discrete chaos in fractional delayed logistic maps, *Nonlinear Dyn.*, **80** (2015), 1697–1703. <https://doi.org/10.1007/s11071-014-1250-3>
17. F. Atici, S. Sengul, Modeling with discrete fractional equations, *J. Math. Anal. Appl.*, **369** (2010), 1–9.
18. B. Ghanbari, On the modeling of the interaction between tumor growth and the immune system using some new fractional and fractional-fractal operators, *Adv. Differ. Equ.*, **2020** (2020), 585. <https://doi.org/10.1186/s13662-020-03040-x>
19. R. Dahal, C. S. Goodrich, A monotonicity result for discrete fractional difference operators, *Arch. Math.*, **102** (2014), 293–299. <https://doi.org/10.1007/s00013-014-0620-x>
20. P. O. Mohammed, T. Abdeljawad, F. K. Hamasalh, On Riemann-Liouville and Caputo fractional forward difference monotonicity analysis, *Mathematics*, **9** (2021), 1303. <https://doi.org/10.3390/math9111303>
21. C. S. Goodrich, M. Muellner, An analysis of the sharpness of monotonicity results via homotopy for sequential fractional operators, *Appl. Math. Lett.*, **98** (2019), 446–452. <https://doi.org/10.1016/j.aml.2019.07.003>
22. B. Jia, L. Erbe, A. Peterson, Two monotonicity results for nabla and delta fractional differences, *Arch. Math.*, **104** (2015), 589–597. <https://doi.org/10.1007/s00013-015-0765-2>
23. C. S. Goodrich, Sharp monotonicity results for fractional nabla sequential differences, *J. Differ. Equ. Appl.*, **25** (2019), 801–814. <https://doi.org/10.1080/10236198.2018.1542431>
24. P. O. Mohammed, O. Almutairi, R. P. Agarwal, Y. S. Hamed, On convexity, monotonicity and positivity analysis for discrete fractional operators defined using exponential kernels, *Fractal Fract.*, **6** (2022), 55. <https://doi.org/10.3390/fractalfract6020055>
25. P. O. Mohammed, F. K. Hamasalh, T. Abdeljawad, Difference monotonicity analysis on discrete fractional operators with discrete generalized Mittag-Leffler kernels, *Adv. Differ. Equ.*, **2021** (2021), 213. <https://doi.org/10.1186/s13662-021-03372-2>
26. L. Erbe, C. S. Goodrich, B. Jia, A. Peterson, Survey of the qualitative properties of fractional difference operators: Monotonicity, convexity, and asymptotic behavior of solutions, *Adv. Differ. Equ.*, **2016** (2016), 43. <https://doi.org/10.1186/s13662-016-0760-3>
27. C. S. Goodrich, A note on convexity, concavity, and growth conditions in discrete fractional calculus with delta difference, *Math. Inequal. Appl.*, **19** (2016), 769–779. <https://doi.org/10.7153/MIA-19-57>
28. C. S. Goodrich, C. Lizama, Positivity, monotonicity, and convexity for convolution operators, *Discrete Contin. Dyn. Syst.*, **40** (2020), 4961–4983. <http://dx.doi.org/10.3934/dcds.2020207>

29. C. S. Goodrich, C. Lizama, A transference principle for nonlocal operators using a convolutional approach: Fractional monotonicity and convexity, *Isr. J. Math.*, **236** (2020), 533–589. <https://doi.org/10.1007/s11856-020-1991-2>
30. D. Mozyrska, D. F. M. Torres, M. Wyrwas, Solutions of systems with the Caputo-Fabrizio fractional delta derivative on time scales, *Nonlinear Anal. Hybrid Syst.*, **32** (2019), 168–176. <https://doi.org/10.1016/j.nahs.2018.12.001>
31. P. O. Mohammed, T. Abdeljawad, F. K. Hamasalh, On discrete delta Caputo-Fabrizio fractional operators and monotonicity analysis, *Fractal Fract.*, **5** (2021), 116. <https://doi.org/10.3390/fractalfract5030116>
32. C. S. Goodrich, J. M. Jonnalagadda, B. Lyons, Convexity, monotonicity and positivity results for sequential fractional nabla difference operators with discrete exponential kernels, *Math. Meth. Appl. Sci.*, **44** (2021), 7099–7120. <https://doi.org/10.1002/mma.7247>
33. P. O. Mohammed, T. Abdeljawad, Discrete generalized fractional operators defined using h -discrete Mittag-Leffler kernels and applications to AB fractional difference systems, *Math. Meth. Appl. Sci.*, 2020. <https://doi.org/10.1002/mma.7083>



AIMS Press

©2022 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)