

NEW MULTI-FUNCTIONAL APPROACH FOR κ TH-ORDER DIFFERENTIABILITY GOVERNED BY FRACTIONAL CALCULUS VIA APPROXIMATELY GENERALIZED (ψ, \hbar) -CONVEX FUNCTIONS IN HILBERT SPACE

MIAO-KUN WANG

*Department of Mathematics, Huzhou University
Huzhou 313000, P. R. China
wmk000@126.com*

SAIMA RASHID

*Department of Mathematics
Government College University
Faisalabad 38000, Pakistan
saimarashid@gcuf.edu.pk*

YELIZ KARACA

*University of Massachusetts Medical School
Worcester, MA 01655, USA
yeliz.karaca@ieee.org*

DUMITRU BALEANU

*Department of Mathematics
Cankaya University, Ankara, Turkey
dumitru@cankaya.edu.tr*

*Corresponding author.

This is an Open Access article in the “Special Issue Section on Fractal AI-Based Analyses and Applications to Complex Systems: Part I”, edited by Yeliz Karaca (University of Massachusetts Medical School, USA), Dumitru Baleanu (Cankaya University, Turkey), Majaz Moonis (University of Massachusetts Medical School, USA), Khan Muhammad (Sejong University, South Korea), Yu-Dong Zhang (University of Leicester, UK) & Osvaldo Gervasi (Perugia University, Italy) published by World Scientific Publishing Company. It is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 (CC-BY-NC-ND) License which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

YU-MING CHU*
College of Science, Hunan City University
Yiyang 413000, P. R. China
chuyuming2005@126.com

Received August 22, 2020
Accepted November 23, 2020
Published March 25, 2021

Abstract

This work addresses several novel classes of convex function involving arbitrary non-negative function, which is known as approximately generalized (ψ, \hbar) -convex and approximately ψ -quasiconvex function, with respect to Raina's function, which are elaborated in Hilbert space. To ensure the feasibility of the proposed concept and with the discussion of special cases, it is presented that these classes generate other classes of generalized (ψ, \hbar) -convex functions such as higher-order strongly (HOS) generalized (ψ, \hbar) -convex functions and HOS generalized ψ -quasiconvex function. The core of the proposed method is a newly developed Simpson's type of identity in the settings of Riemann–Liouville fractional integral operator. Based on the HOS generalized (ψ, \hbar) -convex function representation, we established several theorems and related novel consequences. The presented results demonstrate better performance for HOS generalized ψ -quasiconvex functions where we can generate several other novel classes for convex functions that exist in the relative literature. Accordingly, the assortment in this study aims at presenting a direction in the related fields.

Keywords: Convex Functions; Generalized (ψ, \hbar) -Convex Functions; κ -th Order Differentiability; Raina's Function; Breckner-Type Function; Godunova–Levin-Type Function.

1. INTRODUCTION

In the last three decades, the most intriguing and captivating subject of current research in mathematical sciences is the fractional calculus as well as derivatives and integrals of non-integer order are involved. The fractional operators of the said calculus is the most essential phenomenon in the real world, and it has been treated as the crucial and exceptionally enormous factor in precisely portraying the conduct of oscillators, medicine, mechanical devices, electrical systems, granular soils, circuits, and financial systems has been represented in the continuous composition, e.g. see Refs. 1–3. Subsequently, the dynamical developments of authentic physical structures have been even more fittingly investigated by fractional-order differential and/or difference equations rather than the integer-order ones. Given the way that the high constancy model of physical frameworks can be portrayed by fractional-order frameworks, the region has obtained a lot of enthusiasm for the control

network which has concentrated on stability and control problems in systems represented by non-integer-order differential equations.^{4,5}

On the other hand, the study of the fractional integral inequalities such as in the case of differential and difference equations is of importance in various subjects.^{6–9} Inequalities that are used in structural applications are of practical importance, as they have proven that fractional behavior has a profound impact on the performance of fractional integral inequalities. For instance, Sarikaya *et al.*¹⁰ reported the Hermite–Hadamard-type inequalities for Riemann–Liouville fractional operators. Set *et al.*¹¹ established valuable consequences by utilizing fractional integral operators. The inequality theory is of importance in many fields such as in the field of solid-state physics, materials and metallurgy sciences because it is causally related to different microscopic physics processes and also can be used as an experimental investigation of these procedures.^{7–9} Fractional integral

inequalities play a significant role in both pure and applied sciences because of their wide applications as well as many other natural and human social sciences, while convexity theory has remained as an important tool in the establishment of the theory of integral inequalities. Simpson's inequality, as a member of the family of integral inequalities, is a classic inequality that has long been fascinated by numerous mathematical researchers, which can be stated as follows:

$$\left| \frac{1}{6} \left[\mathcal{Q}(\omega_1) + 4\mathcal{Q}\left(\frac{\omega_1 + \omega_2}{2}\right) + \mathcal{Q}(\omega_2) \right] - \frac{1}{\omega_2 - \omega_1} \int_{\omega_1}^{\omega_2} \mathcal{Q}(z) dz \right| \leq \frac{\|\mathcal{Q}^{(4)}\|_{\infty} (\omega_2 - \omega_1)^4}{2880}, \quad (1)$$

where $\mathcal{Q} : [\omega_1, \omega_2] \rightarrow \mathbb{R}$ is fourth-order differentiable function on (ω_1, ω_2) with the condition that

$$\|\mathcal{Q}^{(4)}\|_{\infty} = \sup_{z \in (\omega_1, \omega_2)} |\mathcal{Q}^{(4)}(z)| < \infty.$$

In recent years, several successful attempts have been made in obtaining the variants and applications of Simpson's type of inequality. For example, Dragomir *et al.*¹² provided many interesting results on its applications in numerical integration. Rashid *et al.*¹³ contemplated the novel Simpson's-type inequalities in the settings of fractional calculus with applications. In Ref. 14, Li *et al.* obtained several extensions of Simpson's-type inequalities via extended (s, m) -convex functions. For more developments, generalizations and variants for Simpson's type of inequality have been the subject of much research, see Refs. 15 and 16.

In Ref. 17, Varosanec proposed a class of convex functions amplify and unifies several existing ideas of classical convex functions, encompassing Breckner-type convex functions,¹⁸ P -functions,¹⁹ Godunova–Levin-type convex, and Q -functions.²⁰ We acknowledge that this class plays a dominant role in convex analysis and this class plays a significant contribution to convexity theory and provides assistance to explore numerous novel classes of a convex function, see Refs. 21 and 22 and the references therein.

In Ref. 23, Polyak introduced and studied a new class of functionals that has significant importance in machine learning models, optimization theory and many other related areas. The existence of

nonlinear complementary problems can be determined by strong convexity.²⁴ The convergence criteria of iterative schemes for variational and equilibrium issues are contemplated by Zu and Marcotte.²⁵ Bynum²⁶ and Chen *et al.*²⁷ introduced the general characterization and utilization of the parallelogram laws for the Banach spaces. Xu²⁸ explored new attributes of p -uniform convexity and q -uniform smoothness of a Banach space utilizing $\|\cdot\|^p$ and $\|\cdot\|^q$, respectively. In Ref. 29, Nikodem and Pales explored the new and novel utilities of the characterization of the inner product space by considering strongly convex functions. Interestingly, the Polyak–Lojasiewicz condition is fulfilled with the assembly of stochastic slope descent for the class of functions based on strongly-convex functions too as a wide scope of non-convex functions incorporating those utilized in machine-learning applications.³⁰ In their recent work, authors³¹ reported results for differentiable higher-order strongly (HOS) \hbar -convex functions. In Ref. 32, the authors established the predominating η -convex functions in a wide broadway and also studied several generalizations for predominating η -quasiconvex functions. For further interesting papers related to HOS convex functions, see Refs. 33–35.

In response to the existing approaches, this paper aims to create refinements of Simpson-type inequalities utilizing approximately generalized (ψ, \hbar) -convex, approximately generalized (ψ, \hbar) -quasiconvex, HOS generalized (ψ, \hbar) -convex and HOS generalized ψ -quasiconvex functions with respect to Raina's function that not only possess the key properties of the classical convex functions but also have exceptional parameters adjustment with and without fixing the non-negative arbitrary function \hbar . Moreover, our consequences are correlated with the auxiliary identity which associates with fractional calculus for κ th-order differentiable functions. Numerous refinements of Simpson's type of inequality are derived that can be used to characterize the uniformly reflex Banach space. The significant advantage of these outcomes is that they used to comprehend the parallelogram laws for L^p spaces. Several remarkable cases are provided to exhibit the novelty of the results given herein.

2. PRELIMINARIES

Initially, suppose a non-empty set $\mathcal{K}_{\mathcal{F}}$ in a real Hilbert space \mathcal{H} . The inner product and norm are denoted by the symbols $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$, respectively.

Moreover, there is an arbitrary non-negative function $h : (0, 1) \rightarrow \mathbb{R}$, $\eta = \{\eta(m)\}_{m=0}^\infty$ be a bounded sequence of real numbers and $\mathcal{F}_{v_1, v_2}^\eta(\cdot)$, $v_1, v_2 > 0$ denotes Raina's function. In Ref. 36, Raina explored a new class of functions stated as follows:

$$\mathcal{F}_{v_1, v_2}^\eta(z) = \mathcal{F}_{v_1, v_2}^{\eta(0), \eta(1), \dots}(z) = \sum_{m=0}^\infty \frac{\eta(m)}{\Gamma(v_1 m + \eta)} z^m, \tag{2}$$

where $v_1, v_2 > 0$, $|z| < \mathcal{R}$ and

$$\eta = (\eta(0), \dots, \eta(m), \dots)$$

is bounded sequence of positive real numbers. Notice that if we choose $v_1 = 1, v_2 = 0$ in Eq. (2), then

$$\eta(m) = \frac{(\delta_1)_m (\delta_2)_m}{(\delta_3)_m} \quad \text{for } m = 0, 1, 2, \dots,$$

where δ_1, δ_2 and δ_3 are parameters which can take arbitrary real and complex values (provided that $\delta_3 \neq 0, -1, -2, \dots$) and we denotes the symbol $(b)_m$ by

$$(b)_m = \frac{\Gamma(b+m)}{\Gamma(b)} = b(b+1) \dots (b+m-1), \quad m = 0, 1, 2, \dots,$$

and restrict its domain to $|z| \leq 1$ (with $z \in \mathbb{C}$), then we have the classical hypergeometric function, which is defined as

$$\begin{aligned} \mathcal{F}_{v_1, v_2}^\eta(z) &= F(\delta_1, \delta_2; \delta_3; z) \\ &= \sum_{m=0}^\infty \frac{(\delta_1)_m (\delta_2)_m}{m! (\delta_3)_m} z^m. \end{aligned}$$

Also, if $\eta = (1, 1, \dots)$ with $\varsigma = \delta$, ($\Re(\delta) > 0$), $\eta = 1$ and restricting its domain to $z \in \mathbb{C}$ in Eq. (2), then we have the classical Mittag-Leffler function:

$$E_{\delta_1}(z) = \sum_{m=0}^\infty \frac{1}{\Gamma(1 + \delta_1 m)} z^{\kappa}.$$

Next, we evoke a new class of set and a new class of functions including Raina's functions.

Definition 1 (Ref. 37). A non-empty set $\mathcal{K}_{\mathcal{F}}$ is said to be generalized ψ -convex set, if

$$\omega_1 + \xi \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \in \mathcal{K}_{\mathcal{F}} \tag{3}$$

for all $\omega_1, \omega_2 \in \mathcal{K}_{\mathcal{F}}, \xi \in [0, 1]$.

We now define the generalized ψ -convex function presented by Cortez et al.³⁷

Definition 2 (Ref. 37). Let a set $\mathcal{K}_{\mathcal{F}} \subseteq \mathbb{R}$ and a function $\mathcal{Q} : \mathcal{K}_{\mathcal{F}} \rightarrow \mathbb{R}$ is said to be generalized ψ -convex, if

$$\mathcal{Q}(\omega_1 + \xi \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) \leq (1 - \xi)\mathcal{Q}(\omega_1) + \xi\mathcal{Q}(\omega_2) \tag{4}$$

for all $\omega_1, \omega_2 \in \mathcal{K}_{\mathcal{F}}, \xi \in [0, 1]$.

Next, we present a new class of generalized ψ -convex functions for an arbitrary non-negative function h .

Definition 3. Let $h : (0, 1) \rightarrow \mathbb{R}$ be a real function and a function $\mathcal{Q} : \mathcal{K}_{\mathcal{F}} \rightarrow \mathbb{R}$ is said to be generalized (ψ, h) -convex function, if

$$\begin{aligned} \mathcal{Q}(\omega_1 + \xi \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) &\leq h(1 - \xi)\mathcal{Q}(\omega_1) \\ &+ h(\xi)\mathcal{Q}(\omega_2) \end{aligned} \tag{5}$$

for all $\omega_1, \omega_2 \in \mathcal{K}_{\mathcal{F}}, \xi \in [0, 1]$.

Further, we demonstrate several novel concepts of generalized ψ -convex functions with respect to an arbitrary non-negative function.

Definition 4. Let $h : (0, 1) \rightarrow \mathbb{R}$ be a real function and a function $\mathcal{Q} : \mathcal{K}_{\mathcal{F}} \rightarrow \mathbb{R}$ is said to be approximately generalized (ψ, h) -convex, if

$$\begin{aligned} \mathcal{Q}(\omega_1 + \xi \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) &\leq h(1 - \xi)\mathcal{Q}(\omega_1) + h(\xi)\mathcal{Q}(\omega_2) + \mathbb{D}(\omega_1, \omega_2) \end{aligned} \tag{6}$$

for all $\omega_1, \omega_2 \in \mathcal{K}_{\mathcal{F}}, \xi \in [0, 1]$.

Some noteworthy cases of Definition 4 are demonstrated as follows:

(I) For some $c \geq 0, p > 2$ and taking $\mathbb{D}(\omega_1, \omega_2) = -c\{\xi^p(1 - \xi) + (1 - \xi)^p\xi\} \|\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)\|^p$, then we have a new concept of HOS generalized (ψ, h) -convex function with respect to an arbitrary non-negative function h .

Definition 5. Let $h : (0, 1) \rightarrow \mathbb{R}$ be a real function and a function $\mathcal{Q} : \mathcal{K}_{\mathcal{F}} \rightarrow \mathbb{R}$ is said to be HOS generalized (ψ, h) -convex having $c \geq 0$, if

$$\begin{aligned} \mathcal{Q}(\omega_1 + \xi \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) &\leq h(1 - \xi)\mathcal{Q}(\omega_1) + h(\xi)\mathcal{Q}(\omega_2) \\ &- c\{\xi^p(1 - \xi) + (1 - \xi)^p\xi\} \\ &\times \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)^p \end{aligned} \tag{7}$$

for all $\omega_1, \omega_2 \in \mathcal{K}_{\mathcal{F}}, \xi \in [0, 1]$.

(II) Taking $\mathbb{D}(\omega_1, \omega_2) = -c\{\xi^p(1 - \xi) + (1 - \xi)^p\xi\}\|\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)\|^p$ with $p = 2$ and $c \geq 0$, then we have a new concept of strongly generalized (ψ, \hbar) -convex function with respect to an arbitrary non-negative function \hbar .

Definition 6. Let $\hbar : (0, 1) \rightarrow \mathbb{R}$ be a real function and a function $\mathcal{Q} : \mathcal{K}_{\mathcal{F}} \rightarrow \mathbb{R}$ is said to be strongly generalized (ψ, \hbar) -convex having $c \geq 0$, if

$$\begin{aligned} \mathcal{Q}(\omega_1 + \xi\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) &\leq \hbar(1 - \xi)\mathcal{Q}(\omega_1) \\ &+ \hbar(\xi)\mathcal{Q}(\omega_2) - c\xi(1 - \xi)\|\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)\|^2 \end{aligned} \tag{8}$$

for all $\omega_1, \omega_2 \in \mathcal{K}_{\mathcal{F}}, \xi \in [0, 1]$.

(III) For some $c \geq 0, p > 2$ with $\hbar(\xi) = \xi$ and taking $\mathbb{D}(\omega_1, \omega_2) = -c\{\xi^p(1 - \xi) + (1 - \xi)^p\xi\}\|\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)\|^p$, then we have a new concept of HOS generalized ψ -convex function.

Definition 7. Let a function $\mathcal{Q} : \mathcal{K}_{\mathcal{F}} \rightarrow \mathbb{R}$ is said to be HOS generalized ψ -convex having $c \geq 0$, if

$$\begin{aligned} \mathcal{Q}(\omega_1 + \xi\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) &\leq (1 - \xi)\mathcal{Q}(\omega_1) \\ &+ \xi\mathcal{Q}(\omega_2) - c\xi(1 - \xi)\|\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)\|^p \end{aligned} \tag{9}$$

for all $\omega_1, \omega_2 \in \mathcal{K}_{\mathcal{F}}, \xi \in [0, 1]$.

We now introduce a new concept of generalized approximately ψ -quasiconvex functions.

Definition 8. Let a function $\mathcal{Q} : \mathcal{K}_{\mathcal{F}} \rightarrow \mathbb{R}$ is said to be approximately generalized ψ -quasiconvex, if

$$\begin{aligned} \mathcal{Q}(\omega_1 + \xi\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) &\leq \max\{\mathcal{Q}(\omega_1), \mathcal{Q}(\omega_2)\} \\ &+ \mathbb{D}(\omega_1, \omega_2) \end{aligned} \tag{10}$$

for all $\omega_1, \omega_2 \in \mathcal{K}_{\mathcal{F}}, \xi \in [0, 1]$.

We now mention some notable cases of Definition 8 as follows:

(I) For some $c \geq 0, p > 2$ with $\hbar(\xi) = \xi$ and taking $\mathbb{D}(\omega_1, \omega_2) = -c\{\xi^p(1 - \xi) + (1 - \xi)^p\xi\}\|\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)\|^p$, then we have a new concept of HOS generalized ψ -quasiconvex function.

Definition 9. Let a function $\mathcal{Q} : \mathcal{K}_{\mathcal{F}} \rightarrow \mathbb{R}$ is said to HOS generalized ψ -quasi-convex having $c \geq 0$, if

$$\begin{aligned} \mathcal{Q}(\omega_1 + \xi\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) &\leq \max\{\mathcal{Q}(\omega_1), \mathcal{Q}(\omega_2)\} \\ &- c\{\xi^p(1 - \xi) + (1 - \xi)^p\xi\}\|\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)\|^p \end{aligned} \tag{11}$$

for all $\omega_1, \omega_2 \in \mathcal{K}_{\mathcal{F}}, \xi \in [0, 1]$.

(II) For some $c \geq 0, p = 2$ and taking $\mathbb{D}(\omega_1, \omega_2) = -c\{\xi^p(1 - \xi) + (1 - \xi)^p\xi\}\|\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)\|^p$, then we have a new concept of strongly generalized ψ -quasiconvex function.

Definition 10. Let a function $\mathcal{Q} : \mathcal{K}_{\mathcal{F}} \rightarrow \mathbb{R}$ is said to be strongly generalized ψ -quasiconvex having $c \geq 0$, if

$$\begin{aligned} \mathcal{Q}(\omega_1 + \xi\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) &\leq \max\{\mathcal{Q}(\omega_1), \mathcal{Q}(\omega_2)\} \\ &- c\xi(1 - \xi)\|\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)\|^2 \end{aligned} \tag{12}$$

for all $\omega_1, \omega_2 \in \mathcal{K}_{\mathcal{F}}, \xi \in [0, 1]$.

We end this section by presenting a notable fractional integral operators in the literature.

Definition 11. Let $\mathcal{Q} \in L_1[\omega_1, \omega_2]$, then the left- and right-sided Riemann–Liouville fractional integrals $\mathcal{J}_{\omega_1^+}^\delta$ and $\mathcal{J}_{\omega_2^-}^\delta$ are defined as

$$\begin{aligned} \mathcal{J}_{\omega_1^+}^\delta \mathcal{Q}(z) &= \frac{1}{\Gamma(\delta)} \int_{\omega_1}^z (z - \xi)^{\delta-1} \mathcal{Q}(\xi) d\xi, \\ &z < \omega_1, \delta > 0 \end{aligned}$$

and

$$\begin{aligned} \mathcal{J}_{\omega_2^-}^\delta \mathcal{Q}(z) &= \frac{1}{\Gamma(\delta)} \int_z^{\omega_2} (\xi - z)^{\delta-1} \mathcal{Q}(\xi) d\xi, \\ &z > \omega_2, \delta > 0, \end{aligned}$$

respectively, $\Gamma(z) := \int_0^\infty \xi^{z-1} e^{-\xi} d\xi, \Re(z) > 0$ is the Gamma function.

Some of our computations need incomplete beta and hypergeometric functions, which are, respectively, stated as

$$\mathbb{B}_z(\omega_1, \omega_2) = \int_0^z \xi^{\omega_1-1} (1 - \xi)^{\omega_2-1} d\xi.$$

The integral representation of the hypergeometric function is

$$\begin{aligned} {}_2\mathfrak{F}_1(\omega_1, \omega_2; c; z) &= \frac{1}{\mathbb{B}(\omega_2, c - \omega_2)} \int_0^1 \xi^{\omega_2-1} \\ &\times (1 - \xi)^{c-\omega_2-1} (1 - z\xi)^{-\omega_1} d\xi \end{aligned}$$

for $|z| < 1, c > \omega_2 > 0$.

3. AN AUXILIARY RESULT FOR GENERALIZED ψ -CONVEX FUNCTIONS

This section deals with the investigation of an integral identity of κ th-order differentiable functions for generalized ψ -convex functions.

Lemma 12. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° (the interior of Ω) such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$ and $\mathcal{Q}^{(\kappa)} \in L_1[\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)]$ (the Lebesgue space). Then the following equality holds:

$$\begin{aligned} &\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q}) \\ &:= \int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right) \\ &\quad \times \mathcal{Q}^{(\kappa)} \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \\ &\quad + \int_0^1 \left(\frac{\xi^{\delta+\kappa-1} - 2(1-\xi)^{\delta+\kappa-1}}{3} \right) \\ &\quad \times \mathcal{Q}^{(\kappa)} \left(\omega_1 + \frac{n-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi, \end{aligned}$$

where

$$\begin{aligned} &\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q}) \\ &:= \frac{1}{6} \Gamma(\delta + \kappa) \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^{\delta+\kappa} \\ &\quad \times \left[\mathcal{J}_{(\omega_1)^+}^\delta \mathcal{Q} \left(\omega_1 + \frac{1}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right. \\ &\quad \left. + (-1)^\kappa \mathcal{J}_{(\omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^-}^\delta \right. \\ &\quad \left. \times \mathcal{Q} \left(\omega_1 + \frac{n}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right] \\ &\quad + \frac{1}{3} \Gamma(\delta + \kappa) \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^{\delta+\kappa} \\ &\quad \times \left[\mathcal{J}_{(\omega_1 + \frac{1}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^-}^\delta \mathcal{Q}(\omega_1) \right. \\ &\quad \left. + (-1)^\kappa \mathcal{J}_{(\omega_1 + \frac{n}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^+}^\delta \right] \end{aligned}$$

$$\begin{aligned} &\times \mathcal{Q}(\omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))] \\ &- \frac{1}{6} \sum_{q=1}^\kappa \frac{\Gamma(\delta + \kappa)}{\Gamma(\delta + \kappa - q + 1)} \\ &\quad \times \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^q \\ &\quad \times \left[(-1)^{q-1} \mathcal{Q}(\omega_1) + 2(-1)^{q-1} \mathcal{Q}^{(\kappa-q)} \right. \\ &\quad \times \left(\omega_1 + \frac{1}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \\ &\quad \left. + 2\mathcal{Q}^{(\kappa-q)} \left(\omega_1 + \frac{n}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right. \\ &\quad \left. + \mathcal{Q}^{\kappa-q}(\omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) \right]. \end{aligned}$$

Proof. Let

$$\begin{aligned} &\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right) \\ &\quad \times \left[\mathcal{Q}^{(\kappa)} \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right. \\ &\quad \left. - \mathcal{Q}^{(\kappa)} \left(\omega_1 + \frac{n-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right] d\xi \\ &= \int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right) \mathcal{Q}^{(\kappa)} \\ &\quad \times \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \\ &\quad - \int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right) \mathcal{Q}^{(\kappa)} \\ &\quad \times \left(\omega_1 + \frac{n-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \\ &= I_1 - I_2. \end{aligned}$$

Now, integrating by parts, we have

$$\begin{aligned} &\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right) \mathcal{Q}^{(\kappa)} \\ &\quad \times \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \\ &= \frac{2}{3} \int_0^1 (1-\xi)^\delta \mathcal{Q}^{(\kappa)} \\ &\quad \times \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \end{aligned}$$

$$\begin{aligned}
 & + \frac{1}{3} \int_0^1 \xi^\delta \mathcal{Q}^{(\kappa)} \\
 & \times \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \\
 = & \frac{2}{3} \left\{ \left[-\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} (1-\xi)^{\delta+\kappa-1} \mathcal{Q}^{(\kappa-1)} \right. \right. \\
 & \times \left. \left. \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right] \right|_0^1 \\
 & - \frac{(n+1)(\delta+\kappa-1)}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \int_0^1 (1-\xi)^{\delta+\kappa-2} \mathcal{Q}^{(\kappa-1)} \\
 & \times \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \Big\} \\
 & - \frac{1}{3} \left\{ \left[-\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \xi^{\delta+\kappa-1} \mathcal{Q}^{(\kappa-1)} \right. \right. \\
 & \times \left. \left. \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right] \right|_0^1 \\
 & + \frac{(n+1)(\delta+\kappa-1)}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \int_0^1 \xi^{\delta+\kappa-2} \mathcal{Q}^{(\kappa-1)} \\
 & \times \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \Big\}. \\
 & \times \left(\omega_1 + \frac{1}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \\
 & + \frac{(n+1)^2(\delta+\kappa-1)}{(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^2} \mathcal{Q}^{(\kappa-2)} \\
 & \times \left(\omega_1 + \frac{1}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \\
 & - \frac{(n+1)^2(\delta+\kappa-1)(\delta+\kappa-2)}{(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^2} \\
 & \times \int_0^1 (1-\xi)^{\delta+\kappa-3} \\
 & \times \mathcal{Q}^{(\kappa-2)} \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \Big\}.
 \end{aligned}$$

Further, integrating by parts successively upto κ -times, we obtain

$$\begin{aligned}
 I_1 = & \frac{2}{3} \left\{ \sum_{q=1}^{\kappa} \frac{(-1)^{q-1}}{\delta+\kappa} \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^q \right. \\
 & \times \mathcal{Q}^{(\kappa-q)} \left(\omega_1 + \frac{1}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \\
 & \times \prod_{\theta=0}^{q-1} (\delta+\kappa-\theta) \\
 & + \frac{(-1)^\kappa}{\delta+\kappa} \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^\kappa \prod_{\theta=0}^{\kappa} (\delta+\kappa-q) \\
 & \times \int_0^1 (1-\xi)^{\delta-1} \mathcal{Q} \\
 & \times \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \Big\} \\
 & - \frac{1}{3} \left\{ \sum_{q=1}^{\kappa} \frac{(-1)^{q-2}}{\delta+\kappa} \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^q \right. \\
 & \times \mathcal{Q}^{(\kappa-q)}(\omega_1) \prod_{\theta=0}^{q-1} (\delta+\kappa-\theta) \\
 & + \frac{(-1)^{\kappa+1}}{\delta+\kappa} \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^\kappa \\
 & \times \prod_{\theta=0}^{\kappa} (\delta+\kappa-q) \int_0^1 (1-\xi)^{\delta-1} \mathcal{Q}
 \end{aligned}$$

Again, through the integration by parts, we obtain

$$\begin{aligned}
 I_1 = & \frac{2}{3} \left\{ \frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \mathcal{Q}^{(\kappa-1)} \right. \\
 & \times \left(\omega_1 + \frac{1}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \\
 & - \frac{(n+1)^2(\delta+\kappa-1)}{(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^2} \mathcal{Q}^{(\kappa-2)} \\
 & \times \left(\omega_1 + \frac{1}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \\
 & + \frac{(n+1)^2(\delta+\kappa-1)(\delta+\kappa-2)}{(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^2} \\
 & \times \int_0^1 (1-\xi)^{\delta+\kappa-3} \\
 & \times \mathcal{Q}^{(\kappa-2)} \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \Big\} \\
 & - \frac{1}{3} \left\{ -\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \mathcal{Q}^{(\kappa-1)} \right.
 \end{aligned}$$

$$\begin{aligned}
 & \times \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \Big\} \\
 = & \frac{2}{3} \left\{ \sum_{q=1}^{\kappa} \frac{(-1)^{q-1}}{\delta + \kappa} \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^q \right. \\
 & \times \mathcal{Q}^{(\kappa-q)} \left(\omega_1 + \frac{1}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \\
 & \times \prod_{\theta=0}^{q-1} (\delta + \kappa - \theta) \\
 & + \frac{(-1)^\kappa \Gamma(\delta + \kappa)}{\Gamma(\delta)} \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^\kappa \\
 & \times \prod_{\theta=0}^{\kappa} (\delta + \kappa - q) \int_0^1 (1-\xi)^{\delta-1} \mathcal{Q} \\
 & \times \left. \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \right\} \\
 - & \frac{1}{3} \left\{ \sum_{q=1}^{\kappa} \frac{(-1)^{q-2}}{\delta + \kappa} \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^q \right. \\
 & \times \mathcal{Q}^{(\kappa-q)}(\omega_1) \prod_{\theta=0}^{q-1} (\delta + \kappa - \theta) \\
 & + \frac{(-1)^{\kappa+1} \Gamma(\delta + \kappa)}{\Gamma(\delta)} \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^\kappa \\
 & \times \prod_{\theta=0}^{\kappa} (\delta + \kappa - q) \int_0^1 (1-\xi)^{\delta-1} \mathcal{Q} \\
 & \times \left. \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \right\} \\
 = & \frac{2}{3} \left\{ \sum_{q=1}^{\kappa} (-1)^{q-1} \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^q \right. \\
 & \times \frac{\Gamma(\delta + \kappa)}{\Gamma(\delta + \kappa - q + 1)} \\
 & \times \mathcal{Q}^{(\kappa-q)} \left(\omega_1 + \frac{1}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \\
 & + (-1)^\kappa \Gamma(\delta + \kappa) \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^{\delta+\kappa} \\
 & \times \left. \mathcal{J}_{\left(\omega_1 + \frac{1}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)\right)^-}^\delta \mathcal{Q}(\omega_1) \right\}
 \end{aligned}$$

$$\begin{aligned}
 & - \frac{1}{3} \left\{ \sum_{q=1}^{\kappa} (-1)^{q-2} \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^q \right. \\
 & \times \frac{\Gamma(\delta + \kappa)}{\Gamma(\delta + \kappa - q + 1)} \mathcal{Q}^{(\kappa-q)}(\omega_1) \\
 & + (-1)^{\kappa+1} \Gamma(\delta + \kappa) \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^{\delta+\kappa} \\
 & \times \left. \mathcal{J}_{\omega_1^+}^\delta \mathcal{Q} \left(\omega_1 + \frac{1}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right\}.
 \end{aligned}$$

Analogously, we have

$$\begin{aligned}
 I_2 = & \int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right) \mathcal{Q}^{(\kappa)} \\
 & \times \left(\omega_1 + \frac{n+\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \\
 = & \frac{2}{3} \left\{ - \frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \mathcal{Q}^{(\kappa-1)} \right. \\
 & \times \left(\omega_1 + \frac{n}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \\
 & - (\delta + \kappa - 1) \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^2 \\
 & \times \mathcal{Q}^{(\kappa-2)} \left(\omega_1 + \frac{n}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \\
 & - (\delta + \kappa - 1)(\delta + \kappa - 2) \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^3 \\
 & \times \mathcal{Q}^{(\kappa-3)} \left(\omega_1 + \frac{n}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \\
 & + (\delta + \kappa - 1)(\delta + \kappa - 2)(\delta + \kappa - 3) \\
 & \times \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^3 \int_0^1 (1-\xi)^{\delta+\kappa-4} \mathcal{Q}^{(\kappa-3)} \\
 & \times \left. \left(\omega_1 + \frac{n+\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \right\} \\
 - & \frac{1}{3} \left\{ \frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \mathcal{Q}^{(\kappa-1)} \right. \\
 & \times (\omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) \\
 & + (\delta + \kappa - 1) \left(\frac{n+1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^2 \\
 & \times \left. \mathcal{Q}^{(\kappa-2)}(\omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) \right\}
 \end{aligned}$$

$$\begin{aligned}
 & + (\delta + \kappa - 1)(\delta + \kappa - 2) \left(\frac{n + 1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^3 \\
 & \times \mathcal{Q}^{(\kappa-3)}(\omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) \\
 & - (\delta + \kappa - 1)(\delta + \kappa - 2)(\delta + \kappa - 3) \\
 & \times \left(\frac{n + 1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^3 \int_0^1 \xi^{\delta + \kappa - 4} \mathcal{Q}^{(\kappa-3)} \\
 & \times \left(\omega_1 + \frac{n + \xi}{n + 1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) d\xi \} \\
 & \vdots \\
 & = \frac{2}{3} \left\{ - \sum_{q=1}^{\kappa} \left(\frac{n + 1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^q \right. \\
 & \times \frac{\Gamma(\delta + \kappa)}{\Gamma(\delta + \kappa - q + 1)} \\
 & \times \mathcal{Q}^{(\kappa-q)} \left(\omega_1 + \frac{n}{n + 1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \\
 & + \Gamma(\delta + \kappa) \left(\frac{n + 1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^{\delta + \kappa} \\
 & \times \mathcal{J}_{(\omega_1 + \frac{n}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^+}^\delta \\
 & \times \mathcal{Q}(\omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) \} \\
 & - \frac{1}{3} \left\{ \sum_{q=1}^{\kappa} \left(\frac{n + 1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^q \right. \\
 & \times \frac{\Gamma(\delta + \kappa)}{\Gamma(\delta + \kappa - q + 1)} \\
 & \times \mathcal{Q}^{(\kappa-q)}(\omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) \\
 & + \Gamma(\delta + \kappa) \left(\frac{n + 1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^{\delta + \kappa} \\
 & \times \mathcal{J}_{(\omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^-}^\delta \\
 & \times \mathcal{Q} \left(\omega_1 + \frac{n}{n + 1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \} . \\
 & \times \left(\frac{n + 1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^q \\
 & \times \left[(-1)^{q-1} \mathcal{Q}(\omega_1) + 2(-1)^{q-1} \mathcal{Q}^{(\kappa-q)} \right. \\
 & \times \left(\omega_1 + \frac{1}{n + 1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \\
 & - 2\mathcal{Q}^{(\kappa-q)} \left(\omega_1 + \frac{n}{n + 1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \\
 & \left. - \mathcal{Q}^{\kappa-q}(\omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) \right] \\
 & + \frac{1}{6} \Gamma(\delta + \kappa) \left(\frac{n + 1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^{\delta + \kappa} \\
 & \times \left[\mathcal{J}_{(\omega_1)^+}^\delta + \mathcal{Q} \left(\omega_1 + \frac{1}{n + 1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right. \\
 & + (-1)^\kappa \mathcal{J}_{(\omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^-}^\delta \\
 & \left. \times \mathcal{Q} \left(\omega_1 + \frac{n}{n + 1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right] \\
 & + \frac{1}{3} \Gamma(\delta + \kappa) \left(\frac{n + 1}{\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)} \right)^{\delta + \kappa} \\
 & \times \left[\mathcal{J}_{(\omega_1 + \frac{1}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^-}^\delta \mathcal{Q}(\omega_1) \right. \\
 & \left. + (-1)^\kappa \mathcal{J}_{(\omega_1 + \frac{n}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^+}^\delta \right. \\
 & \left. \times \mathcal{Q}(\omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)) \right]. \tag{13}
 \end{aligned}$$

This completes the proof of Lemma 12. □

4. CERTAIN ESTIMATES FOR κ TH DIFFERENTIABLE FUNCTIONS

We propose some new generalizations of upper bounds for the mapping $\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})$ for approximately generalized (ψ, \hbar) -convex functions, our main consequences are described in the subsequent theorems.

Theorem 13. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|$ is an approximately generalized (ψ, \hbar) -convex

Therefore,

$$\begin{aligned}
 I_1 - I_2 & = \frac{1}{6} \sum_{q=1}^{\kappa} \frac{\Gamma(\delta + \kappa)}{\Gamma(\delta + \kappa - q + 1)}
 \end{aligned}$$

on Ω , then

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq \Phi_1(\delta, \kappa, \xi)[|\mathcal{Q}^{(\kappa)}(\omega_1)| + |\mathcal{Q}^{(\kappa)}(\omega_2)|] \\
 & \quad + \frac{2\mathbb{D}(\omega_1, \omega_2)}{3(\delta + \kappa)},
 \end{aligned}$$

where

$$\begin{aligned}
 \Phi_1(\delta, \kappa, \xi) := & \int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right) \\
 & \times \left(\hbar \left(\frac{1 - \xi}{n + 1} \right) + \hbar \left(\frac{n + \xi}{n + 1} \right) \right) d\xi.
 \end{aligned}$$

Proof. According to Lemma 12, the triangular property and utilizing the fact of approximately generalized (ψ, \hbar) -convex functions, we obtain

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq \int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right) \left| \right. \\
 & \quad \times \mathcal{Q}^{(\kappa)} \left(\omega_1 + \frac{1 - \xi}{n + 1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \left. \right| d\xi \\
 & \quad + \int_0^1 \left(\frac{\xi^{\delta + \kappa - 1} - 2(1 - \xi)^{\delta + \kappa - 1}}{3} \right) \left| \right. \\
 & \quad \times \mathcal{Q}^{(\kappa)} \left(\omega_1 + \frac{n - \xi}{n + 1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \left. \right| d\xi \\
 & \leq \int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right) \\
 & \quad \times \left[\hbar \left(\frac{n + \xi}{n + 1} \right) |\mathcal{Q}^{(\kappa)}(\omega_1)| \right. \\
 & \quad \left. + \hbar \left(\frac{1 - \xi}{n + 1} \right) |\mathcal{Q}^{(\kappa)}(\omega_2)| + \mathbb{D}(\omega_1, \omega_2) \right] d\xi \\
 & \quad + \int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right) \\
 & \quad \times \left[\hbar \left(\frac{1 - \xi}{n + 1} \right) |\mathcal{Q}^{(\kappa)}(\omega_1)| \right. \\
 & \quad \left. + \hbar \left(\frac{n + \xi}{n + 1} \right) |\mathcal{Q}^{(\kappa)}(\omega_2)| + \mathbb{D}(\omega_1, \omega_2) \right] d\xi \\
 & \leq \Phi_1(\delta, \kappa, \xi)[|\mathcal{Q}^{(\kappa)}(\omega_1)| + |\mathcal{Q}^{(\kappa)}(\omega_2)|] \\
 & \quad + \frac{2\mathbb{D}(\omega_1, \omega_2)}{3(\delta + \kappa)}. \tag{14}
 \end{aligned}$$

This completes the proof. \square

Now, we shall state some special cases of Theorem 13.

(I) Letting $\hbar(\xi) = 1$, then we acquire a new result for HOS generalized ψ - P -convex functions.

Corollary 14. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|$ is a HOS generalized ψ - P -convex on Ω , then

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq \frac{2}{3(\delta + \kappa)} \{ [|\mathcal{Q}^{(\kappa)}(\omega_1)| + |\mathcal{Q}^{(\kappa)}(\omega_2)|] \\
 & \quad + \mathbb{D}(\omega_1, \omega_2) \}.
 \end{aligned}$$

(II) Letting $\hbar(\xi) = \xi$, then we acquire a new result for HOS generalized ψ -convex functions.

Corollary 15. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|$ is a HOS generalized ψ -convex on Ω , then

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq \frac{1}{3(\delta + \kappa)} \{ [|\mathcal{Q}^{(\kappa)}(\omega_1)| + |\mathcal{Q}^{(\kappa)}(\omega_2)|] \\
 & \quad + 2\mathbb{D}(\omega_1, \omega_2) \}.
 \end{aligned}$$

(III) Letting $\hbar(\xi) = \xi^s$, then we acquire a new result for Breckner type of HOS generalized ψ - s -convex functions.

Corollary 16. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|$ is a Breckner type of HOS generalized ψ - s -convex on Ω , then

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq \frac{1}{3(n + 1)^s} \left[\frac{2}{\delta + \kappa + s} + 2n_2^s \mathfrak{F}_1 \right. \\
 & \quad \left. \times \left(-s; 1; \delta + \kappa + s + 1; -\frac{1}{n} \right) \right]
 \end{aligned}$$

$$\begin{aligned} & -n_2^s \mathfrak{F}_1 \left(-s; 1; \delta + \kappa + s + 1; -\frac{1}{n} \right) && \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n} \right) \\ & -\mathbb{B}(\delta + \kappa, s + 1) [|\mathcal{Q}^{(\kappa)}(\omega_1)| + |\mathcal{Q}^{(\kappa)}(\omega_2)|] && -n\mathbb{B}(\delta + \kappa, p + 1)_2 \mathfrak{F}_1 \\ & + \frac{2\mathbb{D}(\omega_1, \omega_2)}{3(\delta + \kappa)}. && \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n} \right) \\ & && + 2n_2^p \mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n} \right) \\ & && -n^p \mathbb{B}(\delta + \kappa, 2)_2 \mathfrak{F}_1 \\ & && \times \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n} \right) \}. \end{aligned}$$

(IV) Letting $\hbar(\xi) = \xi^{-s}$, then we acquire a new result for Godunova–Levin type of HOS generalized ψ - s -convex functions.

Corollary 17. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|$ is a Godunova–Levin type of HOS generalized ψ - s -convex on Ω , then

$$\begin{aligned} & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\ & \leq \frac{1}{3(n+1)^{-s}} \left[\frac{2}{(\delta + \kappa - s)} - \mathbb{B}(1 - s, \delta + \kappa) \right. \\ & \quad + 2n_2^{-s} \mathfrak{F}_1 \left(s; 1; \delta + \kappa; -\frac{1}{n} \right) \\ & \quad - n^{-s}(\delta + \kappa)_2 \mathfrak{F}_1 \\ & \quad \times \left(s; \delta + \kappa; \delta + \kappa + 1; -\frac{1}{n} \right) \left. \right] \\ & \quad \times [|\mathcal{Q}^{(\kappa)}(\omega_1)| + |\mathcal{Q}^{(\kappa)}(\omega_2)|] \\ & \quad + \frac{2\mathbb{D}(\omega_1, \omega_2)}{3(\delta + \kappa)}. \end{aligned}$$

(V) Letting $\hbar(\xi) = \xi$ and $\mathbb{D}(\omega_1, \omega_2) = -\frac{c}{(n+1)^{2p}} \{(n+\xi)^p(1-\xi) + (n+\xi)(1-\xi)^p\} (\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p$, then we acquire HOS ψ -quasiconvex function.

Corollary 18. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|$ is a HOS generalized ψ -quasiconvex on Ω , then

$$\begin{aligned} & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\ & \leq \frac{1}{3(\delta + \kappa)} \{|\mathcal{Q}^{(\kappa)}(\omega_1)| + |\mathcal{Q}^{(\kappa)}(\omega_2)|\} \\ & \quad - \frac{2c(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p}{3(n+1)^{2p}} \left\{ 2n(\delta + \kappa + 1)_2 \mathfrak{F}_1 \right. \end{aligned}$$

(VI) Letting $\hbar(\xi) = 1$ and $\mathbb{D}(\omega_1, \omega_2) = -\frac{c}{(n+1)^{2p}} \{(n + \xi)^p(1 - \xi) + (n + \xi)(1 - \xi)^p\} (\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p$, then we acquire HOS ψ - P -convex function.

Corollary 19. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|$ is a HOS generalized ψ - P -convex on Ω , then

$$\begin{aligned} & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\ & \leq \frac{2}{3(\delta + \kappa)} \{|\mathcal{Q}^{(\kappa)}(\omega_1)| + |\mathcal{Q}^{(\kappa)}(\omega_2)|\} \\ & \quad - \frac{2c(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p}{3(n+1)^{2p}} \left\{ 2n(\delta + \kappa + 1)_2 \mathfrak{F}_1 \right. \\ & \quad \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n} \right) \\ & \quad - n\mathbb{B}(\delta + \kappa, p + 1)_2 \mathfrak{F}_1 \\ & \quad \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n} \right) \\ & \quad + 2n_2^p \mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n} \right) \\ & \quad - n^p \mathbb{B}(\delta + \kappa, 2)_2 \mathfrak{F}_1 \\ & \quad \times \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n} \right) \}. \end{aligned}$$

(VII) Letting $\hbar(\xi) = \xi^s$ and $\mathbb{D}(\omega_1, \omega_2) = -\frac{c}{(n+1)^{2p}} \{(n + \xi)^p(1 - \xi) + (n + \xi)(1 - \xi)^p\} (\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p$,

then we acquire a new result for Breakner type of HOS generalized ψ - s -convex functions.

Corollary 20. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|$ is a Breakner type of HOS generalized ψ - s -convex on Ω , then

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq \frac{1}{3(n+1)^s} \left[\frac{2}{\delta + \kappa + s} + 2n_2^s \mathfrak{F}_1 \right. \\
 & \quad \times \left(-s; 1; \delta + \kappa + s + 1; -\frac{1}{n} \right) - n_2^s \mathfrak{F}_1 \\
 & \quad \times \left(-s; 1; \delta + \kappa + s + 1; -\frac{1}{n} \right) \\
 & \quad \left. - \mathbb{B}(\delta + \kappa, s + 1) \right] [|\mathcal{Q}^{(\kappa)}(\omega_1)| + |\mathcal{Q}^{(\kappa)}(\omega_2)|] \\
 & \quad - \frac{2c(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p}{3(n+1)^{2p}} \left\{ 2n(\delta + \kappa + 1)_2 \mathfrak{F}_1 \right. \\
 & \quad \times / \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n} \right) \\
 & \quad - n\mathbb{B}(\delta + \kappa, p + 1)_2 \mathfrak{F}_1 \\
 & \quad \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n} \right) \\
 & \quad + 2n_2^p \mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n} \right) \\
 & \quad - n^p \mathbb{B}(\delta + \kappa, 2)_2 \mathfrak{F}_1 \\
 & \quad \left. \times \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n} \right) \right\}.
 \end{aligned}$$

(VIII) Letting $\hbar(\xi) = \xi^{-s}$ and $\mathbb{D}(\omega_1, \omega_2) = -\frac{c}{(n+1)^{2p}} \{ (n+\xi)^p(1-\xi) + (n+\xi)(1-\xi)^p \} (\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p$, then we acquire a new result for Godunova–Levin-type HOS generalized ψ - s -convex functions.

Corollary 21. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|$ is a Godunova–Levin type of HOS generalized

ψ - s -convex on Ω , then

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq \frac{1}{3(n+1)^{-s}} \left[\frac{2}{\delta + \kappa - s} + 2n_2^{-s} \mathfrak{F}_1 \right. \\
 & \quad \times \left(s; 1; \delta + \kappa - s + 1; -\frac{1}{n} \right) \\
 & \quad - n_2^{-s} \mathfrak{F}_1 \left(s; 1; \delta + \kappa - s + 1; -\frac{1}{n} \right) \\
 & \quad \left. - \mathbb{B}(\delta + \kappa, 1 - s) \right] [|\mathcal{Q}^{(\kappa)}(\omega_1)| + |\mathcal{Q}^{(\kappa)}(\omega_2)|] \\
 & \quad - \frac{2c(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p}{3(n+1)^{2p}} \\
 & \quad \times \{ 2n(\delta + \kappa + 1)_2 \mathfrak{F}_1 \\
 & \quad \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n} \right) \\
 & \quad - n\mathbb{B}(\delta + \kappa, p + 1)_2 \mathfrak{F}_1 \\
 & \quad \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n} \right) \\
 & \quad + 2n_2^p \mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n} \right) \\
 & \quad - n^p \mathbb{B}(\delta + \kappa, 2)_2 \mathfrak{F}_1 \\
 & \quad \left. \times \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n} \right) \right\}.
 \end{aligned}$$

Theorem 22. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|^{q_2}$ is an approximately generalized (ψ, \hbar) -convex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq \left(\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\
 & \quad \times \{ [\Phi_1^{**}(\delta, \kappa, n, \xi) |\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} \\
 & \quad + \Phi_2^{**}(\delta, \kappa, n, \xi) \\
 & \quad \times |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} + \mathbb{D}(\omega_1, \omega_2)]^{\frac{1}{q_2}} \}
 \end{aligned}$$

$$\begin{aligned}
 & + [\Phi_2^{**}(\delta, \kappa, n, \xi) |\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} \\
 & + \Phi_1^{**}(\delta, \kappa, n, \xi) \\
 & \times |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} + \mathbb{D}(\omega_1, \omega_2)]^{\frac{1}{q_2}}, \tag{15}
 \end{aligned}$$

where

$$\Phi_1^{**}(\delta, \kappa, n, \xi) := \int_0^1 \hbar \left(\frac{n + \xi}{n + 1} \right) d\xi \tag{16}$$

and

$$\Phi_2^{**}(\delta, \kappa, n, \xi) := \int_0^1 \hbar \left(\frac{1 - \xi}{n + 1} \right) d\xi. \tag{17}$$

Proof. According to Lemma 12, the well-known Hölder inequality and utilizing the fact of approximately generalized (ψ, \hbar) -convex functions, we obtain

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq \left(\int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\
 & \times \left(\int_0^1 \left| \mathcal{Q}^{(\kappa)} \left(\omega_1 + \frac{1 - \xi}{n + 1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right. \right. \\
 & \times \left. \left. \left| d\xi \right|^{\frac{1}{q_2}} \right)^{\frac{1}{q_2}} \\
 & + \left(\int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\
 & \times \left(\int_0^1 \left| \mathcal{Q}^{(\kappa)} \left(\omega_1 + \frac{n + \xi}{n + 1} \mathcal{F}_{v_1, v_2}^\eta \right. \right. \right. \\
 & \times \left. \left. \left. (\omega_2 - \omega_1) \right) \right|^{\frac{1}{q_2}} d\xi \right)^{\frac{1}{q_2}} \\
 & \leq \left(\int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\
 & \times \left\{ \left[\int_0^1 \left(\hbar \left(\frac{n + \xi}{n + 1} \right) |\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} \right. \right. \right. \\
 & + \hbar \left(\frac{1 - \xi}{n + 1} \right) |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} \\
 & \left. \left. \left. + \mathbb{D}(\omega_1, \omega_2) \right) d\xi \right]^{\frac{1}{q_2}}
 \end{aligned}$$

$$\begin{aligned}
 & + \left[\int_0^1 \left(\hbar \left(\frac{1 - \xi}{n + 1} \right) |\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} \right. \right. \\
 & + \hbar \left(\frac{n + \xi}{n + 1} \right) |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} + \mathbb{D}(\omega_1, \omega_2) \left. \right. \\
 & \left. \left. \times \xi \right]^{\frac{1}{q_2}} \right\} \\
 & \leq \left(\int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\
 & \times \{ [\Phi_1^{**}(\delta, \kappa, n, \xi) |\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} \\
 & + \Phi_2^{**}(\delta, \kappa, n, \xi) |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} \\
 & + \mathbb{D}(\omega_1, \omega_2)]^{\frac{1}{q_2}} \\
 & + [\Phi_2^{**}(\delta, \kappa, n, \xi) |\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} \\
 & + \Phi_1^{**}(\delta, \kappa, n, \xi) |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} \\
 & + \mathbb{D}(\omega_1, \omega_2)]^{\frac{1}{q_2}} \}. \tag{18}
 \end{aligned}$$

This completes the proof. \square

Some new special cases of Theorem 22 can be described as follows:

(I) Letting $\hbar(\xi) = 1$, then we acquire a new result for HOS generalized ψ - P -convex functions.

Corollary 23. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|^{q_2}$ is a HOS generalized (ψ, \hbar) -convex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq 2 \left(\int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\
 & \times [|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} + |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} \\
 & + \mathbb{D}(\omega_1, \omega_2)]^{\frac{1}{q_2}}.
 \end{aligned}$$

(II) Letting $\hbar(\xi) = \xi$, then we acquire a new result for HOS generalized ψ -convex functions.

Corollary 24. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable

mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|^{q_2}$ is a HOS generalized ψ -convex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned} & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\ & \leq \left(\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\ & \quad \times \left\{ \left[\frac{2n+1}{2(n+1)} |\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} + \frac{1}{2(n+1)} \right. \right. \\ & \quad \times \left. \left. |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} + \mathbb{D}(\omega_1, \omega_2) \right]^{\frac{1}{q_2}} \right. \\ & \quad \left. + \left[\frac{1}{2(n+1)} |\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} + \frac{2n+1}{2(n+1)} \right. \right. \\ & \quad \left. \left. \times |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} + \mathbb{D}(\omega_1, \omega_2) \right]^{\frac{1}{q_2}} \right\}. \end{aligned}$$

(III) Letting $\hbar(\xi) = \xi^s$, then we acquire a new result for Breckner type of HOS generalized ψ - s -convex functions.

Corollary 25. For $s \in (0, 1], n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|^{q_2}$ is a Breckner type of HOS generalized ψ - s -convex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned} & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\ & \leq \left(\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\ & \quad \times \left\{ \left[\frac{n^{s+1} |\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} + |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}}{(n+1)^s(s+1)} \right. \right. \\ & \quad \left. \left. + \mathbb{D}(\omega_1, \omega_2) \right]^{\frac{1}{q_2}} \right. \\ & \quad \left. + \left[\frac{n^{s+1} |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} + |\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}}{(n+1)^s(s+1)} \right. \right. \\ & \quad \left. \left. + \mathbb{D}(\omega_1, \omega_2) \right]^{\frac{1}{q_2}} \right\}. \end{aligned}$$

(IV) Letting $\hbar(\xi) = \xi^s$, then we acquire a new result for Godunova-Levin type of HOS generalized ψ - s -convex functions.

Corollary 26. For $s \in (0, 1], n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|^{q_2}$ is a Godunova-Levin type of HOS generalized ψ - s -convex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned} & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\ & \leq \left(\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\ & \quad \times \left\{ \left[\frac{n^{1-s} |\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} + |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}}{(n+1)^{-s}(1-s)} \right. \right. \\ & \quad \left. \left. + \mathbb{D}(\omega_1, \omega_2) \right]^{\frac{1}{q_2}} \right. \\ & \quad \left. + \left[\frac{n^{1-s} |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} + |\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}}{(n+1)^{-s}(1-s)} \right. \right. \\ & \quad \left. \left. + \mathbb{D}(\omega_1, \omega_2) \right]^{\frac{1}{q_2}} \right\}. \end{aligned}$$

(V) Letting $\hbar(\xi) = 1$ and $\mathbb{D}(\omega_1, \omega_2) = -\frac{c}{(n+1)^{2p}} \left\{ (n + \xi)^p(1 - \xi) + (n + \xi)(1 - \xi)^p \right\} (\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p$, then we acquire a new result for HOS generalized ψ - P -convex functions.

Corollary 27. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|^{q_2}$ is a HOS of generalized (ψ, \hbar) -convex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned} & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\ & \leq 2 \left(\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \end{aligned}$$

$$\begin{aligned} & \times \left[|Q^{(\kappa)}(\omega_1)|^{q_2} + |Q^{(\kappa)}(\omega_2)|^{q_2} \right. \\ & - \frac{2c(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p}{3(n+1)^{2p}} \{2n(\delta + \kappa + 1)_2 \mathfrak{F}_1 \\ & \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ & - n\mathbb{B}(\delta + \kappa, p + 1)_2 \mathfrak{F}_1 \\ & \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ & + 2n_2^p \mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n}\right) \\ & - n^p \mathbb{B}(\delta + \kappa, 2)_2 \mathfrak{F}_1 \\ & \left. \times \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n}\right) \right]^{1/q_2}. \end{aligned}$$

(VI) Letting $\hbar(\xi) = \xi$ and $\mathbb{D}(\omega_1, \omega_2) = -\frac{c}{(n+1)^{2p}} \left\{ (n + \xi)^p (1 - \xi) + (n + \xi)(1 - \xi)^p \right\} (\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p$, then we acquire a new result for HOS generalized ψ -convex functions.

Corollary 28. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $Q : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|Q^{(\kappa)}|^{q_2}$ is a HOS generalized ψ -convex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned} & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(Q)| \\ & \leq \left(\int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right)^{q_1} d\xi \right)^{1/q_1} \\ & \times \left\{ \left[\frac{2n+1}{2(n+1)} |Q^{(\kappa)}(\omega_1)|^{q_2} \right. \right. \\ & + \frac{1}{2(n+1)} |Q^{(\kappa)}(\omega_2)|^{q_2} \\ & - \frac{2c(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p}{3(n+1)^{2p}} \\ & \times \{2n(\delta + \kappa + 1)_2 \mathfrak{F}_1 \\ & \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ & - n\mathbb{B}(\delta + \kappa, p + 1)_2 \mathfrak{F}_1 \end{aligned}$$

$$\begin{aligned} & \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ & + 2n_2^p \mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n}\right) \\ & - n^p \mathbb{B}(\delta + \kappa, 2) \\ & \left. \times {}_2 \mathfrak{F}_1 \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n}\right) \right\}^{1/q_2} \\ & + \left[\frac{1}{2(n+1)} |Q^{(\kappa)}(\omega_1)|^{q_2} \right. \\ & + \frac{2n+1}{2(n+1)} |Q^{(\kappa)}(\omega_2)|^{q_2} \\ & - \frac{2c(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p}{3(n+1)^{2p}} \{2n(\delta + \kappa + 1)_2 \mathfrak{F}_1 \\ & \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ & - n\mathbb{B}(\delta + \kappa, p + 1)_2 \mathfrak{F}_1 \\ & \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ & + 2n_2^p \mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n}\right) \\ & - n^p \mathbb{B}(\delta + \kappa, 2)_2 \mathfrak{F}_1 \\ & \left. \times \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n}\right) \right\}^{1/q_2} \}. \end{aligned}$$

(VII) Letting $\hbar(\xi) = \xi^s$ and $\mathbb{D}(\omega_1, \omega_2) = -\frac{c}{(n+1)^{2p}} \left\{ (n + \xi)^p (1 - \xi) + (n + \xi)(1 - \xi)^p \right\} (\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p$, then we acquire a new result for Breckner type of HOS generalized ψ -s-convex functions.

Corollary 29. For $s \in (0, 1], n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $Q : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|Q^{(\kappa)}|^{q_2}$ is a Breckner type of HOS generalized ψ -s-convex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned} & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(Q)| \\ & \leq \left(\int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right)^{q_1} d\xi \right)^{1/q_1} \end{aligned}$$

$$\begin{aligned} & \times \left\{ \left[\frac{n^{s+1}|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} + |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}}{(n+1)^s(s+1)} \right. \right. \\ & - \frac{2c(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p}{3(n+1)^{2p}} \\ & \times \{2n(\delta + \kappa + 1)_2\mathfrak{F}_1 \\ & \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ & - n\mathbb{B}(\delta + \kappa, p + 1)_2\mathfrak{F}_1 \\ & \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ & + 2n_2^p\mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n}\right) \\ & - n^p\mathbb{B}(\delta + \kappa, 2)_2\mathfrak{F}_1 \\ & \left. \left. \times \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n}\right) \right\} \right]^{\frac{1}{q_2}} \\ & + \left[\frac{n^{s+1}|\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} + |\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}}{(n+1)^s(s+1)} \right. \\ & - \frac{2c(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p}{3(n+1)^{2p}} \\ & \times \{2n(\delta + \kappa + 1)_2\mathfrak{F}_1 \\ & \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ & - n\mathbb{B}(\delta + \kappa, p + 1)_2\mathfrak{F}_1 \\ & \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ & + 2n_2^p\mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n}\right) \\ & - n^p\mathbb{B}(\delta + \kappa, 2)_2\mathfrak{F}_1 \\ & \left. \left. \times \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n}\right) \right\} \right]^{\frac{1}{q_2}} \Bigg\}. \end{aligned}$$

(VIII) Letting $h(\xi) = \xi^{-s}$ and $\mathbb{D}(\omega_1, \omega_2) = -\frac{c}{(n+1)^{2p}} \left\{ (n + \xi)^p(1 - \xi) + (n + \xi)(1 - \xi)^p \right\} (\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p$, then we acquire a new result for Godunova–Levin-type of HOS generalized ψ - s -convex functions.

Corollary 30. For $s \in (0, 1], n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded

sequence of real numbers. Also, assume that a κ th order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|^{q_2}$ is a Godunova–Levin type of HOS generalized ψ - s -convex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned} & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\ & \leq \left(\int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\ & \times \left\{ \left[\frac{n^{1-s}|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} + |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}}{(n+1)^{-s}(1-s)} \right. \right. \\ & - \frac{2c(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p}{3(n+1)^{2p}} \\ & \times \{2n(\delta + \kappa + 1)_2\mathfrak{F}_1 \\ & \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ & - n\mathbb{B}(\delta + \kappa, p + 1)_2\mathfrak{F}_1 \\ & \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ & + 2n_2^p\mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n}\right) \\ & - n^p\mathbb{B}(\delta + \kappa, 2)_2\mathfrak{F}_1 \\ & \left. \left. \times \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n}\right) \right\} \right]^{\frac{1}{q_2}} \\ & + \left[\frac{n^{1-s}|\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} + |\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}}{(n+1)^{-s}(1-s)} \right. \\ & - \frac{2c(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p}{3(n+1)^{2p}} \\ & \times \{2n(\delta + \kappa + 1)_2\mathfrak{F}_1 \\ & \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ & - n\mathbb{B}(\delta + \kappa, p + 1)_2\mathfrak{F}_1 \\ & \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ & + 2n_2^p\mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n}\right) \\ & - n^p\mathbb{B}(\delta + \kappa, 2)_2\mathfrak{F}_1 \\ & \left. \left. \times \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n}\right) \right\} \right]^{\frac{1}{q_2}} \Bigg\}. \end{aligned}$$

Theorem 31. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|^{q_2}$ is a HOS generalized (ψ, \hbar) -convex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned} & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\ & \leq \left(\frac{1}{3(\delta + \kappa)}\right)^{1-\frac{1}{q_2}} \{[\Phi_1^*(\delta, \kappa, n, \xi)|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} \\ & \quad + \Phi_2^*(\delta, \kappa, n, \xi)|\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} \\ & \quad + \mathbb{D}(\omega_1, \omega_2)]^{\frac{1}{q_2}} \\ & \quad + [\Phi_2^*(\delta, \kappa, n, \xi)|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} \\ & \quad + \Phi_1^*(\delta, \kappa, n, \xi)|\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} \\ & \quad + \mathbb{D}(\omega_1, \omega_2)]^{\frac{1}{q_2}}\}, \end{aligned} \tag{19}$$

where

$$\begin{aligned} \Phi_1^*(\delta, \kappa, n, \xi) & := \int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3}\right) \\ & \quad \times \hbar\left(\frac{n+\xi}{n+1}\right) d\xi \end{aligned} \tag{20}$$

and

$$\begin{aligned} \Phi_2^*(\delta, \kappa, n, \xi) & := \int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3}\right) \\ & \quad \times \hbar\left(\frac{1-\xi}{n+1}\right) d\xi. \end{aligned} \tag{21}$$

Proof. According to Lemma 12, the well-known power-mean inequality and utilizing the fact of approximately generalized (ψ, \hbar) -convex functions, we obtain

$$\begin{aligned} & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\ & \leq \left(\int_0^1 \frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} d\xi\right)^{1-\frac{1}{q_2}} \\ & \quad \times \left(\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3}\right) \right. \\ & \quad \times \left. \left|\mathcal{Q}^{(\kappa)}\left(\omega_1 + \frac{1-\xi}{n+1}\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)\right.\right.\right. \\ & \quad \left.\left.\left. \times (\omega_2 - \omega_1)\right)\right|^{q_2} d\xi\right)^{\frac{1}{q_2}} \end{aligned}$$

$$\begin{aligned} & + \left(\int_0^1 \frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} d\xi\right)^{1-\frac{1}{q_2}} \\ & \quad \times \left(\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3}\right) \right. \\ & \quad \times \left. \left|\mathcal{Q}^{(\kappa)}\left(\omega_1 + \frac{n+\xi}{n+1}\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)\right)\right|^{q_2} d\xi\right)^{\frac{1}{q_2}} \\ & \leq \left(\frac{1}{3(\delta + \kappa)}\right)^{1-\frac{1}{q_2}} \\ & \quad \times \left\{ \left[\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3}\right) \right. \right. \\ & \quad \times \left. \left(\hbar\left(\frac{n+\xi}{n+1}\right)|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} \right. \right. \\ & \quad \left. \left. + \hbar\left(\frac{1-\xi}{n+1}\right)|\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} \right. \right. \\ & \quad \left. \left. + \mathbb{D}(\omega_1, \omega_2)\right) d\xi\right]^{\frac{1}{q_2}} \\ & \quad + \left[\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3}\right) \right. \\ & \quad \times \left. \left(\hbar\left(\frac{1-\xi}{n+1}\right)|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} + \hbar\left(\frac{n+\xi}{n+1}\right) \right. \right. \\ & \quad \left. \left. \times |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} + \mathbb{D}(\omega_1, \omega_2)\right) d\xi\right]^{\frac{1}{q_2}} \left. \right\} \\ & \leq \left(\frac{1}{3(\delta + \kappa)}\right)^{1-\frac{1}{q_2}} \{[\Phi_1^*(\delta, \kappa, n, \xi)|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} \\ & \quad + \Phi_2^*(\delta, \kappa, n, \xi)|\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} + \mathbb{D}(\omega_1, \omega_2)]^{\frac{1}{q_2}} \\ & \quad + [\Phi_2^*(\delta, \kappa, n, \xi)|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2} + \Phi_1^*(\delta, \kappa, n, \xi) \\ & \quad \times |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2} + \mathbb{D}(\omega_1, \omega_2)]^{\frac{1}{q_2}}\}. \end{aligned} \tag{22}$$

This completes the proof. \square

Remark 32. By applying the similar arguments as we did for Theorems 13 and 22, we can find several special cases for Theorem 31 by considering the approximately generalized (ψ, \hbar) -convex functions with the appropriate and exceptional selection of function \hbar .

5. NEW GENERALIZATIONS OF APPROXIMATELY GENERALIZED ψ -QUASICONVEX FUNCTIONS HAVING κ th ORDER DIFFERENTIABILITY

We propose some new generalizations of upper bounds for the mapping $\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})$ for approximately generalized ψ -quasiconvex functions by considering Definition 8 and Lemma 12.

Theorem 33. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|$ is an approximately generalized ψ -quasiconvex on Ω , then

$$|\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \leq \left(\frac{1}{3(\delta + \kappa)}\right) \{ \max[|\mathcal{Q}^{(\kappa)}(\omega_1)|, |\mathcal{Q}^{(\kappa)}(\omega_2)|] + \mathbb{D}(\omega_1, \omega_2) \}.$$

Proof. According to Lemma 12, the triangular property and utilizing the fact of approximately generalized ψ -quasiconvex functions, we obtain

$$\begin{aligned} |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| &\leq \int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right) \\ &\times \left| \mathcal{Q}^{(\kappa)} \left(\omega_1 + \frac{1 - \xi}{n + 1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right| d\xi \\ &+ \int_0^1 \left(\frac{\xi^{\delta + \kappa - 1} - 2(1 - \xi)^{\delta + \kappa - 1}}{3} \right) \\ &\times \left| \mathcal{Q}^{(\kappa)} \left(\omega_1 + \frac{n + \xi}{n + 1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right| d\xi \\ &\leq \int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right) \\ &\times [\max\{|\mathcal{Q}^{(\kappa)}(\omega_1)|, |\mathcal{Q}^{(\kappa)}(\omega_2)|\}] \\ &+ \mathbb{D}(\omega_1, \omega_2)] d\xi \\ &+ \int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right) \end{aligned}$$

$$\begin{aligned} &\times [\max\{|\mathcal{Q}^{(\kappa)}(\omega_1)|, |\mathcal{Q}^{(\kappa)}(\omega_2)|\}] \\ &+ \mathbb{D}(\omega_1, \omega_2)] d\xi \\ &= \left(\frac{1}{3(\delta + \kappa)}\right) \{ \max[|\mathcal{Q}^{(\kappa)}(\omega_1)|, |\mathcal{Q}^{(\kappa)}(\omega_2)|] \\ &+ \mathbb{D}(\omega_1, \omega_2) \}. \end{aligned} \tag{23}$$

This completes the proof. \square

Some remarkable cases of Theorem 33 can be discussed as follows:

(I) Letting $\mathbb{D}(\omega_1, \omega_2) = -\frac{c}{(n+1)^{2p}} \{ (n + \xi)^p(1 - \xi) + (n + \xi)(1 - \xi)^p \} (\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p$, then we acquire HOS ψ -quasiconvex function.

Corollary 34. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|$ is a HOS generalized ψ -quasiconvex on Ω , then

$$\begin{aligned} |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| &\leq \left(\frac{1}{3(\delta + \kappa)}\right) \{ \max[|\mathcal{Q}^{(\kappa)}(\omega_1)|, |\mathcal{Q}^{(\kappa)}(\omega_2)|] \\ &- \frac{c(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p}{3(n + 1)^{2p}} \{ 2n(\delta + \kappa + 1) \\ &\times {}_2\mathfrak{F}_1 \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n} \right) \\ &- n\mathbb{B}(\delta + \kappa, p + 1) \\ &\times {}_2\mathfrak{F}_1 \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n} \right) \\ &+ 2n_2^p \mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n} \right) \\ &- n^p \mathbb{B}(\delta + \kappa, 2) {}_2\mathfrak{F}_1 \\ &\times \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n} \right) \} \}. \end{aligned}$$

(II) Letting $\mathbb{D}(\omega_1, \omega_2) = -\frac{c}{(n+1)^{2p}} \{ (n + \xi)^p(1 - \xi) + (n + \xi)(1 - \xi)^p \} (\omega_2 - \omega_1)^p$ along with $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) = \omega_2 - \omega_1$, then we acquire HOS quasiconvex function.

Corollary 35. For $n, \kappa \in \mathbb{N}$ with $\delta > 0$. Let a κ th-order differentiable mapping $\mathcal{Q} : \Omega \rightarrow \mathbb{R}$ be defined

on Ω° . If $|\mathcal{Q}^{(\kappa)}|$ is a HOS quasiconvex on Ω , then

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq \left(\frac{1}{3(\delta + \kappa)} \right) \left\{ \max[|\mathcal{Q}^{(\kappa)}(\omega_1)|, |\mathcal{Q}^{(\kappa)}(\omega_2)|] \right. \\
 & \quad - \frac{c(\omega_2 - \omega_1)^p}{3(n+1)^{2p}} \{2n(\delta + \kappa + 1)_2\mathfrak{F}_1 \\
 & \quad \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\
 & \quad - n\mathbb{B}(\delta + \kappa, p + 1)_2\mathfrak{F}_1 \\
 & \quad \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\
 & \quad + 2n_2^p\mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n}\right) \\
 & \quad - n^p\mathbb{B}(\delta + \kappa, 2) \\
 & \quad \left. \times {}_2\mathfrak{F}_1 \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n}\right) \right\}.
 \end{aligned}$$

Theorem 36. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|^{q_2}$ is an approximately generalized ψ -quasiconvex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq 2 \left(\int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\
 & \quad \times \{ \max\{|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}, |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}\} \\
 & \quad + \mathbb{D}(\omega_1, \omega_2) \}^{\frac{1}{q_2}}.
 \end{aligned} \tag{24}$$

Proof. According to Lemma 12, the well-known Hölder inequality and utilizing the fact of approximately generalized ψ -quasiconvex functions, we obtain

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq \left(\int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}}
 \end{aligned}$$

$$\begin{aligned}
 & \times \left(\int_0^1 \left| \mathcal{Q}^{(\kappa)} \left(\omega_1 + \frac{1 - \xi}{n + 1} \mathcal{F}_{v_1, v_2}^\eta \right. \right. \right. \\
 & \quad \left. \left. \left. \times (\omega_2 - \omega_1) \right) \right|^{q_2} d\xi \right)^{\frac{1}{q_2}} \\
 & + \left(\int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\
 & \times \left(\int_0^1 \left| \mathcal{Q}^{(\kappa)} \left(\omega_1 + \frac{n + \xi}{n + 1} \mathcal{F}_{v_1, v_2}^\eta \right. \right. \right. \\
 & \quad \left. \left. \left. \times (\omega_2 - \omega_1) \right) \right|^{q_2} d\xi \right)^{\frac{1}{q_2}} \\
 & \leq \left(\int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\
 & \times \left\{ \left[\int_0^1 (\max\{|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}, |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}\} \right. \right. \right. \\
 & \quad \left. \left. \left. + \mathbb{D}(\omega_1, \omega_2) \right) d\xi \right]^{\frac{1}{q_2}} + \left[\int_0^1 (\max\{|\mathcal{Q}^{(\kappa)} \right. \right. \right. \\
 & \quad \left. \left. \left. \times (\omega_1)|^{q_2}, |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}\} + \mathbb{D}(\omega_1, \omega_2) \right) d\xi \right]^{\frac{1}{q_2}} \right\} \\
 & \leq 2 \left(\int_0^1 \left(\frac{2(1 - \xi)^{\delta + \kappa - 1} - \xi^{\delta + \kappa - 1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\
 & \times \{ \max\{|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}, |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}\} \\
 & \quad + \mathbb{D}(\omega_1, \omega_2) \}^{\frac{1}{q_2}}.
 \end{aligned} \tag{25}$$

This completes the proof. \square

Some remarkable cases of Theorem 36 can be discussed as follows:

(I) Letting $\mathbb{D}(\omega_1, \omega_2) = -\frac{c}{(n+1)^{2p}} \left\{ (n + \xi)^p(1 - \xi) + (n + \xi)(1 - \xi)^p \right\} (\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p$, then we acquire HOS ψ -quasiconvex function.

Corollary 37. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|^{q_2}$ is a HOS generalized ψ -quasiconvex on Ω

for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq 2 \left(\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\
 & \quad \times [\max\{|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}, |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}\} \\
 & \quad - \frac{c(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p}{3(n+1)^{2p}} \{2n(\delta + \kappa + 1)_2 \mathfrak{F}_1 \\
 & \quad \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\
 & \quad - n\mathbb{B}(\delta + \kappa, p + 1)_2 \mathfrak{F}_1 \\
 & \quad \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\
 & \quad + 2n_2^p \mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n}\right) \\
 & \quad - n^p \mathbb{B}(\delta + \kappa, 2)_2 \mathfrak{F}_1 \\
 & \quad \times \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n}\right) \left. \right\}]^{\frac{1}{q_2}}.
 \end{aligned}$$

(II) Letting $\mathbb{D}(\omega_1, \omega_2) = -\frac{c}{(n+1)^{2p}} \{(n + \xi)^p(1 - \xi) + (n + \xi)(1 - \xi)^p\}(\omega_2 - \omega_1)^p$ along with $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) = \omega_2 - \omega_1$, then we acquire HOS quasiconvex function.

Corollary 38. For $n, \kappa \in \mathbb{N}$ with $\delta > 0$. Let a κ th-order differentiable mapping $\mathcal{Q} : \Omega \rightarrow \mathbb{R}$ be defined on Ω° . If $|\mathcal{Q}^{(\kappa)}|^{q_2}$ is a HOS quasiconvex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq 2 \left(\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right)^{q_1} d\xi \right)^{\frac{1}{q_1}} \\
 & \quad \times [\max\{|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}, |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}\} \\
 & \quad - \frac{c(\omega_2 - \omega_1)^p}{3(n+1)^{2p}} \{2n(\delta + \kappa + 1)_2 \mathfrak{F}_1 \\
 & \quad \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\
 & \quad - n\mathbb{B}(\delta + \kappa, p + 1)_2 \mathfrak{F}_1 \\
 & \quad \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\
 & \quad \times \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n}\right) \left. \right\}]^{\frac{1}{q_2}}.
 \end{aligned}$$

$$\begin{aligned}
 & + 2n_2^p \mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n}\right) \\
 & - n^p \mathbb{B}(\delta + \kappa, 2)_2 \mathfrak{F}_1 \\
 & \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n}\right) \left. \right\}]^{\frac{1}{q_2}}.
 \end{aligned}$$

Theorem 39. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|^{q_2}$ is an approximately generalized ψ -quasiconvex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq 2 \left(\frac{1}{3(\delta + \kappa)} \right) [\max\{|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}, \\
 & \quad \times |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}\} + \mathbb{D}(\omega_1, \omega_2)]^{\frac{1}{q_2}}. \quad (26)
 \end{aligned}$$

Proof. According to Lemma 12, the well-known power-mean inequality and utilizing the fact of approximately generalized (ψ, \hbar) -quasiconvex of $|\mathcal{Q}^{(\kappa)}|^{q_2}$, we obtain

$$\begin{aligned}
 & |\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\
 & \leq \left(\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right) d\xi \right)^{1 - \frac{1}{q_2}} \\
 & \quad \times \left(\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right) \right. \\
 & \quad \times \left| \mathcal{Q}^{(\kappa)} \left(\omega_1 + \frac{1-\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right|^{q_2} \\
 & \quad \times d\xi \left. \right)^{\frac{1}{q_2}} \\
 & \quad + \left(\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right) \right. \\
 & \quad \times d\xi \left. \right)^{1 - \frac{1}{q_2}} \\
 & \quad \times \left(\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3} \right) \right) \left| \mathcal{Q}^{(\kappa)} \right. \\
 & \quad \times \left. \left(\omega_1 + \frac{n+\xi}{n+1} \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) \right) \right|^{q_2} d\xi \left. \right)^{\frac{1}{q_2}}
 \end{aligned}$$

$$\begin{aligned} &\leq \left(\frac{1}{3(\delta + \kappa)}\right)^{1-\frac{1}{q_2}} \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ &\quad \times \left[\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3}\right) - n\mathbb{B}(\delta + \kappa, p + 1)_2\mathfrak{F}_1 \right. \\ &\quad \times (\max\{|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}, |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}\} \\ &\quad \left. + \mathbb{D}(\omega_1, \omega_2))d\xi\right]^{\frac{1}{q_2}} \\ &\quad + \left(\frac{1}{3(\delta + \kappa)}\right)^{1-\frac{1}{q_2}} \\ &\quad \times \left[\int_0^1 \left(\frac{2(1-\xi)^{\delta+\kappa-1} - \xi^{\delta+\kappa-1}}{3}\right) \right. \\ &\quad \times (\max\{|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}, |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}\} \\ &\quad \left. + \mathbb{D}(\omega_1, \omega_2))d\xi\right]^{\frac{1}{q_2}} \\ &= 2 \left(\frac{1}{3(\delta + \kappa)}\right) \\ &\quad \times [\max\{|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}, |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}\} \\ &\quad \left. + \mathbb{D}(\omega_1, \omega_2)\right]^{\frac{1}{q_2}}. \end{aligned} \tag{28}$$

(II) Letting $\mathbb{D}(\omega_1, \omega_2) = -\frac{c}{(n+1)^{2p}} \left\{ (n + \xi)^p(1 - \xi) + (n + \xi)(1 - \xi)^p \right\} (\omega_2 - \omega_1)^p$ along with $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) = \omega_2 - \omega_1$, then we acquire HOS quasiconvex function.

Corollary 41. For $n, \kappa \in \mathbb{N}$ with $\delta > 0$. Let a κ th order differentiable mapping $\mathcal{Q} : \Omega \rightarrow \mathbb{R}$ be defined on Ω° . If $|\mathcal{Q}^{(\kappa)}|^{q_2}$ is a HOS quasiconvex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned} &|\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\ &\leq 2 \left(\frac{1}{3(\delta + \kappa)}\right)^{1-\frac{1}{q_2}} \left[\frac{1}{3(\delta + \kappa)} \right. \\ &\quad \times \max\{|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}, |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}\} \\ &\quad \left. - \frac{c(\omega_2 - \omega_1)^p}{3(n + 1)^{2p}} \{2n(\delta + \kappa + 1)_2\mathfrak{F}_1 \right. \\ &\quad \times \left(-1; 1, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ &\quad \left. - n\mathbb{B}(\delta + \kappa, p + 1)_2\mathfrak{F}_1 \right. \\ &\quad \times \left(-1; \delta + \kappa, \delta + \kappa + p + 1, -\frac{1}{n}\right) \\ &\quad \left. + 2n_2^p\mathfrak{F}_1 \left(-p; 1, \delta + \kappa + 2, -\frac{1}{n}\right) \right. \\ &\quad \left. - n^p\mathbb{B}(\delta + \kappa, 2) \right. \\ &\quad \left. \times 2\mathfrak{F}_1 \left(-p; \delta + \kappa, \delta + \kappa + 2, -\frac{1}{n}\right) \right]^{\frac{1}{q_2}}. \end{aligned} \tag{29}$$

This completes the proof. \square

Some remarkable cases of Theorem 39 can be discussed as follows:

(I) Letting $\mathbb{D}(\omega_1, \omega_2) = -\frac{c}{(n+1)^{2p}} \left\{ (n + \xi)^p(1 - \xi) + (n + \xi)(1 - \xi)^p \right\} (\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p$, then we acquire HOS ψ -quasiconvex function.

Corollary 40. For $n, \kappa \in \mathbb{N}$ with $v_1, v_2, \delta > 0$ and $\eta = \{\eta(m)\}_{m=0}^\infty$ a bounded sequence of real numbers. Also, assume that a κ th-order differentiable mapping $\mathcal{Q} : \Omega = [\omega_1, \omega_1 + \mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1)] \rightarrow \mathbb{R}$ is defined on Ω° such that $\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1) > 0$. If $|\mathcal{Q}^{(\kappa)}|^{q_2}$ is a HOS generalized ψ -quasiconvex on Ω for $q_2 > 1$ with $q_1^{-1} + q_2^{-1} = 1$, then

$$\begin{aligned} &|\mathcal{Y}(\kappa, n, \delta, \omega_1, \omega_2)(\mathcal{Q})| \\ &\leq 2 \left(\frac{1}{3(\delta + \kappa)}\right)^{1-\frac{1}{q_2}} \left[\frac{1}{3(\delta + \kappa)} \right. \\ &\quad \times \max\{|\mathcal{Q}^{(\kappa)}(\omega_1)|^{q_2}, |\mathcal{Q}^{(\kappa)}(\omega_2)|^{q_2}\} \\ &\quad \left. - \frac{c(\mathcal{F}_{v_1, v_2}^\eta(\omega_2 - \omega_1))^p}{3(n + 1)^{2p}} \{2n(\delta + \kappa + 1)_2\mathfrak{F}_1 \right. \end{aligned}$$

6. CONCLUSION

Among the numerous definitions presented in convex analysis, it can be seen that the approximately generalized (ψ, \hbar) -convex, approximately generalized ψ -quasiconvex, HOS generalized (ψ, \hbar) -convex functions and HOS generalized ψ -quasiconvex functions are one of the main definitions of convex analysis that have been used successfully in optimization theory, coding theory, and machine learning. This is the most crucial reason for the need for new approaches in this field as a future direction. In the present scenario, we have established a new integral identity for κ th-order differentiable functions to deal with fractional operators involving the Riemann–Liouville fractional integral and special Raina’s function. New consequences and numerical comparisons demonstrate that the proposed technique is eligible to generate several existing outcomes in the relative literature by the suitable selection of parameters.^{31,32} The suggested scheme has concrete application in uniformly reflex Banach spaces and the parallelogram law in L^p spaces.^{26–28} For further investigation, taking into account the advanced convexity properties, in the preinvexity context, we may extend this study in inequality theory, quantum calculus, artificial intelligence, robotics and forecasting applications in different areas which are promising areas that need potential investigations. Considering these points and the demonstration of better performance with our proposed analysis, a bridge between theory and application can be established, which will eventually generate robust and optimal solutions.

REFERENCES

1. A. Atangana and D. Baleanu, New fractional derivatives with nonlocal and non-singular kernel: Theory and application to heat transfer model, *Therm. Sci.* **20**(2) (2016) 763–769.
2. J. E. Solis-Perez, J. F. Gomez-Aguilar and A. Atangana, Novel numerical method for solving variable-order fractional differential equations with power, exponential and Mittag–Leffler laws, *Chaos Solitons Fractals* **115** (2018) 175–185.
3. B. Ghanbari and A. Atangana, A new application of fractional Atangana–Baleanu derivatives: Designing ABC-fractional masks in image processing, *Physica A* **542** (2020), Article ID: 123516, 25 pp.
4. S. Kumar and A. Atangana, A numerical study of the nonlinear fractional mathematical model of tumor cells in presence of chemotherapeutic treatment, *Int. J. Biomath.* **13**(3) (2020) 2050021, doi.org/10.1142/S1793524520500217.
5. J. Danane, K. Allali and Z. Hammouch, Mathematical analysis of a fractional differential model of HBV infection with antibody immune response, *Chaos Solitons Fractals* **136** (2020) 109787, doi.org/10.1016/j.chaos.2020.109787.
6. T. Abdeljawad, S. Rashid, H. Khan and Y.-M. Chu, On new fractional integral inequalities for p -convexity within interval-valued functions, *Adv. Differ. Equ.* **2020** (2020) 330, doi.org/10.1186/s13662-020-02782-y.
7. H. Khan, A. Khan, T. Abdeljawad and A. Alkhazzan, Existence results in Banach space for a nonlinear impulsive system, *Adv. Differ. Equ.* **2019** (2019) 18.
8. T. Abdeljawad and D. Baleanu, Monotonicity results for fractional difference operators with discrete exponential kernels. *Adv. Differ. Equ.* **2017** (2017) 78.
9. H. Khan, T. Abdeljawad, C. Tunc, A. Alkhazzan and A. Khan, Minkowski’s inequality for the AB-fractional integral operator, *J. Inequal. Appl.* **2019** (2019) 96, doi.org/10.1186/s13660-019-2045-3.
10. M. Z. Sarikaya, E. Set, H. Yaldiz and N. Basak, Hermite–Hadamard’s inequalities for fractional integrals and related fractional inequalities, *Math. Comput. Model.* **57** (2013) 2403–2407.
11. E. Set, İ. İşcan, M. Z. Sarikaya and M. Özdemir, On new inequalities of Hermite–Hadamard–Fejer type for convex functions via fractional integrals, *Appl. Math. Comput.* **259** (2015) 875–888.
12. S. S. Dragomir, R. P. Agarwal and P. Cerone, On Simpson’s inequality and applications, *J. Inequal. Appl.* **5** (2000) 533–579.
13. S. Rashid, A. O. Akdemir, F. Jarad, M. A. Noor and K. I. Noor, Simpson’s type integral inequalities for k -fractional integrals and their applications, *AIMS Math.* **4**(4) (2019) 1087–1100.
14. Y. J. Li and T. S. Du, On Simpson type inequalities for functions whose derivatives are extended (s, m) -GA-convex functions, *Pure Appl. Math. China* **31** (2015) 487–497.
15. Z. Q. Yang, Y. J. Li and T. S. Du, A generalization of Simpson type inequality via differentiable functions using (s, m) -convex functions, *Italian J. Pure Appl. Math.* **35** (2015) 327–338.
16. Y. J. Li and T. S. Du, Some Simpson type integral inequalities for functions whose third derivatives are (s, m) -GA-convex functions, *J. Egypt. Math. Soc.* **24** (2016) 175–180.
17. S. Varosanec, On h -convexity. *J. Math. Anal. Appl.* **326** (2007) 26–35.

18. W. W. Breckner, Stetigkeitsaussagen für eine Klasse verallgemeinerter konvexer Funktionen in topologischen linearen Räumen, *Publ. Inst. Math.* **23** (1978) 13–20.
19. S. S. Dragomir, J. Pecaric and L. E. Persson, Some inequalities of Hadamard type, *Soochow J. Math.* **21** (1995) 335–341.
20. E. K. Godunova and V. I. Levin, Inequalities for functions of a broad class that contains convex, monotone and some other forms of functions, *Numer. Math. Math. Phys.* (1985) 138–142.
21. F. Chen and S. Wu, Several complementary inequalities to inequalities of Hermite–Hadamard type for s -convex functions, *J. Nonlinear Sci. Appl.* **9** (2016) 705–716.
22. S. S. Dragomir and R. P. Agarwal, Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula, *Appl. Math. Lett.* **11** (1998) 91–95.
23. B. T. Polyak, Existence theorems and convergence of minimizing sequences in extremum problems with restrictions, *Sov. Math. Dokl.* **7** (1966) 72–75.
24. S. Karamardian, The nonlinear complementarity problems with applications, Part 2, *J. Optim. Theory Appl.* **4** (1969) 167–181.
25. D. L. Zu and P. Marcotte, Co-coercivity and its role in the convergence of iterative schemes for solving variational inequalities, *SIAM J. Optim.* **6** (1996) 714–726.
26. W. L. Bynum, Weak parallelogram laws for Banach spaces, *Can. Math. Bull.* **19** (1976) 269–275.
27. R. Cheng and W. T. Ross, Weak parallelogram laws on Banach spaces and applications to prediction, *Period. Math. Hung.* **71** (2015) 45–58.
28. H.-K. Xu, Inequalities in Banach spaces with applications, *Nonlinear Anal. Theory Methods Appl.* **16** (1991) 1127–1138.
29. K. Nikodem and Z. Pales, Characterizations of inner product spaces by strongly convex functions, *Banach J. Math. Anal.* **5** (2011) 83–87.
30. R. Bassily, M. Belkin and S. Ma, On exponential convergence of SGD in non-convex over-parametrized learning, preprint (2018), arXiv:1811.02564.
31. S. Rashid, M. A. Latif, Z. Hammouch and Y. -M. Chu, Fractional integral inequalities for strongly h -preinvex functions for a k th order differentiable functions, *Symmetry* **11** (2019) 1448.
32. S. Rashid, H. Kalsoom, Z. Hammouch, R. Ashraf, D. Baleanu and Y.-M. Chu, New multi-parametrized estimates having p th-order differentiability in fractional calculus for predominating h -convex functions in Hilbert space, *Symmetry* **12** (2020) 222, doi:10.3390/sym12020222.
33. M. U. Awan, M. A. Noor, K. I. Noor and F. Safdar, On strongly generalized convex functions, *Filomat* **31** (2017) 5783–5790.
34. S. K. Mishra and N. Sharma, On strongly generalized convex functions of higher order, *Math. Inequal. Appl.* **22** (2019) 111–121.
35. B. B. Mohsen, M. A. Noor, K. I. Noor and M. Postolache, Strongly convex functions of higher order involving bifunction, *Mathematics* **7** (2019) 1028.
36. R. K. Raina, On generalized Wright’s hypergeometric functions and fractional calculus operators, *East Asian Math. J.* **21** (2015) 191–203.
37. M. J. Vivas–Cortez, A. Kashuri, J. E. H. Hernández, Trapezium-type inequalities for the Raina’s fractional integrals operator via generalized convex, *J. Math. Inequal.* (2019), in press.
38. S. Mubeen and G. M. Habibullah, On k -fractional integrals and application. *Int. J. Contemp. Math. Sci.* **7** (2012) 89–94.