

## Optical solitons for conformable space-time fractional nonlinear model



Muhammad Imran Asjad<sup>a,\*</sup>, Naeem Ullah<sup>a</sup>, Hamood ur Rehman<sup>b</sup>, Dumitru Baleanu<sup>c,d</sup>

<sup>a</sup>Department of Mathematics, University of Management and Technology, Lahore, Pakistan.

<sup>b</sup>Department of Mathematics, University of Okara, Okara, Pakistan.

<sup>c</sup>Department of Mathematics, Cankaya University, 06530 Balgat, Ankara, Turkey.

<sup>d</sup>Institute of Space Sciences, R76900 Magurele-Bucharest, Romania.

### Abstract

In search of the exact solutions of nonlinear partial differential equations in solitons form has become most popular to understand the internal features of physical phenomena. In this paper, we discovered various type of solitons solutions for the conformable space-time nonlinear Schrödinger equation (CSTNLSE) with Kerr law nonlinearity. To seek such solutions, we applied two proposed methods which are Sardar-subequation method and new extended hyperbolic function method. In this way several types of solitons obtained for example bright, dark, periodic singular, combined dark-bright, singular, and combined singular solitons. Some of the acquired solutions are interpreted graphically. These solutions are specific, novel, correct and may be beneficial for edifying precise nonlinear physical phenomena in nonlinear dynamical schemes. It is revealed that the proposed methods offer a straightforward and mathematical tool for solving nonlinear conformable space-time nonlinear Schrödinger equation. These results support in attaining nonlinear optical fibers in the future.

**Keywords:** Sardar-subequation method, conformable space-time nonlinear Schrödinger equation, the new extended hyperbolic function method, optical solitons.

**2020 MSC:** 35Q51, 35Q53.

©2022 All rights reserved.

### 1. Introduction

Study of optical solitons have definitely increased momentum in the arena of the solitary waves. Analysis of numerous solutions have been made to the nonlinear Schrödinger equations with low group velocity dispersion, Kerr nonlinearities, dispersion terms, self-steepening, spatiotemporal dispersion, etc. Usually, these solutions are qualifying chirped free and chirped solitons, combo solitons, and dark combo soliton [4, 9, 10, 31, 33]. Although these results have many applications but communication by optic fibers is one of them. Also, solitons have revolutionized the communication system through the wave guides more recently. It is clear that the solitons establish the pillar of data transfer and communication at unbelievable distances. However, all the power of the optical system lies on famous effects, which at the same time establish conditions restrictions. Maximum of the time, pulse propagation in optical fibers

\*Corresponding author

Email address: [imran.asjad@umt.edu.pk](mailto:imran.asjad@umt.edu.pk) (Muhammad Imran Asjad)

doi: [10.22436/jmcs.027.01.03](https://doi.org/10.22436/jmcs.027.01.03)

Received: 2021-09-08 Revised: 2021-11-07 Accepted: 2021-12-02

can be concerned by nonlinearity, group velocity dispersion (GVD), and polarization type dispersion. In recent times, fractional calculus looks in many regions of engineering and science such as rheology, biology, control theory, physics, electrochemistry, systems identification, signal processing, viscoelasticity and so on. In order to define nonlinear (NL) physical phenomena the acquiring the analytical solutions for fractional differential equation (FDE) is one of the significant features. This physical phenomenon may rest on both the time instant and the time history, which can be effectively demonstrated by using the concept of derivatives and integrals of fractional order. Recently, numerous approaches have been utilized to seek exact solutions of FPDEs in the literature (Zhou et al. [54]; Eslami et al. [18]; Mirzazadeh et al. [40]; Vajargah et al. [51]; Sonomezoglu et al. [47, 48]; Zerrad et al. [52, 53]; Biswas and Suarez [49]; Gazizov et al. [21]; Biswas et al. [12]; Lukashchuk [38]; Hashemi [22]; Hosseini and Ansari [25]; Baleanu et al. [6–8]; Hosseini et al. [26]; Inc et al. [34]; Akgül et al. [3]; Baleanu et al. [5]; Rehman et al. [43]; Inc et al. [32]; Tchier et al. [50]; Hashemi et al. [24]). It is famous that some of the belongings of the fractional derivatives are very tough in contrast with the typical ones, so there is a massive inspiration to crack in discovering the solutions of some general equations similar to the fractional nonlinear Schrodinger equation (NLSE). The NLSE is an essential model which defines many type of phenomena, such as optics of nonlinear media, condensed matter physics, plasmas and photonics, (Eslami et al. [15, 16]; Eslami and Rezazadeh [20]; Khodadad et al. [36]; Eslami [14, 17]; Ekici et al. [13]; Biswas et al. [11]; Mirzazadeha et al. [39]; Eslami and Neirameh [19]; Neirameh and Eslami [41]). Additionally, NLSE explains the dynamics behavior of solitons using optical fibers. The dimensionless NLSE with STD is given by

$$iu_t + au_{tx} + bu_{xx} + cF(|u|^2)u = 0, \quad (1.1)$$

where  $a$ ,  $b$ , and  $c$  denote the coefficients of STD, GVD, and NL term, respectively and

$$F(|u|^2)u \in \bigcup_{r,s=1}^{\infty} c^k \left( (-s, s) \times (-r, r); \mathbb{R}^2 \right). \quad (1.2)$$

The layouts of this paper are as follows. The governing equation is discussed in Section 2. Section 3 presents the analysis of the proposed methods SSM and new EHFM. In Section 4 SSM and the new EHFM are applied and Section 5 consists the results and discussion. In Section 6 conclusion of this paper is given.

## 2. Governing equation

In this section, we examine and analyze (1.1) given by Hashemi and Akgul [23] and

$$i \frac{\partial^\alpha u}{\partial t^\alpha} + a \frac{\partial^{\alpha+\beta} u}{\partial t^\alpha \partial x^\beta} + b \frac{\partial^{2\beta} u}{\partial x^{2\beta}} + cF(|u|^2)u = 0. \quad (2.1)$$

We apply two methods which are Sardar-subequation method (SSM) (Rezazadeh et al. [44]) and new extended hyperbolic function method (EHFM) (Shang [45, 46]; Nestor [42]) to discover optical solitons solutions for the CSTNLSE with Kerr law nonlinearity. It is appropriate that many models in engineering and science have an experiential parameters. Therefore, exact solutions provide authority to scientists to plan and perform experiments, by launching suitable or natural situations, to decide these parameters. Thus, exploration and finding solitons solutions is much effective in nonlinear sciences. The freshly familiarized definition of fractional derivative called conformable derivative (Hosseini et al. [27–30]; Korkmaz and Hosseini [37]; Khalil et al. [35]; Hammad and Khalil [2]; Abdeljawad [1]) is applied to transform fractional equations into ODEs, therefore discovering optical solitons solutions via two unlike analytical techniques.

## 3. Analysis of the methods

In this section, we have analyzed two methods which are applied to construct novel solitons solutions of the given model.

### 3.1. Sardar-subequation method

Let the FPDE

$$H(u, u_t^\alpha, u_x^\beta, u_x^{2\beta}, \dots) = 0, \quad (3.1)$$

to explore the travelling wave solutions, using following wave transformations in (1.1)

$$u(x, t) = u(\eta), \quad \eta = \frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha}, \quad (3.2)$$

Using (3.2) into (3.1), we obtain the following ODE

$$P(u, u', u'', \dots) = 0. \quad (3.3)$$

Consider (3.3) has a solution as follows

$$u(\eta) = \sum_{i=0}^N F_i \Phi^i(\eta),$$

where  $F_i$  ( $0 \leq i \leq N$ ) are constants and  $\Phi(\eta)$  admits the ODE as given below

$$(\Phi'(\eta))^2 = \epsilon + \delta \Phi^2(\eta) + \Phi^4(\eta), \quad (3.4)$$

where  $\epsilon$  and  $\delta$  are constants. (3.4) gives the solution as follows.

**Case 1:** When  $\delta > 0$  and  $\epsilon = 0$ , then

$$\Phi_1^\pm(\eta) = \pm \sqrt{-\delta pq} \operatorname{sech}_{pq}(\sqrt{\delta} \eta), \quad \Phi_2^\pm(\eta) = \pm \sqrt{\delta pq} \operatorname{csch}_{pq}(\sqrt{\delta} \eta),$$

where  $\operatorname{sech}_{pq}(\eta) = \frac{2}{pe^\eta + qe^{-\eta}}$ ,  $\operatorname{csch}_{pq}(\eta) = \frac{2}{pe^\eta - qe^{-\eta}}$ .

**Case 2:** When  $\delta < 0$  and  $\epsilon = 0$ , then

$$\Phi_3^\pm(\eta) = \pm \sqrt{-\delta pq} \operatorname{sec}_{pq}(\sqrt{-\delta} \eta), \quad \Phi_4^\pm(\eta) = \pm \sqrt{-\delta pq} \operatorname{csc}_{pq}(\sqrt{-\delta} \eta),$$

where  $\operatorname{sec}_{pq}(\eta) = \frac{2}{pe^\eta + qe^{-\eta}}$ ,  $\operatorname{csc}_{pq}(\eta) = \frac{2i}{pe^\eta - qe^{-\eta}}$ .

**Case 3:** When  $\delta < 0$  and  $\epsilon = \frac{\delta^2}{4b}$ , then

$$\begin{aligned} \Phi_5^\pm(\eta) &= \pm \sqrt{-\frac{\delta}{2}} \operatorname{tanh}_{pq}\left(\sqrt{-\frac{\delta}{2}} \eta\right), \\ \Phi_6^\pm(\eta) &= \pm \sqrt{-\frac{\delta}{2}} \operatorname{coth}_{pq}\left(\sqrt{-\frac{\delta}{2}} \eta\right), \\ \Phi_7^\pm(\eta) &= \pm \sqrt{-\frac{\delta}{2}} \left( \operatorname{tanh}_{pq}(\sqrt{-2\delta} \eta) \pm i\sqrt{pq} \operatorname{sech}_{pq}(\sqrt{-2\delta} \eta) \right), \\ \Phi_8^\pm(\eta) &= \pm \sqrt{-\frac{\delta}{2}} \left( \operatorname{coth}_{pq}(\sqrt{-2\delta} \eta) \pm \sqrt{pq} \operatorname{csch}_{pq}(\sqrt{-2\delta} \eta) \right), \\ \Phi_9^\pm(\eta) &= \pm \sqrt{-\frac{\delta}{8}} \left( \operatorname{tanh}_{pq}\left(\sqrt{-\frac{\delta}{8}} \eta\right) + \operatorname{coth}_{pq}\left(\sqrt{-\frac{\delta}{8}} \eta\right) \right), \end{aligned}$$

where  $\operatorname{tanh}_{pq}(\eta) = \frac{pe^\eta - qe^{-\eta}}{pe^\eta + qe^{-\eta}}$ ,  $\operatorname{coth}_{pq}(\eta) = \frac{pe^\eta + qe^{-\eta}}{pe^\eta - qe^{-\eta}}$ .

**Case 4:** When  $\delta > 0$  and  $\epsilon = \frac{\delta^2}{4}$ , then

$$\Phi_{10}^\pm(\eta) = \pm \sqrt{\frac{\delta}{2}} \operatorname{tan}_{pq}\left(\sqrt{\frac{\delta}{2}} \eta\right),$$

$$\begin{aligned}\Phi_{11}^{\pm}(\eta) &= \pm\sqrt{\frac{\delta}{2}} \cot_{pq}\left(\sqrt{\frac{\delta}{2}} \eta\right), \\ \Phi_{12}^{\pm}(\eta) &= \pm\sqrt{\frac{\delta}{2}} \left( \tan_{pq}(\sqrt{2\delta} \eta) \pm \sqrt{pq} \sec_{pq}(\sqrt{2\delta} \eta) \right), \\ \Phi_{13}^{\pm}(\eta) &= \pm\sqrt{\frac{\delta}{2}} \left( \cot_{pq}(\sqrt{2\delta} \eta) \pm \sqrt{pq} \csc_{pq}(\sqrt{2\delta} \eta) \right), \\ \Phi_{14}^{\pm}(\eta) &= \pm\sqrt{\frac{\delta}{8}} \left( \tan_{pq}\left(\sqrt{\frac{\delta}{8}} \eta\right) + \cot_{pq}\left(\sqrt{\frac{\delta}{8}} \eta\right) \right),\end{aligned}$$

where  $\tan_{pq}(\eta) = -\iota \frac{pe^{\iota\eta} - qe^{-\iota\eta}}{pe^{\iota\eta} + qe^{-\iota\eta}}$ ,  $\cot_{pq}(\eta) = \iota \frac{pe^{\iota\eta} + qe^{-\iota\eta}}{pe^{\iota\eta} - qe^{-\iota\eta}}$ .

### 3.2. New extended hyperbolic function method

The phases of the new EHF method are taken as follows.

**Form 1:** Let PDE as given in (3.1) with the wave transformation in (3.2) using wave transformation ODE is obtained as in (3.3). We assume that (3.3) has a solution in the next form

$$u(\eta) = \sum_{i=0}^N F_i \Phi^i(\eta), \quad (3.5)$$

where the coefficients  $F_i$  ( $i = (1, 2, 3, \dots, N)$ ) are constants and  $\Phi(\eta)$  admits the ODE in next form, as

$$\frac{d\Phi}{d\eta} = \Phi \sqrt{\Lambda + \Theta \Phi^2}, \quad \Lambda, \Theta \in \mathbb{R}. \quad (3.6)$$

By using balancing rule in (3.3) the value of  $N$  is found. Replacing (3.3) into (3.5) with (3.6), gives a set of equations for  $F_i$  ( $i = (0, 1, 2, 3, \dots, N)$ ). On solving this set, we yield set of solutions that admits (3.6), as follows.

**Set 1:** When  $\Lambda > 0$  and  $\Theta > 0$ ,

$$\Phi(\eta) = -\sqrt{\frac{\Lambda}{\Theta}} \operatorname{csch}(\sqrt{\Lambda}(\eta + \eta_0)).$$

**Set 2:** When  $\Lambda < 0$  and  $\Theta > 0$ ,

$$\Phi(\eta) = \sqrt{\frac{-\Lambda}{\Theta}} \sec(\sqrt{-\Lambda}(\eta + \eta_0)).$$

**Set 3:** When  $\Lambda > 0$  and  $\Theta < 0$ ,

$$\Phi(\eta) = \sqrt{\frac{\Lambda}{-\Theta}} \operatorname{sech}(\sqrt{\Lambda}(\eta + \eta_0)).$$

**Set 4:** When  $\Lambda < 0$  and  $\Theta > 0$ ,

$$\Phi(\eta) = \sqrt{\frac{-\Lambda}{\Theta}} \csc(\sqrt{-\Lambda}(\eta + \eta_0)).$$

**Set 5:** When  $\Lambda > 0$  and  $\Theta = 0$ ,

$$\Phi(\eta) = \exp(\sqrt{\Lambda}(\eta + \eta_0)).$$

**Set 6:** When  $\Lambda < 0$  and  $\Theta = 0$ ,

$$\Phi(\eta) = \cos(\sqrt{-\Lambda}(\eta + \eta_0)) + \iota \sin(\sqrt{-\Lambda}(\eta + \eta_0)).$$

**Set 7:** When  $\Lambda = 0$  and  $\Theta > 0$ ,

$$\Phi(\eta) = \pm \frac{1}{(\sqrt{\Theta}(\eta + \eta_0))}.$$

**Set 8:** When  $\Lambda = 0$  and  $\Theta < 0$ ,

$$\Phi(\eta) = \pm \frac{1}{(\sqrt{-\Theta}(\eta + \eta_0))}.$$

**Form 2:** Using the same pattern as previous, we adopt that  $\Phi(\eta)$  admits the ODE as follows

$$\frac{d\Phi}{d\eta} = \Lambda + \Theta\Phi^2, \quad \Lambda, \Theta \in \mathbb{R}. \tag{3.7}$$

Substituting (3.5) into (3.3) along with (3.7) with value of  $N$ , makes a set of (equations with the values of  $F_i (i = 1, 2, 3, \dots, N)$ ). Let the (3.7) accepts the solutions as follows

**Set 1:** When  $\Lambda\Theta > 0$ ,

$$\Phi(\eta) = \text{sgn}(\Lambda) \sqrt{\frac{\Lambda}{\Theta}} \tan(\sqrt{\Lambda\Theta}(\eta + \eta_0)).$$

**Set 2:** When  $\Lambda\Theta > 0$ ,

$$\Phi(\eta) = -\text{sgn}(\Lambda) \sqrt{\frac{\Lambda}{\Theta}} \cot(\sqrt{\Lambda\Theta}(\eta + \eta_0)).$$

**Set 3:** When  $\Lambda\Theta < 0$ ,

$$\Phi(\eta) = \text{sgn}(\Lambda) \sqrt{\frac{\Lambda}{-\Theta}} \tanh(\sqrt{-\Lambda\Theta}(\eta + \eta_0)).$$

**Set 4:** When  $\Lambda\Theta < 0$ ,

$$\Phi(\eta) = \text{sgn}(\Lambda) \sqrt{\frac{\Lambda}{-\Theta}} \coth(\sqrt{-\Lambda\Theta}(\eta + \eta_0)).$$

**Set 5:** When  $\Lambda = 0$  and  $\Theta > 0$ ,

$$\Phi(\eta) = -\frac{1}{\Theta(\eta + \eta_0)}.$$

**Set 6:** When  $\Lambda \in \mathbb{R}$  and  $\Theta = 0$ ,

$$\Phi(\eta) = \Lambda(\eta + \eta_0).$$

**Note:**  $\text{sgn}$  is the famous sign function.

#### 4. Application

In this section, we employ the under consideration methods to obtain optical solitons for the given model, for Kerr law nonlinearity, we have that  $F(u) = u$ . Suppose the next wave transformation

$$u(x, t) = u(\eta) e^{i \left( k \frac{x^\beta}{\beta} + w \frac{t^\alpha}{\alpha} \right)}, \quad \eta = \frac{x^\beta}{\beta} - V \frac{t^\alpha}{\alpha}, \tag{4.1}$$

substituting (4.1) into (2.1) and splitting into imaginary and real parts,

$$V = \frac{aw - 2bk}{1 - ak},$$

and

$$k^2(b - aV)u''(\eta) - (awk - w + bk^2)u(\eta) + cu^3 = 0. \tag{4.2}$$

4.1. Application of the SSM

Here, we employ the SSM for the solutions of NLSE. Using balancing principal on terms of  $u''$  and  $u^3$  in (4.2), we get  $N = 1$ , so (3.1) converts to

$$u(\eta) = F_0 + F_1\Phi(\eta), \tag{4.3}$$

where  $F_0$  and  $F_1$  are constants. Substituting (4.3) into (4.2) and comparing the coefficients of polynomials of  $\Phi(\eta)$  to zero, we make a set of equations in  $F_0, F_1, c$ , and  $a$ . On resolving the set of equations, we get

$$F_0 = 0, \quad F_1 = \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}}, \quad c \neq 0, \quad a = \frac{-bk^2 + \delta bk^2 + w}{k(w + kV\delta)}.$$

**Case 1:** When  $\delta > 0$  and  $\epsilon = 0$ , then

$$z_{1,1}^{\pm}(x, t) = \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}} \left( \pm \sqrt{-pq\delta} \operatorname{sech}_{pq}(\sqrt{\delta}(\eta)) \right) e^{i\left(k\frac{x^\beta}{\beta} + w\frac{t^\alpha}{\alpha}\right)}, \tag{4.4}$$

$$z_{1,2}^{\pm}(x, t) = \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}} \left( \pm \sqrt{pq\delta} \operatorname{csch}_{pq}(\sqrt{\delta}(\eta)) \right) e^{i\left(k\frac{x^\beta}{\beta} + w\frac{t^\alpha}{\alpha}\right)}.$$

**Case 2:** When  $\delta < 0$  and  $\epsilon = 0$ , then

$$z_{1,3}^{\pm}(x, t) = \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}} \left( \pm \sqrt{-pq\delta} \operatorname{sec}_{pq}(\sqrt{-\delta}(\eta)) \right) e^{i\left(\frac{x^\beta}{\beta} + \lambda\frac{t^\alpha}{\alpha}\right)},$$

$$z_{1,4}^{\pm}(x, t) = \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}} \left( \pm \sqrt{-pq\delta} \operatorname{csc}_{pq}(\sqrt{-\delta}(\eta)) \right) e^{i\left(\frac{x^\beta}{\beta} + \lambda\frac{t^\alpha}{\alpha}\right)}.$$

**Case 3:** When  $\delta < 0$  and  $\epsilon = \frac{\delta^2}{4b}$ , then

$$z_{1,5}^{\pm}(x, t) = \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}} \left( \pm \sqrt{-\frac{\delta}{2}} \operatorname{tanh}_{pq}\left(\sqrt{-\frac{\delta}{2}}(\eta)\right) \right) e^{i\left(\frac{x^\beta}{\beta} + \lambda\frac{t^\alpha}{\alpha}\right)}, \tag{4.5}$$

$$z_{1,6}^{\pm}(x, t) = \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}} \left( \pm \sqrt{-\frac{\delta}{2}} \operatorname{coth}_{pq}\left(\sqrt{-\frac{\delta}{2}}(\eta)\right) \right) e^{i\left(\frac{x^\beta}{\beta} + \lambda\frac{t^\alpha}{\alpha}\right)},$$

$$z_{1,7}^{\pm}(x, t) = \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}} \left( \pm \sqrt{-\frac{\delta}{2}} \left( \operatorname{tanh}_{pq}(\sqrt{-2\delta}(\eta)) \right. \right. \\ \left. \left. \pm i\sqrt{pq} \operatorname{sech}_{pq}(\sqrt{-2\delta}(\eta)) \right) \right) e^{i\left(\frac{x^\beta}{\beta} + \lambda\frac{t^\alpha}{\alpha}\right)}, \tag{4.6}$$

$$z_{1,8}^{\pm}(x, t) = \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}} \left( \pm \sqrt{-\frac{\delta}{2}} \left( \operatorname{coth}_{pq}(\sqrt{-2\delta}(\eta)) \right. \right. \\ \left. \left. \pm \sqrt{pq} \operatorname{csch}_{pq}(\sqrt{-2\delta}(\eta)) \right) \right) e^{i\left(\frac{x^\beta}{\beta} + \lambda\frac{t^\alpha}{\alpha}\right)}, \tag{4.7}$$

$$z_{1,9}^{\pm}(x, t) = \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}} \left( \pm \sqrt{-\frac{\delta}{8}} \left( \operatorname{tanh}_{pq}\left(\sqrt{-\frac{\delta}{8}}(\eta)\right) \right. \right. \\ \left. \left. + \operatorname{coth}_{pq}\left(\sqrt{-\frac{\delta}{8}}(\eta)\right) \right) \right) e^{i\left(\frac{x^\beta}{\beta} + \lambda\frac{t^\alpha}{\alpha}\right)}. \tag{4.8}$$

**Case 4:** When  $\delta > 0$  and  $\epsilon = \frac{\delta^2}{4}$ , then

$$\begin{aligned}
 z_{1,10}^{\pm}(x, t) &= \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}} \left( \pm \sqrt{\frac{\delta}{2}} \tan_{pq}\left(\sqrt{\frac{\delta}{2}}(\eta)\right) \right) e^{t\left(\frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha}\right)}, \\
 z_{1,11}^{\pm}(x, t) &= \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}} \left( \pm \sqrt{\frac{\delta}{2}} \cot_{pq}\left(\sqrt{\frac{\delta}{2}}(\eta)\right) \right) e^{t\left(\frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha}\right)}, \\
 z_{1,12}^{\pm}(x, t) &= \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}} \left( \pm \sqrt{\frac{\delta}{2}} \left( \tan_{pq}(\sqrt{2\delta}(\eta)) \pm \sqrt{pq} \sec_{pq}(\sqrt{2\delta}(\eta)) \right) \right) e^{t\left(\frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha}\right)}, \\
 z_{1,13}^{\pm}(x, t) &= \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}} \left( \pm \sqrt{\frac{\delta}{2}} \left( \cot_{pq}(\sqrt{2\delta}(\eta)) \pm \sqrt{pq} \csc_{pq}(\sqrt{2\delta}(\eta)) \right) \right) e^{t\left(\frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha}\right)}, \\
 z_{1,14}^{\pm}(x, t) &= \frac{k\sqrt{-2b + \frac{2V(-bk^2+w+bk^2\delta)}{k(w+kV\delta)}}}{\sqrt{c}} \left( \pm \sqrt{\frac{\delta}{8}} \left( \tan_{pq}\left(\sqrt{\frac{\delta}{8}}(\eta)\right) \right. \right. \\
 &\quad \left. \left. + \cot_{pq}\left(\sqrt{\frac{\delta}{8}}(\eta)\right) \right) \right) e^{t\left(\frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha}\right)}. \tag{4.9}
 \end{aligned}$$

#### 4.2. Application of the new EHFm

**Form 1:** Here, we employ the new EHFm for the solutions NLSE. Using balancing method in (4.2), we get  $N = 1$ , so (3.5) gives

$$u(\eta) = F_0 + F_1\Phi(\eta), \tag{4.10}$$

where  $F_0$  and  $F_1$  are constants. Replacing (3.7) into (4.2) and equating the coefficients polynomials of  $\Phi(\eta)$  to zero, we get a set of equations in  $F_0$ ,  $F_1$ ,  $\Lambda$  and  $\Theta$ . On resolving the set of equations, we attain

$$\begin{aligned}
 F_0 = 0, \quad F_1 &= \frac{\sqrt{2}\sqrt{-b\Theta k^2 + a\Theta k^2 V}}{\sqrt{c}}, \quad c \neq 0, \\
 V \neq 0, \quad \Lambda &= \frac{bk^2 - w + akw}{\Theta k^2(b - aV)}, \quad \Theta = \Theta. \tag{4.11}
 \end{aligned}$$

**Set 1:** When  $\Lambda > 0$  and  $\Theta > 0$ ,

$$z_1(x, t) = \frac{\sqrt{2}\sqrt{-b\Theta k^2 + a\Theta k^2 V}}{\sqrt{c}} \left( -\sqrt{\frac{bk^2 - w + akw}{\Theta^2 k^2(b - aV)}} \operatorname{csch}\left(\sqrt{\frac{bk^2 - w + akw}{\Theta k^2(b - aV)}}(\eta + \eta_0)\right) \right) e^{t\left(\frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha}\right)}. \tag{4.12}$$

**Set 2:** When  $\Lambda < 0$  and  $\Theta > 0$ ,

$$z_2(x, t) = \frac{\sqrt{2}\sqrt{-b\Theta k^2 + a\Theta k^2 V}}{\sqrt{c}} \left( \sqrt{-\frac{bk^2 - w + akw}{\Theta^2 k^2(b - aV)}} \operatorname{sec}\left(\sqrt{-\frac{bk^2 - w + akw}{\Theta k^2(b - aV)}}(\eta + \eta_0)\right) \right) e^{t\left(\frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha}\right)}.$$

**Set 3:** When  $\Lambda > 0$  and  $\Theta < 0$ ,

$$z_3(x, t) = \frac{\sqrt{2}\sqrt{-b\Theta k^2 + a\Theta k^2 V}}{\sqrt{c}} \left( \sqrt{\frac{bk^2 - w + akw}{\Theta^2 k^2(b - aV)}} \operatorname{sech}\left(\sqrt{\frac{bk^2 - w + akw}{\Theta k^2(b - aV)}}(\eta + \eta_0)\right) \right) e^{t\left(\frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha}\right)}. \tag{4.13}$$

**Set 4:** When  $\Lambda < 0$  and  $\Theta < 0$ ,

$$z_4(x, t) = \frac{\sqrt{2}\sqrt{-b\Theta k^2 + a\Theta k^2 V}}{\sqrt{c}} \left( \sqrt{-\frac{bk^2 - w + akw}{\Theta^2 k^2(b - aV)}} \operatorname{csc}\left(\sqrt{-\frac{bk^2 - w + akw}{\Theta k^2(b - aV)}}(\eta + \eta_0)\right) \right) e^{t\left(\frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha}\right)}.$$

**Set 5:** When  $\Lambda = 0$  and  $\Theta > 0$ ,

$$z_7(x, t) = \frac{\sqrt{2\sqrt{-b\Theta k^2 + a\Theta k^2 V}}}{\sqrt{c}} \left( \pm \frac{1}{(\sqrt{\Theta}(\eta + \eta_0))} \right) e^{\iota \left( \frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha} \right)}.$$

**Set 6:** When  $\Lambda = 0$  and  $\Theta < 0$ ,

$$z_8(x, t) = \frac{\sqrt{2\sqrt{-b\Theta k^2 + a\Theta k^2 V}}}{\sqrt{c}} \left( \pm \frac{\iota}{(\sqrt{-\Theta}(\eta + \eta_0))} \right) e^{\iota \left( \frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha} \right)},$$

where  $\eta = \frac{x^\beta}{\beta} - V \frac{t^\alpha}{\alpha}$ .

**Form 2:** Operating balancing rule in (4.2), gives  $N = 1$ , so (3.5) reduces to

$$u(\eta) = F_0 + F_1 \Phi(\eta), \quad (4.14)$$

where  $F_0$  and  $F_1$  are constants. Replacing (4.14) into (4.2) and equating the coefficients of polynomials of  $\Phi(\eta)$  to zero, we get a set of equations in  $F_0$ ,  $F_1$ ,  $\Lambda$ , and  $\Theta$ . On resolving the set of equations, we attain

$$\begin{aligned} F_0 = 0, \quad F_1 &= \frac{\Theta k \sqrt{-2b + 2aV}}{\sqrt{c}}, \quad c \neq 0, \\ V \neq 0, \quad \Lambda &= \frac{bk^2 - w + akw}{2\Theta k^2(b - aV)}, \quad \Theta = \Theta. \end{aligned} \quad (4.15)$$

**Set 1:** When  $\Lambda \Theta > 0$ ,

$$z_9(x, t) = \frac{\Theta k \sqrt{-2b + 2aV}}{\sqrt{c}} \left( \chi \sqrt{\frac{bk^2 - w + akw}{2\Theta^2 k^2(b - aV)}} \tan \left( \sqrt{\frac{bk^2 - w + akw}{2k^2(b - aV)}} (\eta + \eta_0) \right) \right) e^{\iota \left( \frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha} \right)}. \quad (4.16)$$

**Set 2:** When  $\Lambda \Theta > 0$ ,

$$z_{10}(x, t) = \frac{\Theta k \sqrt{-2b + 2aV}}{\sqrt{c}} \left( -\chi \sqrt{\frac{bk^2 - w + akw}{2\Theta^2 k^2(b - aV)}} \cot \left( \sqrt{\frac{bk^2 - w + akw}{2k^2(b - aV)}} (\eta + \eta_0) \right) \right) e^{\iota \left( \frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha} \right)}. \quad (4.17)$$

**Set 3:** When  $\Lambda \Theta < 0$ ,

$$z_{11}(x, t) = \frac{\Theta k \sqrt{-2b + 2aV}}{\sqrt{c}} \left( \chi \sqrt{\frac{bk^2 - w + akw}{2\Theta^2 k^2(b - aV)}} \tanh \left( \sqrt{\frac{bk^2 - w + akw}{2k^2(b - aV)}} (\eta + \eta_0) \right) \right) e^{\iota \left( \frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha} \right)}.$$

**Set 4:** When  $\Lambda \Theta < 0$ ,

$$z_{12}(x, t) = \frac{\Theta k \sqrt{-2b + 2aV}}{\sqrt{c}} \left( \chi \sqrt{\frac{bk^2 - w + akw}{2\Theta^2 k^2(b - aV)}} \coth \left( \sqrt{\frac{bk^2 - w + akw}{2k^2(b - aV)}} (\eta + \eta_0) \right) \right) e^{\iota \left( \frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha} \right)}.$$

**Set 5:** When  $\Lambda = 0$  and  $\Theta > 0$ ,

$$z_{13}(x, t) = \frac{\Theta k \sqrt{-2b + 2aV}}{\sqrt{c}} \left( -\frac{1}{\Theta(\eta + \eta_0)} \right) e^{\iota \left( \frac{x^\beta}{\beta} + \lambda \frac{t^\alpha}{\alpha} \right)}.$$

where  $\chi = \text{sgn} \left( \frac{bk^2 - w + akw}{2\Theta k^2(b - aV)} \right)$ ,  $\eta = \frac{x^\beta}{\beta} - V \frac{t^\alpha}{\alpha}$ .



## 5. Results and discussions

In this study, we successfully construct novel solitons solutions along with hyperbolic and trigonometric function solutions for the CSTNLSE with Kerr law of nonlinearity using SSM and new EHFm. These methods are considered as most recent schemes in this arena and that are not utilized to this equation earlier. For physical analysis, 3-dim and 2-dim plots of some of these solutions are included with appropriate parameters. These acquired solutions discover their application in communication to convey information because solitons have the capability to spread over long distances without reduction and without changing their forms.

We only added particular figures to avoid overfilling the document. Moreover, we effectively construct various solitary wave and solitons solutions along with hyperbolic and trigonometric function solutions. Absolutely the developed results are fresh and different from that reported results. These acquired solutions discover their application in communication to convey information because solitons have the capability to spread over long distances without reduction and without changing their forms. The authors proposed different analytic approach in newly issued article and reported some fascinating findings. We can understand from all the graphs that the SSM and new EHFm are very effectual and more specific in assessing the equation under consideration. In this paper, we only added particular figures to avoid overfilling the document. The authors proposed different analytic approaches in newly issued article and reported some fascinating findings. We can understand from all the graphs that the SSM and new EHFm are very effectual and more specific in assessing the equation under consideration. For graphical representation for (1.1), the physical behavior of (4.4) using the proper values of parameters  $F_2 = 2.45$ ,  $\alpha = 1.75$ ,  $\nu = 0.65$ ,  $p = 1$ ,  $q = 1$ ,  $\alpha = 0.75$ ,  $k = 0.8$ , and  $t = 1$  are shown in Fig. 1, the physical behavior of (4.5) using the appropriate values of parameters  $F_2 = 2.45$ ,  $\alpha = -1.75$ ,  $\nu = 0.65$ ,  $p = 1$ ,  $q = 1$ ,  $\alpha = 0.75$ ,  $k = 0.8$ , and  $t = 1$  are shown in Fig. 2, the physical behavior of (4.9) using the proper values of parameters  $F_2 = 2.45$ ,  $\alpha = 1.75$ ,  $\nu = 0.65$ ,  $p = 1$ ,  $q = 1$ ,  $\alpha = 0.75$ ,  $k = 0.8$ , and  $t = 1$  are shown in Fig. 3, the entire behavior of (4.12) using the proper values of parameters  $F_2 = 2.45$ ,  $\alpha = 1.75$ ,  $\nu = 0.65$ ,  $p = 1$ ,  $q = 1$ ,  $\alpha = 0.75$ ,  $k = 0.8$ , and  $t = 1$  are shown in Fig. 4, the complete behavior of (4.13) using the proper values of parameters  $F_2 = 1.4$ ,  $\alpha = 0.4$ ,  $\nu = 0.5$ ,  $\Theta = 0.75$ ,  $k = 0.65$ ,  $\beta = 0.2$ ,  $\gamma = 0.8$ , and  $t = 1$  are revealed in Fig. 5, the complete behavior of (4.16) with the proper values of parameters  $F_2 = 1.4$ ,  $\alpha = -0.4$ ,  $\nu = 0.5$ ,  $\Theta = -0.75$ ,  $k = 0.65$ ,  $\beta = 0.2$ ,  $\gamma = 0.8$ , and  $t = 1$  are presented in Fig. 6, the absolute behavior of (4.17) using the proper values of parameters  $F_2 = 2.45$ ,  $\sigma = 0.95$ ,  $\nu = 0.5$ ,  $\Theta = -0.85$ ,  $k = 0.45$ , and  $t = 1$  are revealed in Fig. 7.




Figure 1: (a) 3D graph of (4.6) with  $\alpha = 0.45$ ,  $\beta = 0.65$ ,  $w = 2$ ,  $V = 2$ ,  $p = 0.98$ ,  $q = 0.95$ ,  $k = 2$ ,  $A = 3$ ,  $b = 2$ ,  $\delta = 2$ ,  $c = 4$ ; (a-1) 2D plot of (4.4) with  $t = 1$ ; (a-2) Contour graph of (4.4).




Figure 2: (b) 3D graph of (4.10) with  $\alpha = 0.45$ ,  $\beta = 0.65$ ,  $w = 2$ ,  $p = 0.98$ ,  $q = 0.95$ ,  $k = 2$ ,  $\Lambda = 2.7$ ,  $b = 2$ ,  $\delta = 2$ ,  $c = 4$ ,  $V = 2$ ; (b-1) 2D plot of (4.5) with  $t = 1$ ; (b-2) Contour graph of (4.5).




Figure 3: (c) 3D graph of (4.19) with  $\alpha = 0.45$ ,  $\beta = 0.65$ ,  $w = 2$ ,  $p = 0.98$ ,  $q = 0.95$ ,  $k = 2$ ,  $\Lambda = 2.7$ ,  $b = 2$ ,  $\delta = 2$ ,  $c = 4$ ,  $V = 2$ ; (c-1) 2D plot of (4.9) with  $t = 1$ ; (c-2) Contour graph of (4.9).




Figure 4: (d) 3D graph of (4.22) with  $b = 2.5$ ,  $a = 1.5$ ,  $v = 1$ ,  $k = 2$ ,  $p = 0.98$ ,  $q = 0.95$ ,  $\alpha = 0.75$ ,  $\Lambda = 2.7$ ,  $\alpha = 0.45$ ,  $\beta = 0.65$ ,  $\theta = 4$ ,  $V = 3$ ,  $w = 3$ ; (d-1) 2D plot of (4.12) with  $t = 1$ ; (d-2) Contour graph of (4.12).




Figure 5: (e) 3D graph of (4.24) with  $b = 2.5$ ,  $a = 1.5$ ,  $v = 1$ ,  $k = 2$ ,  $p = 0.98$ ,  $q = 0.95$ ,  $\alpha = 0.75$ ,  $A = 2.7$ ,  $\alpha = 0.45$ ,  $\beta = 0.65$ ,  $\theta = 4$ ,  $V = 3$ ,  $w = 3$ ; (e-1) 2D plot of (4.13) with  $t = 1$ ; (e-2) Contour graph of (4.13).




Figure 6: (f) 3D graph of (4.30) with  $b = 2.5$ ,  $a = 1.5$ ,  $v = 1$ ,  $k = 2$ ,  $p = 0.98$ ,  $q = 0.95$ ,  $\alpha = 0.75$ ,  $A = 2.7$ ,  $\alpha = 0.45$ ,  $\beta = 0.65$ ,  $\theta = 4$ ,  $V = 3$ ,  $w = 3$ ; (f-1) 2D plot of (4.16) with  $t = 1$ ; (f-2) Contour graph of (4.16).




Figure 7: (g) 3D graph of (4.31) with  $b = 2.5$ ,  $a = 1.5$ ,  $v = 1$ ,  $k = 2$ ,  $p = 0.98$ ,  $q = 0.95$ ,  $\alpha = 0.75$ ,  $A = 2.7$ ,  $\alpha = 0.45$ ,  $\beta = 0.65$ ,  $\theta = 4$ ,  $V = 3$ ,  $w = 3$ ; (g-1) 2D plot of (4.17) with  $t = 1$ ; (g-2) Contour graph of (4.17).

## 6. Conclusion

We constructed novel optical soliton solutions for the CSTNLSE with Kerr law nonlinearity by using SSM and new EHF. Different cases of the constraints are used to describe specific solutions. The main

aim of this study is to acquire optical solitons that could satisfy the constraint conditions posed on the different parameters of the CSTNLSE. In order to describe the behavior of acquired solutions of the model, we plotted some selected solutions by giving appropriate values to the involved parameters. The obtained results may have much influence on numerous fields of nonlinear sciences. By obtained results, we can recognize that the proposed methods are proficient, consistent and beneficial for discovering the exact solutions of nonlinear FPDEs in a wide range. It has been detected that all the solutions of governing model in this study are novel and unique. In future, we will be more interested in birefringence aspect and cross-phase modulation to construct ultrashort optical pulses and soliton pulses. We need to investigate some new type optical solitons and modulation instability analysis of some fractional order NLSE type equations in the future.

## Acknowledgment

The authors would like to grateful to acknowledge the University of Management and Technology Lahore, Pakistan for the financial support for this research.

## References

- [1] T. Abdeljawad, *On conformable fractional calculus*, J. Comput. Appl. Math., **279** (2015), 57–66. 2
- [2] M. Abu Hammad, R. Khalil, *Conformable fractional heat differential equation*, Int. J. Pure Appl. Math., **94** (2014), 215–221. 2
- [3] A. Akgül, *A novel method for a fractional derivative with non-local and non-singular kernel*, Chaos Soliton Fractal, **114** (2018), 478–482. 1
- [4] M. Arshad, A. R. Seadawy, D. Lu, *Bright-dark solitary wave solutions of generalized higher-order nonlinear Schrödinger equation and its applications in optics*, J Electromagnet Waves Appl., **31** (2017), 1711–1721. 1
- [5] D. Baleanu, A. Fernandez, A. Akgül, *On a fractional operator combining proportional and classical differintegrals*, Mathematics, **8** (2020), 13 pages. 1
- [6] D. Baleanu, M. Inc, A. Yusuf, A. I. Aliyu, *Lie symmetry analysis, exact solutions and conservation laws for the time fractional modified Zakharov-Kuznetsov equation*, Nonlinear Anal. Model. Control, **22** (2017), 861–876. 1
- [7] D. Baleanu, M. Inc, A. Yusuf, A. I. Aliyu, *Lie symmetry analysis, exact solutions and conservation laws for the time fractional Caudrey-Dodd-Gibbon-Sawada-Kotera equation*, Commun. Nonlinear Sci. Numer. Simulat., **59** (2018), 222–234.
- [8] D. Baleanu, M. Inc, A. Yusuf, A. I. Aliyu, *Time fractional third-order evolution equation: symmetry analysis, explicit solutions, and conservation laws*, J. Comput. Nonlinear Dyn., **13** (2018), 021011-5. 1
- [9] A. Biswas, *Soliton solutions of the perturbed resonant nonlinear Schrödinger equation with full nonlinearity by semi-inverse variational principle*, Quantum Phys. Lett. **1** (2012), 79–89. 1
- [10] A. Biswas, M. O. AlAmr, H. Rezazadeh, M. Mirzazadeh, M. Eslami, Q. Zhou, *Resonant optical solitons with dual-power law nonlinearity and fractional temporal evolution*, Optik, **165** (2018), 233–239. 1
- [11] A. Biswas, M. Mirzazadeh, M. Eslami, D. Milovic, *Solitons in optical metamaterials by functional variable method and first integral approach*, Frequenz, **68** (2014), 525–530. 1
- [12] A. Biswas, R. Morris, A. H. Kara, *Soliton solution and conservation laws of the Zakharov-Kuznetsov equation in plasmas with power law nonlinearity*, Nonlinear Anal. Model. Control, **18** (2013), 153–159. 1
- [13] M. Ekici, M. Mirzazadeh, M. Eslami, *Solitons and other solutions to Boussinesq equation with power law nonlinearity and dual dispersion*, Nonlinear Dyn., **84** (2016), 669–676. 1
- [14] M. Ekici, M. Mirzazadeh, M. Eslami, Q. Zhou, S. P. Moshokoa, A. Biswas, M. Belic, *Optical soliton perturbation with fractional-temporal evolution by first integral method with conformable fractional derivatives*, Optik, **127** (2016), 10659–10669. 1
- [15] M. Eslami, *Exact traveling wave solutions to the fractional coupled nonlinear Schrodinger equations*, Appl. Math. Comput., **285** (2016), 141–148. 1
- [16] M. Eslami, *Trial solution technique to chiral nonlinear Schrodinger's equation in (1 + 2)-dimensions*, Nonlinear Dyn., **85** (2016), 813–816. 1
- [17] M. Eslami, F. S. Khodadad, F. Nazari, H. Rezazadeh, *The first integral method applied to the Bogoyavlenskii equations by means of conformable fractional derivative*, Opt. Quant. Electron., **49** (2017), 18 pages. 1
- [18] M. Eslami, M. Mirzazadeh, *Optical solitons with Biswas-Milovic equation for power law and dual-power law nonlinearities*, Nonlinear Dyn., **83** (2016), 731–738. 1
- [19] M. Eslami, A. Neirameh, *New exact solutions for higher order nonlinear Schrodinger equation in optical fibers*, Opt. Quant. Electron., **50** (2017), 8 pages. 1

- [20] M. Eslami, H. Rezazadeh, *The first integral method for Wu-Zhang system with conformable time-fractional derivative*, *Calcolo*, **53** (2016), 475–485. 1
- [21] R. K. Gazizov, N. H. Ibragimov, S. Y. Lukashchuk, *Nonlinear self-adjointness, conservation laws and exact solutions of time-fractional Kompaneets equations*, *Commun. Nonlinear Sci. Numer. Simul.*, **23** (2014), 153–163. 1
- [22] M. S. Hashemi, *Group analysis and exact solutions of the time fractional Fokker-Planck equation*, *Phys. A Stat. Mech. Appl.*, **417** (2015), 141–149. 1
- [23] M. S. Hashemi, A. Akgül, *Solitary wave solutions of time-space nonlinear fractional Schrödinger's equation: Two analytical approaches*, *J. Comput. Appl. Math.*, **339** (2017), 147–160. 2
- [24] M. S. Hashemi, M. Inc, B. Kilic, A. Akgül, *On solitons and invariant solutions of the Magneto-electro-elastic circular rod*, *Waves in Random and Complex Media*, **26** (2016), 259–271. 1
- [25] K. Hosseini, R. Ansari, *New exact solutions of nonlinear conformable time-fractional Boussinesq equations using the modified Kudryashov method*, *Waves Random Complex Media*, **27** (2017), 628–636. 1
- [26] K. Hosseini, A. Bekir, R. Ansari, *New exact solutions of the conformable time-fractional Cahn-Allen and Cahn-Hilliard equations using the modified Kudryashov method*, *Optik*, **132** (2017), 203–209. 1
- [27] K. Hosseini, A. Bekir, M. Kaplan, Ö. Güner, *A new technique for solving the nonlinear conformable time-fractional differential equations*, *Opt. Quant. Electron.*, **49** (2017), 12 pages. 2
- [28] K. Hosseini, P. Mayeli, R. Ansari, *Bright and singular soliton solutions of the conformable time-fractional Klein-Gordon equations with different nonlinearities*, *Waves Random Complex Media*, **28** (2017), 426–434.
- [29] K. Hosseini, P. Mayeli, A. Bekir, O. Guner, *Density-dependent conformable space-time fractional diffusion-reaction equation and its exact solutions*, *Commun. Theor. Phys.*, **69** (2018), 4 pages.
- [30] K. Hosseini, Y. J. Xu, P. Mayeli, A. Bekir, P. Yao, Q. Zhou, O. Guner, *A study on the conformable time-fractional Klein-Gordon equations with quadratic and cubic nonlinearities*, *Optoelectron. Adv. Mater. Rapid Commun.*, **11** (2017), 423–429. 2
- [31] A. Houwe, M. B. Hubert, S. Nestor, D. Jerome, M. Justin, G. Betchewe, S. Y. Doka, K. T. Crepin, S. Khan, A. Biswas, M. Ekici, S. Adesanya, S. P. Moshokoa, M. Belic, *Optical solitons for higher-order nonlinear Schrödinger equation with three exotic integration architectures*, *Optik*, **17** (2019), 861–866. 1
- [32] M. Inc, A. Akgül, *Classifications of Soliton Solutions of the Generalized Benjamin-Bona-Mahony Equation with Power-Law Nonlinearity*, *J. Adv. Phys.*, **7** (2018), 130–134. 1
- [33] M. Inc, I. A. Aliyu, A. Yusuf, D. Baleanu, *Dispersive optical solitons and modulation instability analysis of Schrödinger-Hirota equation with spatio-temporal dispersion and Kerr law nonlinearity*, *Superlattices Microstruct.*, **113** (2018), 319–327. 1
- [34] M. Inc, A. Yusuf, A. I. Aliyu, D. Baleanu, *Time-fractional Cahn-Allen and time-fractional Klein-Gordon equations: lie symmetry analysis, explicit solutions and convergence analysis*, *Phys. A*, **493** (2018), 94–106. 1
- [35] R. Khalil, A. L. M. Horani, A. Yousef, M. Sababheh, *A new definition of fractional derivative*, *J. Comput. Appl. Math.*, **264** (2014), 65–70. 2
- [36] F. S. Khodadad, F. Nazari, M. Eslami, H. Rezazadeh, *Soliton solutions of the conformable fractional Zakharov-Kuznetsov equation with dual-power law nonlinearity*, *Opt. Quant. Electron.*, **49** (2017), 384. 1
- [37] A. Korkmaz, K. Hosseini, *Exact solutions of a nonlinear conformable time-fractional parabolic equation with exponential nonlinearity using reliable method*, *Opt. Quant. Electron.*, **49** (2017), 10 pages. 2
- [38] S. Y. Lukashchuk, *Conservation laws for time-fractional sub-diffusion and diffusion-wave equations*, *Nonlinear Dyn.*, **80** (2015), 791–802. 1
- [39] M. Mirzazadeh, M. Eslami, A. Biswas, *1-Soliton solution of KdV6 equation*, *Nonlinear Dyn.* **80** (2015), 387–396. 1
- [40] M. Mirzazadeh, M. Eslami, B. F. Vajargah, A. Biswas, *Application of the first integral method to fractional partial differential equations*, *Indian J. Phys.*, **88** (2014), 177–184. 1
- [41] A. Neirameh, M. Eslami, *An analytical method for finding exact solitary wave solutions of the coupled (2 + 1)-dimensional Painleve Burgers equation*, *Sci. Iran.*, **24** (2017), 715–726. 1
- [42] S. Nestor, A. Houwe, G. Betchewe, M. Inc, S. Y. Doka, *A series of abundant new optical solitons to the conformable space-time fractional perturbed nonlinear Schrödinger equation*, *Physica Scripta*, **95** (2020), 11 pages. 2
- [43] H. U. Rehman, M. A. Imran, M. Bibi, A. Akgül, *New soliton solutions of the 2D-chiral nonlinear Schrödinger equation using two integration schemes* *Math. Methods Appl. Sci.*, **44** (2021), 5663–5682. 1
- [44] H. Rezazadeh, M. Inc, D. Baleanu, *New Solitary Wave Solutions For Variants Of (3 + 1)-Dimensional Wazwaz-Benjamin-Bona-Mahony Equations*, *Frontiers Phys.*, **8** (2020), 12 pages. 2
- [45] Y. Shang, *The extended hyperbolic function method and exact solutions of the long-short wave resonance equations*, *Chaos Solitons Fractals*, **36** (2008), 762–771. 2
- [46] Y. Shang, Y. Huang, W. Yuan, *The extended hyperbolic functions method and new exact solutions to the Zakharov equations*, *Appl. Math. Comput.*, **200** (2008), 110–122. 2
- [47] A. Sonomezoglu, M. Eslami, Q. Zhou, E. Zerrad, A. Biswas, M. Belic, M. Mirzazadeh, M. Ekici, *Optical solitons in nano-fibers with fractional temporal evolution*, *J. Comput. Theor. Nanosci.*, **13** (2016), 5361–5374. 1
- [48] A. Sonomezoglu, S. Ortakaya, M. Eslami, A. Biswas, M. Mirzazadeh, M. Ekici, *Solitons solutions to a few fractional nonlinear evolution equations in shallow water wave dynamics*, *Eur. Phys. J. Plus*, **131** (2016), 166–177. 1
- [49] P. Suarez, A. Biswas, *Exact 1-soliton solution of the Zakharov equation in plasmas with power law nonlinearity*, *Appl. Math. Comput.*, **217** (2011), 7372–7375. 1

- [50] F. Tchier, M. Inc, B. Kilic, A. Akgül, *On soliton structures of generalized resonance equation with time dependent coefficients*, *Optik*, **128** (2017), 218–223. 1
- [51] B. F. Vajargah, M. Mirzazadeh, M. Eslami, A. Biswas, *Solitons and periodic solutions to a couple of fractional nonlinear evolution equations*, *Pramana*, **82** (2014), 465–476. 1
- [52] E. Zerrad, A. Biswas, R. Kohl, D. Milovic, *Optical solitons by He’s variational principle in a non-Kerr law media*, *J. Infrared Millim. Terahertz Waves*, **30** (2009), 526–537. 1
- [53] E. Zerrad, A. Biswas, M. Song, B. Ahmed, *Domain wall and bifurcation analysis of the klein-gordon Zakharov-Kuznetsov equation in  $(1 + 2)$ -dimensions with power law nonlinearity*, *Chaos*, **23** (2013), 15 pages. 1
- [54] Q. Zhou, S. P. Moshokoa, A. Biswas, M. Belic, M. Ekici, M. Mirzazadeh, *Solitons in optical metamaterials with fractional temporal evolution*, *Optik*, **127** (2016), 10879–10897. 1